



# **Initial Assessment of Facial Nerve Paralysis based on Motion Analysis using Optical Flow Method**

by

**WAN SYAHIRAH BINTI W.SAMSUDIN  
(1140610618)**

A thesis submitted in fulfillment of the requirements for the degree of  
Master of Science in Mechatronic Engineering

**School of Mechatronic Engineering  
UNIVERSITI MALAYSIA PERLIS**

2015

## **TABLE OF CONTENTS**

	<b>PAGE</b>
<b>THESIS DECLARATION</b>	ii
<b>ACKNOWLEDGEMENT</b>	iii
<b>TABLE OF CONTENTS</b>	iv
<b>LIST OF TABLES</b>	vii
<b>LIST OF FIGURES</b>	x
<b>LIST OF ABBREVIATION</b>	xi
<b>ABSTRAK</b>	xii
<b>ABSTRACT</b>	xiii
<b>CHAPTER 1 INTRODUCTION</b>	
1.1 Introduction	1
1.2 Problem Statement	4
1.3 Objectives of Research	5
1.4 Scope of Research	6
1.5 Thesis Outline	7
<b>CHAPTER 2 LITERATURE REVIEW</b>	
2.1 Introduction	8
2.2 Facial Nerve	8
2.3 Facial Paralysis	9
2.3.1 Incidence and Prevalence	10
2.3.2 Causes	11
2.3.3 Symptoms	12
2.3.4 Effects	13
2.4 Facial Nerve Assessment Methods	14
2.4.1 Subjective assessment methods	14
2.4.2 Objective assessment methods	19
2.5 Previous Kanade-Lucas-Tomasi (KLT) Research	27

2.6	Summary	29
-----	---------	----

## CHAPTER 3 METHODOLOGY

3.1	Introduction	34
3.2	Ethical Statement	34
3.3	Equipment Installation and Experimental Setup	35
3.4	Landmarks Chosen and Facial Exercise Development Based On Muscle Movement and House-Brackmann (HB) Score	36
3.5	Data Acquisition	40
3.5.1	Database	40
3.5.2	Subjects	41
3.5.3	Protocols of Experiment	42
3.5.4	Extracting Frames from Video Data	43
3.6	Facial Tracking System	44
3.6.1	Kanade-Lucas-Tomasi (KLT) Tracking Algorithm	45
3.6.2	Facial Tracking System using Optical Flow Method	48
3.6.3	Detect and track the landmarks chosen on facial video data	50
3.7	Compute motion changes using Kanade-Lucas-Tomasi (KLT) method	52
3.8	Data Analysis	55
3.9	Summary	59

## CHAPTER 4 RESULTS AND DISCUSSION

4.1	Introduction	60
4.2	Experiment Overview	61
4.2.1	Results Evaluation	61
4.3	Experimental Results	62
4.3.1	Experiment 1: Distance Measurement of Three Exercises in Four Conditions	62
4.3.2	Experiment 2: Area Measurement of Three Exercises in Four Conditions	82
4.4	Summary	99

<b>CHAPTER 5 CONCLUSIONS</b>	
5.1 Conclusion	101
5.2 Contribution of Research	102
5.3 Research Limitations	103
5.4 Suggestions for Future Work	103
<b>REFERENCES</b>	105
<b>APPENDIX</b>	111
<b>LIST OF PUBLICATIONS</b>	113

## LIST OF TABLES

NO		PAGE
2.1	House-Brackmann Grading Scale	16
2.2	Sunnybrook Scale	17
2.3	Yanagihara Grading System	18
2.4	Peitersen Grading System	19
2.5	Nottingham Grading System	22
2.6	Summarization of Subjective Assessment Methods	31
2.7	Summarization of Objective Assessment Methods	31
3.1	List of Selected Landmarks for Distance Measurement	38
3.2	List of Selected Landmarks for Area Measurement	39
3.3	Details on Subjects for Database	41
3.4	Tracking of Landmarks in Measuring Distance	51
3.5	Tracking of Landmarks in Measuring Area	52
3.6	Individual Score Chart	56
3.7	Total Score for Three Exercises	58
3.8	Grading paralysis based on Total Score	58
4.1	Distance in the Rise Forehead Exercise in Four Cases	63
4.2	Percentage and Individual Score in the Rise Forehead Exercise in Four Cases of Distance Measurement	65
4.3	Distance in the Smile Exercise in Four Cases	66
4.4	Percentage and Individual Score in the Smile Exercises in Four Cases of Distance Measurement	67
4.5	Distance in the Close Eyes Exercise in Four Cases	68
4.6	Percentage and Individual Score in the Close Eyes Exercise in Four Cases of Distance Measurement	70
4.7	Initial Facial Assessment Table for Normal Subjects in the First Case of Distance Measurement	72

4.8	Initial Facial Assessment Table for Patient Subjects in the First Case of Distance Measurement	73
4.9	Initial Facial Assessment Table for Normal Subjects in the Second Case of Distance Measurement	74
4.10	Initial Facial Assessment Table for Patient Subjects in the Second Case of Distance Measurement	75
4.11	Initial Facial Assessment Table for Normal Subjects in the Third Case of Distance Measurement	76
4.12	Initial Facial Assessment Table for Patient Subjects in the Third Case of Distance Measurement	77
4.13	Initial Facial Assessment Table for Normal Subjects in the Fourth Condition of Distance Measurement	78
4.14	Initial Facial Assessment Table for Patient Subjects in the Fourth Case of Distance Measurement	79
4.15	Accuracy Table for Four Cases of Distance Measurement	80
4.16	Area in the Rise Forehead Exercise in Four Cases	82
4.17	Percentage and Individual Score in the Rise Forehead Exercise in the Four Cases of Area Measurement	84
4.18	Area in the Smile Exercise in Four Cases	85
4.19	Percentage and Individual Score in the Smile Exercises in Four Cases of Area Measurement	86
4.20	Area in the Close Eyes Exercise in Four Cases	87
4.21	Percentage and Individual Score in the Close Eyes Exercises in Four Cases of Area Measurement	89
4.22	Initial Facial Assessment Table for Normal Subjects in the First Case of Area Measurement	90
4.23	Initial Facial Assessment Table for Patient Subjects in the First Case of Area Measurement	91
4.24	Initial Facial Assessment Table for Normal Subjects in the Second Case of Area Measurement	92
4.25	Initial Facial Assessment Table for Patient Subjects in the Second Case of Area Measurement	93

4.26	Initial Facial Assessment Table for Normal Subjects of Third Case of Area Measurement	94
4.27	Initial Facial Assessment Table for Patient Subjects in the Third Case of Area Measurement	95
4.28	Initial Facial Assessment Table for Normal Subjects in the Fourth Case of Area Measurement	96
4.29	Initial Facial Assessment Table for Patient Subjects in the Fourth Case of Area Measurement	97
4.30	Accuracy Table for Four Cases of Area Measurement	98

## **LIST OF FIGURES**

<b>NO</b>		<b>PAGE</b>
1.1	Head and facial muscles	1
2.1	Anatomy of facial nerve	9
2.2	The measurement of Nottingham Grading System	21
3.1	Flowchart of Methodology	35
3.2	Flowchart of Experiment Protocol for Data Acquisition	42
3.2	(A) and (B) are two rectangle features; (C) Three-rectangle features; (D) Four-rectangle features	35
3.3	Extracted Frames from Video Data of Subject	44
3.4	Image Registration Problem	45
3.6	Illustration of distance measurement in Rise Forehead Exercise	53
3.7	Illustration of area measurement in Rise Forehead Exercise	54



## **LIST OF ABBREVIATIONS**

AFA	Automated Facial Analysis
AU	Action Units
AVI	Audio Video Interleave
FNF	Facial Nerve Paralysis
HB	House Brackmann
HBGS	House Brackmann Grading System
KLT	Kanade-Lucas-Tomasi
LBP	Local Binary Pattern
MREC	Medical Research and Ethics Committee
SRA	Maximum Static Response Array
OP	Optical Flow
RAD	Resistor-Average-Distance
RBF	Radial Basis Function
SLFS	Stennert-Limberg-Frentrup Scale

# **Penilaian Awal Kelumpuhan Saraf Muka Berdasarkan Analisis Pergerakan Menggunakan Kaedah Aliran Optik**

## **ABSTRAK**

Pesakit yang mengalami lumpuh saraf muka menghadapi masalah fungsi, kosmetik, dan psikologi yang serius dengan keupayaan untuk berkomunikasi secara lisan dan bukan lisan terjejas. Rehabilitasi untuk kelumpuhan muka bermula dengan penilaian klinikal yang teliti dalam menilai tahap lumpuh. Secara umumnya, untuk penilaian ketidakfungsian saraf muka pada muka pesakit dalam aplikasi klinikal setiap hari, system pemerhatian yang subjektif oleh pakar klinikal telah digunakan. Penilaian subjektif terhadap fungsi saraf muka ini dicapai dengan membuat kesimpulan melalui pemerhatian kepada pergerakan muka yang tidak dipaksa. Kaedah-kaedah ini biasanya menghasilkan satu nilai dimana hipotesisnya sepadan dengan tahap kelumpuhan muka. Setakat ini, tiada kaedah penilaian secara subjektif yang boleh menghasilkan maklumat kuantitatif mengenai fungsi saraf muka dengan tepat. Kajian sebelum ini telah mencadangkan pelbagai kaedah bagi memperolehi data objektif untuk mengukur dan menentukan tahap kelumpuhan muka. Walau bagaimanapun, tiada kaedah kuantitatif yang mengikut piawaian bagi penerangan fungsi saraf muka secara objektif. Mengakui kepentingan penilaian muka ini, kajian ini dijalankan untuk membangunkan penilaian awal yang boleh mengelaskan subjek kepada normal dan pesakit berdasarkan analisis pergerakan muka menggunakan algoritma aliran optik serta mengkategorikan pesakit kepada enam peringkat mengikut sistem House-Brackmann (HB). Sebelum penilaian awal dibangunkan, dua eksperimen telah dijalankan untuk mencari parameter yang terbaik dan juga ukuran terbaik untuk menilai kelumpuhan muka. Jarak dan luas kawasan telah dipilih sebagai dua parameter yang disiasat dalam kajian ini. Beberapa analisis matematik dan statistic telah dijalankan untuk menentukan ukuran yang terbaik daripada parameter-parameter ini. Keputusan menunjukkan bahawa jarak adalah parameter yang terbaik dan nilai pada bingkai awal senaman adalah pengukuran yang paling penting dalam mengesan perubahan pergerakan muka. Tesis ini juga membentangkan penilaian awal muka yang mengandungi markah individu untuk setiap senaman yang terlibat dalam kajian ini dan juga menggredkan tahap lumpuh untuk setiap pesakit sepadan dengan sistem HB. Kajian ini telah mendapat keputusan yang memuaskan yang telah disahkan oleh pakar perubatan dalam Otorinolaringologi. Penilaian awal muka ini memainkan peranan penting dalam permulaan system penilaian akhir untuk kelumpuhan muka dan membangunkan program rehabilitasi yang lebih baik untuk pesakit.

# **Initial Assessment of Facial Nerve Paralysis Based on Motion Analysis using Optical Flow Method**

## **ABSTRACT**

Patients with facial nerve paralysis suffer serious functional, cosmetic and psychological problems with impaired ability to communicate both verbally and non-verbally. Rehabilitation for facial paralysis begins with a thorough clinical evaluation in accessing the degree of paralysis. Generally, for assessment of facial nerve dysfunction on patient's face in daily clinical application, observation-based subjective grading systems by clinicians is been employed. Subjective assessments of facial nerve function are accomplished by inferring nerve function through the observation of voluntary facial movement. These methods usually yield a single value which hypothetically corresponds to the severity of facial paralysis. To date, there are no subjective assessment methods that can reliably produce quantitative information regarding facial nerve function. Previous works have proposed a wide range of methods to obtain objective data to quantify and determine the severity of facial paralysis. However, there is no standardized quantitative method for objective description of facial nerve function. Acknowledging the importance of this facial assessment, this research was conducted to develop an initial assessment method that can classify normal and patient subjects based on facial motion analysis using Optical Flow algorithm as well as categorizing the severity of patient into six levels according to House-Brackmann (HB) system. Prior to initial assessment method been developed, two experiments were conducted to find the best parameter and best measurement for assessing facial paralysis. Distance and area were selected as two measurement parameters which were investigated in this research. A number of mathematical and statistical analyses were performed to determine the best measurement of these parameters. The results indicate that the distance is the best parameter and the value of initial exercise frame is the most important measurement in tracking the changes of facial movement. This thesis also presents the initial facial assessment method which contains the individual scores for each exercise involved in this research and also grading the paralysis for each patient corresponds to HB system. This research showed satisfactory results which was validated by a medical professional in Otorhinolaryngology. This initial facial assessment method may play a pivotal role in initiation of final assessment system for facial paralysis and develop better rehabilitation program for patients.

# CHAPTER 1

## INTRODUCTION

### 1.1 Introduction

Facial nerve incorporates about 7,000 individual nerve fibers. Each fiber brings the electrical impulses to a specific facial muscle. All information through the fibers of this nerve permits us to make any facial expression such as laugh, cry, smile, and frown. Human express the emotions such as sadness, surprise, happiness, excitement and confusion through facial expressions. For example, a smile can be interpreted as the sign of happiness while frowning shows some disagreement and sadness. Facial nerve paralysis will take place when some of the individual nerve fibers are disrupted. Besides, the movement of facial muscles starts spasming or twitching if these fibers are irritated. Figure 1.1 shows some of the facial muscles that are responsible for the facial expressions.




Figure 1.1: Head and facial muscles (Knysh B., 2015).

Facial nerve paralysis (FNP) is a common problem which involves the paralysis of any structure innervated by the facial nerve. Literature shows that the prevalent cause of this paralysis is Bell's palsy, named after Sir Charles Bell (1774-1842). Bell's palsy is an idiopathic disease category, reporting approximately 50% of the cases (Brach & VanSwearingen, 1999; Singhi & Jain, 2003).

The other possible causes of facial paralysis are due to birth, trauma, neurologic syndromes, infection, metabolic, neoplastic, toxic and iatrogenic (Benecke, 2002). Patients with these paralyses suffer serious functional, cosmetic and psychological problems with impaired ability to communicate both verbally and non-verbally. Numbness can occur on the affected side of the face although no actual sensory loss occurs (Singhi, 2003). Besides, they will be unable to close the eye on the affected side, which can lead to irritation and corneal ulceration. Because of that, the eye should be lubricated with artificial tears until facial paralysis ends (Piercy, 2005). However, the most dramatic impact of the paralysis is its psychological effect where the patients may have low confidence and fear when interacting with others.

A thorough medical history of patients and physical examination are the earliest steps in making a facial diagnosis. Clinicians examine whether the forehead is involved in motor defect or not. This is commonly accomplished by assessing how well a patient can raise his or her eyebrows. The results from this action help in determining which part the lesion is in whether in the upper motor neuron or lower motor neuron of facial nerve component. Rehabilitation is suggested by doctors after patient's treatment, depending on their condition. It is to regain the function of facial nerves and improve both the strength and flexibility of the nerves. The failure in rehabilitation procedures may lead to the continued weakness and inability to function of facial nerves.

The availability of facial rehabilitation is limited, and most individuals with facial movement disorders have been told to await (spontaneous) recovery or told no effective intervention exists. Consequently, individuals with this paralysis will deal with physical, psychological, and social disability daily. The rehabilitation for facial paralysis begins with a thorough clinical evaluation in accessing the degree of paralysis. It is important to measure the facial disability from onset to the various stages of recovery and also to detect changes over time or after treatment. In the past few decades, several internationally accepted systems have been proposed by different researchers, yet most of the existing systems are subjective. This subjective evaluation refers to various facial nerve grading systems, of which the most widely used is the House-Brackmann (HB) system (Dellanoy& Ward, 2010; Dulguerov, Wang, Perneger, Marchal, & Lehmann, 2003). The facial nerve evaluation will vary over many clinicians and it will results in inaccurate assessment of paralysis.

There is no standardized system yet for facial nerve evaluation which has been accepted for world-wide used. Thus, an objective standardized method, which is easy to perform, low cost and fast, and simple can be a useful clinical tool to detect the level of paralysis in patients with facial palsy and monitor their improvement or performance during and after the rehabilitation procedures. In this study, an initial facial assessment method is proposed to assist clinicians in assessing the facial nerve function. Comparisons have been made between two measurement parameters to determine which one is better to be implemented in the assessment system.

## **1.2 Problem Statement**

The problems that motivate this study are the limitation of standard facial nerve assessment, subjective clinical assessment by clinicians or physiotherapist, and the requirement of printed scores during evaluation of facial nerve function.

### **A. Limitation of Standard Facial Nerve Assessment**

There are many grading systems such as House-Brackmann (HB), Yanagihara and Sunnybrook system and also assessment methods available to evaluate the facial nerve function. However, no standardized system is used by clinicians because the previous systems have not achieved the requirement to be an ideal system for assessing facial paralysis. Mostly, all the assessment methods are based on the HB system. So, an initial assessment method which can give an objective measurement for both sides of the face, simple, easy to use, and based on the HB system is an ideal assessment method to evaluate the facial nerve function.

### **B. Subjective Clinical Assessment by Clinicians**

No objective assessment of patient's nerve functions in real life application is clinically available. The results vary depending on the clinicians' observation and the evaluation is not standardized (Meier-Gallati, Scriba, & Fisch, 1998). Sometimes, the person who is still in a paralyzed state is treated as normal because the evaluation is only based on the clinicians' naked eyes. An objective assessment which provides quantitative measurement can reduce the time for grading the paralysis and make it reliable to all the patients, even with different clinicians or physiotherapists'

assessment (Murty, Diver, Kelly, & O'Donoghue, 1994). Even though the patient is still in the same level of HB scores, the clinicians are able to see how much their patients have improved in performance of rehabilitation based on the objective measurement.

### **C. Requirement of Printed Scores during Evaluation of Facial Nerve Function**

The clinicians and doctors have to refer to the printed scores on sheets of paper and make their subjective evaluation based on the patients' movements (Isono, Murata, Tanaka, Kawamoto, & Azuma, 1996; He Soraghan, O'Reilly, & Xing, 2009). Apart from referring to the scores, some of them need to compare the snapshots taken during the rehabilitation program to evaluate the performance of patients. The initial objective assessment will assist the clinicians in detecting the level of paralysis and record the performance of patients. It will help clinicians to acknowledge the patients' improvement during rehabilitation program.

### **1.3 Objectives of Research**

The main objective of this project is to develop an initial facial assessment for facial nerve paralysis based on motion analysis using Optical Flow (OF) method. The sub-objectives of this research include:

1. To explore and identify all possible facial muscles which are involved in the desired rehabilitation facial exercises.
2. To determine potential landmarks on the face based on muscle movement and medical professional's recommendation



3. To track the landmarks over the time (in the image frames) successfully by applying optical flow algorithm.
4. To investigate the most suitable measurement parameter based on motion analysis to be implemented in the initial assessment of facial nerve function.
5. To validate the performance of the initial facial assessment system, which is able to differentiate between normal and patient subjects, detect the level of paralysis with the HB system and compare the results with a clinician's evaluation.

#### **1.4 Scope of Research**

The scope of this research is to develop an initial facial nerve assessment to evaluate the facial nerve function. This research will analyze two parameters which are possible to evaluate the facial nerve function, which is distance and area. The proposed assessment method will be able to differentiate between normal and patient subjects, determine which side of face that is paralyzed and also able to predict the level of paralysis based on the HB scoring system. The data for this research are normal data (healthy subjects) and patient data (Bell's palsy patients from Hospital Tunku Fauziah (HTF)). Bell's palsy is the common cause of facial paralysis which is reported in HTF. This assessment method will be validated by appropriate medical professionals from HTF.

## **1.5 Thesis Outline**

In Chapter 1, the general idea of this research which is to develop an initial facial nerve assessment for facial paralysis has been briefly presented. The problem statement, research's objectives and scope are explained in this chapter. Theoretical studies and relevant literature are presented in Chapter 2. The physiology study on facial nerve and facial paralysis are explained thoroughly in this chapter. Previous works done on facial assessment method for facial paralysis is also discussed in this chapter. Chapter 3 focuses on the research methodology which is to develop an initial facial assessment for facial paralysis. The process of data acquisition, landmark placement, tracking the landmarks on the face, and the scores of the facial nerve function based on the exercise done by the subjects are discussed in detailed in this chapter. Results and discussion will be presented in Chapter 4. The results from the experiment are analyzed and explained in this chapter. Validation of the results by a medical professional from the Otorhinolaryngology department of HTF is also presented. Overall conclusion is summarized in Chapter 5. Contributions of the research are highlighted along with the challenges experienced during the research period. Finally, suggestions for future work are given at the end of this chapter.

## CHAPTER 2

### LITERATURE REVIEW

#### 2.1 Introduction

This chapter presents the overview of facial nerve and the type of facial nerve paralysis. The incidence and prevalence, causes, symptoms, and effects to patients who have facial nerve paralysis are discussed in this chapter. The literature on facial nerve assessment methods is also discussed.

#### 2.2 Facial Nerve

The facial nerve consists of a motor and also a sensory part, which being frequently described as the nervus intermedius (*pars intermedii of Wrisberg*). Anatomy of the facial nerve is shown in Figure 2.1. The nerve contains 10,000 fibers out of which 7,000 are myelinated motor fibers (Feghali, Joseph, Fayad Y., & Murphy M., 2013). Most of the motor fibers travel to the extratemporal portion of the facial nerve and innervate the muscle of the face, scalp, and auricle, the buccinators and platysma, the stapedius, the stylohyoideus, and posterior belly of the digastricus (Kecskes, 2012).




Figure 2.1: Anatomy of facial nerve (Tiemstra & Khatkhate, 2007).

The main function of the facial nerve is to express the voluntary behavior and spontaneous emotions via innervating 23 facial muscles on each side of the face. The facial muscles are inserted directly into the skin and the contraction of these muscles will cause the skin to move. Signals from the complex array of the nerves to the various muscles instruct the muscles to move in combinations and also as individually (Kecskes, 2012).

### 2.3 Facial Paralysis

Facial paralysis is the loss or impairment of motor function of facial muscles due to damage of the facial nerve (Cranial nerve VII), brainstem nuclei of the facial nerve, and/or the neuromuscular system innervated by this nerve. The degree of paralysis is from minor weakness to complete paralysis depends on the site and extent of the lesion (Dunwald, 2010).

Facial paralysis or facial nerve palsy is a serious problem, for both the afflicted patient and the physician attempting to make a proper diagnosis (Benecke,

2002). Facial paralysis is a disability of communication because as a human, the non-verbal communication is relied on our facial expression which reveals our innermost feelings. Those who work with this facial paralysis patients will aware for the need of rehabilitation in both the physiological and psychosocial aspects of this disability. Restoring the function and expression to the highest level possible will results in improved health, self-esteem, self-acceptance, acceptance by others, and quality of life (Tiemstra & Khatkhate, 2007).

### **2.3.1 Incidence and Prevalence**

The annual prevalence of facial paralysis is approximately between 15 to 40 cases per 100,000 in a general population (Kanerva, Poussa, & Pitkaranta, 2006; Meier-Gallati, Scriba, & Fisch, 1998; Ross, Fradet, & Nedzelski, 1996) of which 30% of them will developed a facial nerve paresis with sequelae (asymmetry of the face at rest and during movement, problems with speaking, eating and drinking and psychosocial problems) varying from very mild to very severe (Beurskens, 2004; Peitersen, 1994). Among 400 Dutch physiotherapists (response rate 76%) in 1996, 25% of them were involved in the treatment of patients with facial nerve paresis (Beurskens, 2004).

Both genders are included in this approximation with the peak ages is between 30 to 50 and 60 to 70 years old (Scriba, Stoeckli, Veraguth, Pollak, & Fisch, 1999). By referring to the US Census Bureau in their International Database, extrapolation on incidence rate of Bell's palsy has been made. Results show that China has the highest incidence rate, which is 191,007 cases followed by India and USA, which have 156,628 and 43,184 respectively.

### 2.3.2 Causes

Before a patient of facial paralysis undergone the rehabilitation process, the etiology and duration of the paralysis is recorded by clinicians. Some possible causes of facial paralysis are (Benecke, 2002; Kecskes, 2012; Schaitkin & May, 2000).

- i) Birth (molding, forceps, Möbius)
- ii) Trauma (skull fractures, penetrating injuries, facial injuries)
- iii) Neurologic (multiple sclerosis, Guillain-Barre syndrome)
- iv) Infection (otitis, mastoiditis, mumps, herpes virus, HIV, Epstein-Barr virus, Lyme disease)
- v) Metabolic (diabetes mellitus, thyroid disease, pregnancy)
- vi) Neoplastic (schwannoma, paraganglioma, meningioma, carcinoma, leukemia, cholesteatoma)
- vii) Toxic (thalidomide, carbon monoxide, tetanus, diphtheria)
- viii) Iatrogenic (anesthesia, vaccine, parotid surgery, otology surgery, dental, embolic)
- ix) Idiopathic (Bell's palsy, Melkersson Rosenthal paralysis, autoimmune disease, Guillain-Barre syndrome, multiple sclerosis, sarcoid, osteopetrosis)

The prevalent cause of this paralysis is Bell's palsy which is about 50% of the cases reported (Beurskens, 2004; Brach & VanSwearingen, 1999). However, it is important for clinicians to consider all potential causes to avoid overlook of any threatening diseases (Bleicher, Hamiel, & Genglon, 1996; Brach & VanSwearingen, 1996). Although at least 80 known causes of facial paralysis exist, the cause of Bell's palsy remains unknown. Reviews of the literature suggest that no population is immune to idiopathic facial paralysis like Bell's palsy.

### 2.3.3 Symptoms

Issues resulting from facial paralysis include those associated with: (a) somatic function, (b) social function, (c) psychological wellbeing, and (d) physiological function (Devriese, 1998). Physical symptoms of facial paralysis in the upper face include the inability to close the eyes and sagging lower eyelid. Tears may run from the affected eye, or alternatively, tearing may be reduced. Failure of the eyelids to completely close, combined with the loss of the blink reflex may cause irritation, exposure keratopathy, corneal ulcerations, and blindness (Tate & Tollefson, 2006).

The sagging of the mouth corner is the most notable characteristic in the lower half of the face. The lack of control over lip closure can result difficulties in eating, drinking, speech (especially with sounds that require the lips such as /p/ and /b/) and control of drooling. Also, the nostril on the affected side may collapse, causing nasal obstruction (Schaitkin & May, 2000). Additionally, patients with facial paralysis associated with inflammation, as in the case with Bell's palsy and herpes zoster oticus, may have pain in the mastoid process, neck, ear, or the face (Schaitkin & May, 2000). Patients will also develop synkinesis which is one sequelae of facial paralysis (Nakamura, Toda, Sakamaki, Kashima, & Takeda, 2003). Common examples of synkinesis include eye closure with volitional contraction of mouth muscles and involuntary movement of the mouth during eye closure. These abnormal movements in muscle contractions have massive cosmetic and functional implications.

#### **2.3.4 Effects**

One of the most socially devastating effects of facial paralysis is the inability to produce facial expression of emotion. Patients commonly report personal and work-related problems as well as limited social integration and interpersonal communication. These problems stem from facial disfigurement and difficulties in eating, drinking and communicating effectively in social settings. Also, people with facial paralysis are often introvert and may become isolated (Goldberg, DeLorie, Zuker, & Mantkelow, 2003).

The face also is a crucial component of beauty, sexual attractiveness, and sexual interest (Ekman, 1986). Therefore, patients with facial paralysis experience pronounce psychological distress. Oftentimes, they must cope with feelings of shame, decreased self-esteem, anxiety, depression, guilt, anger, and/or fear (Devriese, 1998; Ekman, 1986).

Nutritional impairment also may be apparent when facial paralysis exists. Routine and seemingly easy tasks such as eating can be quite challenging for patients with facial paralysis, who often demonstrate swallowing difficulties that occur in both oral and pharyngeal phases of deglutition. More specifically, a survey conducted by Sjogreen, Andersson-Norinder, & Jacobson (2001) revealed that this group has difficulties in the oral phase including getting food off spoon with lips, food and liquid leak out of the corners of the mouth and takes a long time to swallow bites of food. Choking on food and coughing when receiving liquid are some of the problems in the pharyngeal phase of the swallow.

Verbal communication also may be impaired in this population of patients. Bilabial incompetence produces characteristics speech error patterns including substitution, distortions, and omissions of bilabial sounds (/p/, /b/, /m/) and the



## CHAPTER 1

### INTRODUCTION

#### 1.1 Introduction

Facial nerve incorporates about 7,000 individual nerve fibers. Each fiber brings the electrical impulses to a specific facial muscle. All information through the fibers of this nerve permits us to make any facial expression such as laugh, cry, smile, and frown. Human express the emotions such as sadness, surprise, happiness, excitement and confusion through facial expressions. For example, a smile can be interpreted as the sign of happiness while frowning shows some disagreement and sadness. Facial nerve paralysis will take place when some of the individual nerve fibers are disrupted. Besides, the movement of facial muscles starts spasming or twitching if these fibers are irritated. Figure 1.1 shows some of the facial muscles that are responsible for the facial expressions.




Figure 1.1: Head and facial muscles (Knysh B., 2015).

Facial nerve paralysis (FNP) is a common problem which involves the paralysis of any structure innervated by the facial nerve. Literature shows that the prevalent cause of this paralysis is Bell's palsy, named after Sir Charles Bell (1774-1842). Bell's palsy is an idiopathic disease category, reporting approximately 50% of the cases (Brach & VanSwearingen, 1999; Singhi & Jain, 2003).

The other possible causes of facial paralysis are due to birth, trauma, neurologic syndromes, infection, metabolic, neoplastic, toxic and iatrogenic (Benecke, 2002). Patients with these paralyses suffer serious functional, cosmetic and psychological problems with impaired ability to communicate both verbally and non-verbally. Numbness can occur on the affected side of the face although no actual sensory loss occurs (Singhi, 2003). Besides, they will be unable to close the eye on the affected side, which can lead to irritation and corneal ulceration. Because of that, the eye should be lubricated with artificial tears until facial paralysis ends (Piercy, 2005). However, the most dramatic impact of the paralysis is its psychological effect where the patients may have low confidence and fear when interacting with others.

A thorough medical history of patients and physical examination are the earliest steps in making a facial diagnosis. Clinicians examine whether the forehead is involved in motor defect or not. This is commonly accomplished by assessing how well a patient can raise his or her eyebrows. The results from this action help in determining which part the lesion is in whether in the upper motor neuron or lower motor neuron of facial nerve component. Rehabilitation is suggested by doctors after patient's treatment, depending on their condition. It is to regain the function of facial nerves and improve both the strength and flexibility of the nerves. The failure in rehabilitation procedures may lead to the continued weakness and inability to function of facial nerves.

The availability of facial rehabilitation is limited, and most individuals with facial movement disorders have been told to await (spontaneous) recovery or told no effective intervention exists. Consequently, individuals with this paralysis will deal with physical, psychological, and social disability daily. The rehabilitation for facial paralysis begins with a thorough clinical evaluation in accessing the degree of paralysis. It is important to measure the facial disability from onset to the various stages of recovery and also to detect changes over time or after treatment. In the past few decades, several internationally accepted systems have been proposed by different researchers, yet most of the existing systems are subjective. This subjective evaluation refers to various facial nerve grading systems, of which the most widely used is the House-Brackmann (HB) system (Dellanoy& Ward, 2010; Dulguerov, Wang, Perneger, Marchal, & Lehmann, 2003). The facial nerve evaluation will vary over many clinicians and it will results in inaccurate assessment of paralysis.

There is no standardized system yet for facial nerve evaluation which has been accepted for world-wide used. Thus, an objective standardized method, which is easy to perform, low cost and fast, and simple can be a useful clinical tool to detect the level of paralysis in patients with facial palsy and monitor their improvement or performance during and after the rehabilitation procedures. In this study, an initial facial assessment method is proposed to assist clinicians in assessing the facial nerve function. Comparisons have been made between two measurement parameters to determine which one is better to be implemented in the assessment system.

## **CHAPTER 3**

### **METHODOLOGY**

#### **3.1 Introduction**

In this chapter, the methodology for the development of an initial assessment method for facial nerve paralysis will be discussed. The goal of this study is to develop an assessment method which is capable in grading the severity of the paralysis. For the facial tracking process, the Kanade-Lucas-Tomasi (KLT) optical flow algorithm was applied to track the motion of desired landmarks on the face over time. No markers were attached on the subject's face to aid tracking. A few experiments were conducted to select the best measurement parameter for the possible implementation towards facial nerve assessment. The flow of the methodology of this research is presented in Figure 3.1.

#### **3.2 Ethical Statement**

All the procedures in this study were approved by the Medical Research and Ethics Committee (MREC), Ministry of Health, Bangsar, Kuala Lumpur, Malaysia (Ref no.: KKM/NIHSEC/800-2/2/2/P13-96) (Appendix). All subjects were provided with the Patient Information Sheet and Patient Consent Form before the data acquisition process begun. The Patient Information Sheet is the document containing information about the intended research for prospective subjects whereas the Patient Consent Form is a form to document the subject's consent to participate in this study.




Figure 3.1: Flowchart of Methodology

### 3.3 Equipment Installation and Experimental Setup

For the experimental setup, no standard is available for collecting the video databases. So, protocol for the experiment is made based on the requirement for the research purpose. For collecting the facial exercises data, a digital camera (SONY DSC-W620) with  $640 \times 480$  resolution was used to capture the videos of the subject's face from the frontal view at the center of the face. The digital camera was placed in front of the subject and distance between the subject and the camera was 1.0 meter. This value of the distance is most suitable in capturing the video of the full head and partial shoulder of the subjects. Two different locations of experiment have been decided for normal and patient subjects. The experiment for normal subjects

was carried out in a good lighting condition in a laboratory of Universiti Malaysia Perlis (UniMAP) whereas the experiment for patient subjects was carried out in the physiotherapy department of Hospital Tunku Fauziah (HTF). The space for this experiment is limited and requires additional lighting support. Hence, a lighting source of white light was added to improve the video capturing. The white light used for lighting is not in any way harmful to the patient. The luminance of the light is 12 kcd/m<sup>2</sup> and the wattage is only 11W.

### **3.4 Landmarks Chosen and Facial Exercise Development Based On Muscle Movement and House-Brackmann (HB) Score**

In this study, the HB grading system was used as a reference score in assessing the facial nerve functions. It is the grading system which is the most used by clinicians all over the world as stated before in Chapter 2. The HB grading consists of three main parts of our face which is the forehead, eyes and mouth parts. The movement of these three parts has covered all of facial muscles. To evaluate these three parts, a study was designed to choose some desirable landmarks based on both muscle movements and also the HB score. The selection of the landmarks which have to be tracked later is a very crucial procedure since the goal of this study is to evaluate the facial nerve functions whether they are in normal condition or otherwise. The landmarks in Table 3.1 and Table 3.2 have been chosen by considering the muscle movement and the selection was agreed by a medical professional at the Otorhinolaryngology Department from Hospital Tunku Fauziah (HTF), Kangar. For Rise Forehead Exercise, the muscle involved is Frontalis. To evaluate the functionality of this muscle, the landmarks are placed based on the movement of the exercise. Rise Forehead Exercise is a movement of eyebrows or