IMMOBILIZATION OF α-AMYLASE (BAN) FOR SAGO STARCH HYDROLYSIS

KHAIRIL NAZUAN BIN MOHD

A report submitted in partial fulfillment Of the requirements for the award Of the degree of Bachelor of Chemical Engineering

Faculty of Chemical Engineering & Natural Resources University College of Engineering & Technology Malaysia

NOVEMBER 2006

DECLARATION

I declare that this thesis entitled "Immobilization of α -Amylase (BAN) for sago starch hydrolysis" is the result of my own research except as cited in the references. The thesis has not been accepted for any degree and is not concurrently submitted in candidature of any other degree.

Signature	:
Name	: KHAIRIL NAZUAN BIN MOHD
Date	: 20 November 2006

ACKNOWLEDGEMENT

In order to complete this research, I was in contact with many peoples, researchers, academicians and practitioners. All of them have assisted me in many ways towards completing this research. They also have contributed towards my understanding and thoughts. I would like to express my sincere appreciation to my supervisor, Mr Lau Sing Hui dan Miss Nina Suhaity Binti Azmi for their encouragement, guidance, critics and friendship in finishing my research.

I also would like to thanks the personnel of Faculty of Chemical Engineering and Natural Resource (FKKSA), especially lectures for their assistance and cooperation. Not forgotten to Mr. Joharizal Johari for their advices, motivation and ideas. Without their continued support and interest, this research would not have been the same as presented here.

My biggest thanks to the staff of FKKSA Chemical Laboratory especially Madam Norlia Mohammad, Miss Idayu and Mr. Anuar Ramli for their directly or indirectly influential and supportive in finishing this research.

My sincere appreciation also extends to all my colleagues, especially Mohd Rosli Ramly and others who have provided assistance at various occasions. Their views and tips are useful indeed. Your kindness is really appreciate and always in my mind forever.

Finally, my special appreciation is dedicated to my parents, Mr. Mohd bin Omar dan Madam Zainun binti Ismail and also my family for their tireless effort and endless moral support. Thank you.

TABLE OF CONTENTS

CHAPTER

TITTLE

PAGE

ORGANIZATION OF A THESIS

TITLE	i
DECLARATION	ii
DEDICATION	iii
ACKNOWLEDGEMENT	iv
ABSTRACT	v
ABSTRAK	vi
TABLE OF CONTENTS	vii
LIST OF TABLE	х
LIST OF FIGURES	xi
LIST OF APPENDICES	xii
LIST OF ABBREVIATION	xiii

1 INTRODUCTION

1.1	Introduction	1
1.2	Problem Statement	2
1.3	Objective	2
1.4	Scope	3

2 LITERATURE REVIEW

2.1	Introduction	4
2.2	Method of immobilization	5
	2.2.1 Carrier binding	5

		2.2.1.1 Physical adsorption	7
		2.2.1.2 Ionic bonding	8
		2.2.1.3 Covalent binding	9
	2.2.2	Cross linking	11
	2.2.3	Entrapping	12
2.3	Prope	rties of immobilized enzymes	14
	2.3.1	Stability	15
	2.3.2	Kinetic properties	15
2.4	Hydro	olysis process	16
2.5	α-Am	ylase (BAN)	17
2.6	Sago s	starch	18
2.7	Enzyn	ne assay	19
2.8	Туре	of enzyme assay	19
	2.8.1	Continuous assay	20
		2.8.1.1 Spectrophotometric	20
		2.8.1.2 Fluorometric	20
		2.8.1.3 Calorimetric	21
		2.8.1.4 Chemiluminescent	21
	2.8.2	Discontinuous assay	21
		2.8.2.1 Radiometric	22
		2.8.2.2 Chromatographic	22
2.9	Factor	rs to control in assays	22
	2.9.1	Temperature	22
	2.9.2	Enzyme concentration	23
	2.9.3	Substrate concentration	23

3 METHODOLOGY

3.1	Introduction		25
3.2	Overall methodology		25
3.3	Experi	mental design	27
	3.3.1	Raw material preparation	27
	3.3.2	Estimation of enzyme activities	27
	3.3.3	Entrapment of BAN in alginate beads	27
	3.3.4	Effect of sodium alginate concentration on the gel	27
		capsule permeability	
	3.3.5	Effect of Cacl2 concentration on the rigidity of the	28
		beads	
	3.3.6	Effects of the enzyme concentration in the capsules	28
3.4	Operat	ional efficiencies	28

4 **RESULT AND DISCUSSIONS**

Introduction	29
Sago starch hydrolysis	29
Effects of sodium alginate concentrations	32
Bead size	33
Kinetic analysis	34
	Introduction Sago starch hydrolysis Effects of sodium alginate concentrations Bead size Kinetic analysis

5 CONCLUSION AND RECOMMENDATIONS

5.1 Conclusion and Recommendations	35
------------------------------------	----

REFERENCES	37

APPENDICES 39-42

LIST OF TABLES

TABLE NO	TITLE	PAGE
4.2	Data for different concentration of sodium alginate	33
4.3	Data for different concentration of Cacl2	33
4.4	Data for different number of unit of α -amylase	34
4.5	Kinetic constant of enzyme	37

LIST OF FIGURES

FIGURE NO	TITLE P.	4GE
2.2.1	Carrier binding	6
2.2.3	Entrapping	13
3.1	Overall methodology	28
4.3	Effects of alginate concentration on immobilization	n 35
4.4	Effects of bead size on rate of starch hydrolysis	36

LIST OF APPENDICES

APPENDICES A	44
APPENDICES B	45
APPENDICES C	46

LIST OF ABBREVIATION

BAN	α-Amylase
рН	A measure of acidity
UGI	A multi-component reaction in organic chemistry
MTT	A laboratory test and a standard colorimetric assay
UV	Ultraviolet
NAD+	Nicotinamide adenine dinucleotide
NADH	Reduced form of NAD^+

IMMOBILIZATION OF α-AMYLASE (BAN) FOR SAGO STARCH HYDROLYSIS

KHAIRIL NAZUAN BIN MOHD

A report submitted in partial fulfillment Of the requirements for the award Of the degree of Bachelor of Chemical Engineering

Faculty of Chemical Engineering & Natural Resources University College of Engineering & Technology Malaysia

NOVEMBER 2006

ABSTRACT

 α -Amylase produced from *Baccilus Amyliquifaciens* (termamyl) was immobilized by entrapment in calcium alginate gel capsules and it was used repeatedly in batch processes of starch hydrolysis. The degree of starch degradation and operational stability of the immobilized system were increased by tailoring the characteristics of the capsules. Capsules prepared from 2% (w/v) sodium alginate and 5% (w/v) CaCl2 were suitable for up to 20 repeated uses, losing only 30% of their initial efficiency. These alginate/silica capsules carrying α -Amylase retained 90% of their initial efficiency after 20 starch hydrolysis batches and released more than 10,700 mg of reducing sugars during a processing period of 160 h.

ABSTRACT

 α -Amylase daripada *Baccilus Amyliquifaciens* (termamyl) dipegunkan dengan cara pemerangkapan di dalam kapsul kalsium alginat dan digunakan berulang-kali dalam proses hidrolisis kanji. Darjah penurunan kanji dan kestabilan operasi dalam sistem hidrolisis ini telah dapat ditingkatkan dengan menala karekteristik kapsul tersebut. Kapsul yang disediakan dengan menggunakan 2% (w/v) sodium alginat dan 5% (w/v) CaCl₂ adalah sangat sesuai digunakan sehingga 20 kali dengan hanya kehilangan 30% daripada kecekapannya pada permulaan proses. Kapsul alginat ini yang mengandungi α -Amylase dapat mencapai 90% daripada kecekapan permulaannya selepas 20 kali proses hidrolisis kanji dilakukan dan dapat membebaskan lebih daripada 10,700 mg gula penurun sepanjang tempoh proses selama 160 jam.

CHAPTER 1

INTRODUCTION

1.1 Introduction

Enzymes are protein molecules which serve to accelerate the chemical reactions of living cells (often by several orders of magnitude). Without enzymes, most biochemical reactions would be too slow to even carry out life processes. Enzymes display great specificity and are not permanently modified by their participation in reactions. Since they are not changed during the reactions, it is cost-effective to use them more than once. However, if the enzymes are in solution with the reactants and/or products it is difficult to separate them. Therefore, if they can be attached to the reactor in some way, they can be used again after the products have been removed. The term "immobilized" means unable to move or stationary. And that is exactly what an immobilized enzyme is: an enzyme that is physically attached to a solid support over which a substrate is passed and converted to product.

Enzymes can denature due to solvent effects and mechanical shear forces. Recovery of enzymes from reaction solutions and separation of the enzymes from substrates and products are in general very difficult. These problems can be successfully tackled by immobilization of the enzyme.

The main advantages of immobilized enzymes are:

• Easy separation from reaction mixture, providing the ability to control reaction times and minimize the enzymes lost in the product.

• Re-use of enzymes for many reaction cycles, lowering the total production cost of enzyme mediated reactions.

• Ability of enzymes to replace multiple standard chemical steps and provide enatomerically pure products.

1.2 Problem Statement

Enzymes are protein molecules which serve to accelerate the chemical reactions of living cells (often by several orders of magnitude). Without enzymes, most biochemical reactions would be too slow to even carry out life processes. Enzymes display great specificity and are not permanently modified by their participation in reactions. Since they are not changed during the reactions, it is cost-effective to use them more than once. However, if the enzymes are in solution with the reactants and/or products it is difficult to separate them. Therefore, if they can be attached to the reactor in some way, they can be used again after the products have been removed. The term "immobilized" means unable to move or stationary. And that is exactly what an immobilized enzyme is: an enzyme that is physically attached to a solid support over which a substrate is passed and converted to product

1.3 Objective

The objective of this research is to study the immobilization of α -amylase (BAN) and in alginate beads for sago starch hydrolysis.

1.4 Scope

- 1 To study the immobilization of α -amylase and amyloglucosidase for sago starch hydrolysis
- 2 To investigate the relationship between bead size and alginate concentration with alginate capsules
- 3 To optimize the capsule's characteristic.

REFERENCES

- Bentley IS, Williams EC. Starch conversion. In: Godfrey T, West S, editors. Industrial enzymology. London: Macmillan Press; 1996. p. 337–57.
- [2] Bernfeld, P. (1955), Amylases, alpha and beta. In: Colowick, S. P. and Kaplan, *Methods in enzymology*. New York : Academic Press. v. 1. pp. 149-158.
- [3] Cheetham PSJ. Principles of industrial biocatalysis and bioprocessing. In: Wiseman A, editor. Handbook of enzyme biotechnology. London: Ellis Horwood Ltd.; 1985. p. 83–233.
- [4] Chew Tek-Ann, Abu Hassan Isa and Mohd. Ghazali Mohayidin. 1999. Sago (*Metroxylon sagu* Rottboll), The Forgotten Palm, Journal of Sustainable Agriculture, forthcoming.
- [5] Dekker LA. Worthington enzyme manual. Freehold, New Jersey: Worthington Biochemical Corp.; 1977. p. 173–6.
- [6] Dixon, M , and Webb, E.c , *Enzymes*, Academic Press, Inc., New York, 1958.p.479.
- [7] Dziezak, J. D. (1991), Enzymes: Catalyst for food process. *Food Technol.*,
 45, 78-85

- [8] Emne'us, J. and Gorton, L. (1990), Effects on the hydrolysis of native starch and glycogen by a thermostable a-amylase after immobilization on solid supports. *Anal. Chim. Acta.*, 234, 97-106.
- [9] F. Godia and C. Sole, Fluidized bed bioreactors. Biotechnol. Prog. 11 (1995), pp. 479–497.
- [10] G.L. Miller, Use of dinitrosalicylic acid reagent for determination of reducing sugar. Anal. Chem. 31 (1959), pp. 426–428.
- [11] Linko, Y. Y.; Saarinen, P and Linko, M. (1975), Starch conversion by soluble and immobilized a-amylase. *Biotechnol. Bioeng.*, 17, 153-159
- [12] Mosbach K, Mattiasson B. Multistep enzyme systems. In: Mosbach K, editor. Methods in enzymology, vol. 44. New York: Academic Press Inc.; 1976. p. 453–78.
- [13] O. Smirsod and G. Skjak-Braek, Alginate as immobilization matrix for cells. Trends Biotechnol. 8 (1990), pp. 71–78.
- [14] W.S. Crabb and J.K. Shetty, Commodity scale production of sugars from starches. Curr. Opin. Microbiol. 2 (1999), pp. 252–256.
- [15] Zanin, G. M.; Kambara, L. M.; Calsavara, L. P. V. and De Moraes, F. F. (1994), Performance of fixed and fluidized bed reactors with immobilized enzyme. *Appl.Biochem. Biotechnol.*, 45/46, 627-640.
- [16] <u>http://www.sigmaaldrich.com/Brands/Fluka</u> <u>Riedel_Home/OrganicSynthetic/Enzymes/Organic_Chemistry/Immobilizatio</u> <u>n.html</u>
- [17] <u>http://en.wikipedia.org/wiki/Amylase</u>