

## REFERENCES

- Aarabi, A., Honarvar, M., Mizani, M., Faghihian, H., & Gerami, A. (2016). Extraction and purification of ferulic acid as an antioxidant from sugar beet pulp by alkaline hydrolysis. *Italian Journal of Food Science*, 28(3), 362–375.
- Abdullah, N., Sulaiman, F., Miskam, M. A., & Taib, R. M. (2014). Characterization of banana (*Musa* spp.) pseudo-stem and fruit-bunch-stem as a potential renewable energy resource. *International Journal of Biological, Veterinary and Agricultural Engineering*, 8(8), 794–798.
- Adler, E. (1977). Lignin chemistry-past, present and future. *Wood Science and Technology*, 11(3), 169–218.
- Adom, K. K., & Liu, R. H. (2002). Antioxidant activity of grains. *Journal of Agricultural and Food Chemistry*, 50(21), 6182–6187.
- Agustini, L., Efiyanti, L., Faulina, S. A., & Santoso, E. (2012). Isolation and characterization of cellulase- and xylanase-producing microbes isolated from tropical forests in Java and Sumatra. *International Journal of Environment and Bioenergy*, 3(3), 154-167.
- Ahmad, M. N., & Holland, C. R. (1995). Growth kinetics of single-cell protein in batch fermenters. *Journal of Food Engineering*, 26(4), 443-452.
- Ahmed, I., Yokota, A., Yamazoe, A., & Fujiwara, T. (2007). Proposal of *Lysinibacillus boronitolerans* gen. nov. sp. nov. and transfer of *Bacillus fusiformis* to *Lysinibacillus fusiformis* comb. nov. and *Bacillus sphaericus* to *Lysinibacillus sphaericus* comb. nov. *International Journal of Systematic and Evolutionary Microbiology*, 57, 1117–1125.
- Akinbile, C. O., El-Latif, K. A., Abdullah, R., & Yusoff, M. S. (2011). Rice production and water use efficiency for self-sufficiency in Malaysia: A review. *Trends in Applied Sciences Research*, 6(10), 1127.
- Akpınar, Ö., & Usal, G. (2015). Investigation of the effect of temperature and alkaline concentration on the solubilization of phenolic acids from dilute acid-pretreated wheat straw. *Food and Bioproducts Processing*, 95, 272-280.
- Alwi, H., Idris, J., Musa, M., & Ku Hamid, K. H. (2013). A preliminary study of banana stem juice as a plant-based coagulant for treatment of spent coolant wastewater. *Journal of Chemistry*. 2013, 1-7.

- Amin, M., Rakhisi, Z., & Ahmady, A. Z. (2015). Isolation and identification of *Bacillus* species from soil and evaluation of their antibacterial properties. *Avicenna J Clin Microb Infec*, 2, 10–13.
- Ascher, U. M., & Petzold, L. R. (1998). *Computer methods for ordinary differential equations and differential-algebraic equations*. Philadelphia: Society for Industrial and Applied Mathematics (SIAM).
- Aslan, N. (2008). Application of response surface methodology and central composite rotatable design for modeling and optimization of a multi-gravity separator for chromite concentration. *Powder Technology*, 185(1), 80–86.
- Aziz, N. A. A., Ho, L. H., Azahari, B., Bhat, R., Cheng, L.-H., & Ibrahim, M. N. M. (2011). Chemical and functional properties of the native banana (*Musa acuminata* × *balbisiana* Colla cv. Awak) pseudo-stem and pseudo-stem tender core flours. *Food Chemistry*, 128(3), 748–753.
- Baciu, I. E., & Jordening, H. J. (2003). Kinetics of galacturonic acid release from sugar beet pulp. *Enzyme Microb. Technol.*, 34, 505-512.
- Bader, J., Mast-Gerlach, E., Popović, M. K., Bajpai, R., & Stahl, U. (2010). Relevance of microbial coculture fermentations in biotechnology. *Journal of Applied Microbiology*. 109(2), 371-387.
- Badoei-Dalfard, A., & Karami, Z. (2013). Screening and isolation of an organic solvent tolerant-protease from *Bacillus* sp. JER02: Activity optimization by response surface methodology. *Journal of Molecular Catalysis B: Enzymatic*, 89, 15-23.
- Balasubramanian, N., & Simoes, N. (2014). *Bacillus pumilus* S124A carboxymethyl cellulase; a thermo stable enzyme with a wide substrate spectrum utility. *International Journal of Biological Macromolecules*, 67, 132-139.
- Barberousse, H., Roiseux, O., Robert, C., Paquot, M., Deroanne, C., & Blecker, C. (2008). Analytical methodologies for quantification of ferulic acid and its oligomers. *Journal of the Science of Food and Agriculture*. 88, 1494–1511.
- Bartolome, B., Faulds, C. B., & Williamson, G. (1997). Enzymic release of ferulic acid from barley spent grain. *Journal of Cereal Science*, 25, 285–288.
- Bartolomé, B., Gómez-Cordovés, C., Sancho, A. I., Díez, N., Ferreira, P., Soliveri, J., & Copa-Patiño, J. L. (2003). Growth and release of hydroxycinnamic acids from brewer's spent grain by *Streptomyces avermitilis* CECT 3339. *Enzyme and Microbial Technology*, 32(1), 140–144.

- Bates, M. K., Phillips, D. S., & O'Bryan, J. (2011). Shaker agitation rate and orbit affect growth of cultured bacteria. Retrieved from Fisher Scientific website: <https://tools.thermofisher.com/content/sfs/brochures/D02594~.pdf>
- Baucher, M., Monties, B., Montagu, M. Van, & Boerjan, W. (1998). Biosynthesis and genetic engineering of lignin. *Critical Reviews in Plant Sciences*, 17(2), 125–197.
- Berg, J. M., Tymoczko, J. L., & Stryer, L. (2002). *The Michaelis-Menten Model Accounts for the Kinetic Properties of Many Enzymes* (5th ed). New York: W H Freeman.
- Bertrand, S., Bohni, N., Schnee, S., Schumpp, O., Gindro, K., & Wolfender, J. L. (2014). Metabolite induction via microorganism co-culture: A potential way to enhance chemical diversity for drug discovery. *Biotechnology Advances*, 32, 1180-1204.
- Beyenal, H., & Babauta, J. T. (2015). *Biofilms in Bioelectrochemical Systems: From Laboratory Practice to Data Interpretation*. Hoboken, USA: John Wiley & Sons.
- Bezerra, R. M., Fraga, I. & Dias, A. A. (2013). Utilization of integrated Michaelis–Menten equations for enzyme inhibition diagnosis and determination of kinetic constants using Solver supplement of Microsoft Office Excel. *Computer methods and programs in biomedicine*, 109(1), 26-31.
- Bezerra, M. A., Santelli, R. E., Oliveira, E. P., Silveira, V. L., & Escalera, L. A. (2008). Response surface methodology (RSM) as a tool for optimization in analytical chemistry. *Talanta*, 76, 965–977.
- Bidlack, J., Malone, M., & Benson, R. (1992). Molecular structure and component integration of secondary cell walls in plants. *Proceedings of the Oklahoma Academy of Science*, 72, 51–56.
- Bilba, K., Arsene, M. A., & Ouensanga, A. (2007). Study of banana and coconut fibers botanical composition, thermal degradation and textural observations. *Bioresource Technology*, 98(1), 58–68.
- Bisswanger, H. (2014). Enzyme assays. *Perspectives in Science*, 1(1), 41-55.
- Box, G. E. P., Hunter, W. J., & Hunter, J. S. (1978). *Statistics for Experimenters: An Introduction to Design, Data Analysis, and Model Building*. USA: John Wiley & Sons, Inc.
- Box, G. E. P., & Wilson, K. B. (1951). On the experimental attainment of optimum conditions. *Journal of the Royal Statistical Society*, 13(1), 1–45.

- Brown, S., & Coleman, G. (1975). Messenger ribonucleic acid content of *Bacillus amyloliquefaciens* throughout its growth cycle compared with *Bacillus subtilis* 168. *Journal of Molecular Biology*, 96(2), 345-352.
- Buranov, A. U., & Mazza, G. (2008). Lignin in straw of herbaceous crops. *Industrial Crops and Products*, 28(3), 237–259.
- Butler, N. A. (2008). Defining equations for two-level factorial designs. *Journal of Statistical Planning and Inference*, 138(10), 3157-3163.
- Chai, L., Zhang, H., Yang, W., Zhu, Y., Yang, Z., Zheng, Y., & Chen Y. (2013). *Journal of Central South University of Technology*. 20(7), 1964-1970.
- Chakuton, K., Puangpronpitag, D., & Nakornriab, M. (2012). Phytochemical content and antioxidant activity of colored and non-colored Thai rice cultivars. *Asian Journal of Plant Sciences*, 11(6), 285.
- Chamkha, M., Patel, B. K. C., Garcia, J. L., & Labat, M. (2001). Isolation of *Clostridium bifermentans* from oil mill wastewaters converting cinnamic acid to 3-phenylpropionic acid and emendation of the species. *Anaerobe*, 7(4), 189-197.
- Chang, S. H., Teng, T. T., & Ismail, N. (2011). Screening of factors influencing Cu(II) extraction by soybean oil-based organic solvents using fractional factorial design. *Journal of Environmental Management*, 92(10), 2580–2585.
- Chapra, S. C., & Canale, R. P. (2010). *Numerical methods for engineers*. New York: McGraw-Hill.
- Claus, D. (1992). A standardized gram staining procedure. *World Journal of Microbiology and Biotechnology*, 8(4), 451-452.
- Crepin, V. F., Faulds, C. B., & Connerton, I. F. (2003). A non-modular type B feruloyl esterase from *Neurospora crassa* exhibits concentration-dependent substrate inhibition. *The Biochemical Journal*, 370, 417–427.
- Cruz, J. M., Domínguez, J. M., Domínguez, H., & Parajó, J. C. (2001). Antioxidant and antimicrobial effects of extracts from hydrolysates of lignocellulosic materials. *Journal of Agricultural and Food Chemistry*, 49(5), 2459–64.

- Damáσιο, A. R. L., Braga, C. M. P., Brenelli, L. B., Citadini, A. P., Mandelli, F., Cota, J., ... Squina, F. M. (2013). Biomass-to-bio-products application of feruloyl esterase from *Aspergillus clavatus*. *Applied Microbiology and Biotechnology*, 97(15), 6759–6767.
- Dodić, J. M., Vučurović, D. G., Dodić, S. N., Grahovac, J. A., Popov, S. D., & Nedeljković, N. M. (2012). Kinetic modelling of batch ethanol production from sugar beet raw juice. *Applied Energy*, 99, 192–197.
- Donaghy, J., Kelly, P. F., & McKay, A. M. (1998). Detection of ferulic acid esterase production by *Bacillus* spp. and lactobacilli. *Applied Microbiology and Biotechnology*, 50, 257–260.
- Duza, M. B., & Mastan, S. A. (2013). Microbial enzymes and their applications—a review. *American Journal of Pharm Research*, 3(8), 6208-6219.
- El-Mansi, E. M. T., Bryce, C. F., Demain, A. L., & Allman, A. R. (2011). *Fermentation microbiology and biotechnology*. USA: CRC press.
- Fadel, M. (2000). Production physiology of cellulases and  $\beta$ -glucosidase enzymes of *Aspergillus niger* grown under solid state fermentation conditions. *OnLine journal of biological sciences*, 1(5), 401-411.
- Faulds, C. B., & Williamson, G. (1995). Release of ferulic acid from wheat bran by a ferulic acid esterase (FAE-III) from *Aspergillus niger*. *Applied Microbiology and Biotechnology*, 43, 1082–1087.
- Fazary, A. E., & Ju, Y. H. (2007). Feruloyl esterases as biotechnological tools: Current and future perspectives. *Acta Biochimica et Biophysica Sinica*, 39, 811–828.
- Fazary, A. E., Ismadji, S., & Ju, Y. H. (2010). Studies on temperature dependent kinetics of *Aspergillus awamori* feruloyl esterase in water solutions. *Kinetics and Catalysis*, 51(1), 31–37.
- Felix, E., Clara, O., & Vincent, A. O. (2014). A kinetic study of the fermentation of cane sugar using *Saccharomyces cerevisiae*. *Open Journal of Physical Chemistry*, 4(1), 26-31.
- Ferreira, P., Diez, N., Faulds, C. B., Soliveri, J., & Copa-Patiño, J. L. (2007). Release of ferulic acid and feruloylated oligosaccharides from sugar beet pulp by *Streptomyces tendae*. *Bioresource Technology*, 98(8), 1522–1528.

- Ferreira, P., Diez, N., Gutierrez, C., Soliveri, J., & Copa-Patino, J. L. (1999). *Streptomyces avermitilis* CECT 3339 produces a ferulic acid esterase able to release ferulic acid from sugar beet pulp soluble feruloylated oligosaccharides. *Journal of the Science of Food and Agriculture*, 79, 440–442.
- Fogler, H. S. (2010). *Elements of chemical reaction engineering*. United Sate: Pearson Education.
- Furuya, T., & Kino, K. (2014). Catalytic activity of the two-component flavin-dependent monooxygenase from *Pseudomonas aeruginosa* toward cinnamic acid derivatives. *Applied Microbiology and Biotechnology*, 98(3), 1145–1154.
- Ganesapillai, M., Mathew, M., Singh, A., & Simha, P. (2016). Influence of microwave and ultrasound pretreatment on solvent extraction of bio-components from walnut (*Julgans regia L.*) shells. *Periodica Polytechnica Chemical Engineering*, 60(1), 1-9.
- Georgiou, S. D. (2007). New two-variable full orthogonal designs and related experiments with linear regression models. *Statistics and Probability Letters*, 77(1), 25–31.
- Gibson, L. J. (2012). The hierarchical structure and mechanics of plant materials. *Journal of the Royal Society Interface*, 9(76), 2749–2766.
- Godoy, L., Martínez, C., Carrasco, N., & Ganga, M. A. (2008). Purification and characterization of a p-coumarate decarboxylase and a vinylphenol reductase from *Brettanomyces bruxellensis*. *International Journal of Food Microbiology*, 127(1), 6-11.
- Goers, L., Freemont, P., & Polizzi, K. M. (2014). Co-culture systems and technologies: taking synthetic biology to the next level. *Journal of the Royal Society Interface*, 11(96), 1-13.
- Golshani, T., Jorjani, E., Chelgani, S. C., Shafaei, S. Z., & Nafechi, Y. H. (2013). Modeling and process optimization for microbial desulfurization of coal by using a two-level full factorial design. *International Journal of Mining Science and Technology*, 23, 261-265.
- Golub, K. W., Forrest, A. K., Wales, M. E., Hammett, A. J. M., Cope, J. L., Wilkinson, H. H., & Holtzapple, M. T. (2013). Comparison of three screening methods to select mixed-microbial inoculum for mixed-acid fermentations. *Bioresource Technology*, 130, 739–749.

- Gómez, J. M., & Cantero, D. (2003). Kinetic study of biological ferrous sulphate oxidation by iron-oxidising bacteria in continuous stirred tank and packed bed bioreactors. *Process Biochemistry*, 38(6), 867–875.
- Gopalan, N., Rodríguez-Duran, L. V., Saucedo-Castaneda, G., & Nampoothiri, K. M. (2015). Review on technological and scientific aspects of feruloyl esterases: A versatile enzyme for biorefining of biomass. *Bioresource Technology*, 193, 534-544.
- Gottipati, R., & Mishra, S. (2010). Process optimization of adsorption of Cr(VI) on activated carbons prepared from plant precursors by a two-level full factorial design. *Chemical Engineering Journal*, 160(1), 99–107.
- Guimaraes, J. L., Frollini, E., Da Silva, C. G., Wypych, F., & Satyanarayana, K. G. (2009). Characterization of banana, sugarcane bagasse and sponge gourd fibers of Brazil. *Industrial Crops and Products*, 30(3), 407–415.
- Guowei, S., Chunji, D., He, C., & Xu, W. (2013). Application of Box-Behnken design in optimization for crude polysaccharides from fruits of *Tribulusteristris L.* *Journal of Chemical and Pharmaceutical Research*, 5(10), 342-350.
- Gutierrez-Correa, M., Portal, L., Moreno, P., & Tengerdy, R. P. (1999). Mixed culture solid substrate fermentation of *Trichoderma reesei* with *Aspergillus niger* on sugar cane bagasse. *Bioresource Technology*, 68(2), 173–178.
- Hagedorn, S., & Kaphammer, B. (1994). Microbial biocatalysis in the generation of flavor and fragrance chemicals. *Annual Review of Microbiology*, 48(19), 773–800.
- Hall, B. G., Acar, H., Nandipati, A., & Barlow, M. (2014). Growth rates made easy. *Molecular Biology and Evolution*, 31(1), 232-238.
- Hamzaoui, A. H., Jamoussi, B., & M'nif, A. (2008). Lithium recovery from highly concentrated solutions: Response surface methodology (RSM) process parameters optimization. *Hydrometallurgy*, 90, 1-7.
- Hashizume, K., Ito, T., Ishizuka, T., & Takeda, N. (2013). Formation of ethyl ferulate by rice koji enzyme in sake and mirin mash conditions. *Journal of bioscience and bioengineering*, 116(2), 209-213.
- Herrick, J. B., Madsen, E. L., Batt, C. A., & Ghiorse, W. C. (1993). Polymerase chain reaction amplification of naphthalene-catabolic and 16S rRNA gene sequences from indigenous sediment bacteria. *Applied and Environmental Microbiology*, 59(3), 687-694.

- Hooda, A., Nanda, A., Jain, M., Kumar, V., & Rathee, P. (2012). Optimization and evaluation of gastroretentive ranitidine HCl microspheres by using design expert software. *International journal of biological macromolecules*, 51(5), 691-700.
- Hoorman, J. J., & Islam, R. (2010). Understanding soil microbes and nutrient recycling (Fact sheet SAG-16-10). Retrieved from The Ohio State University Extension website: <http://ohioline.osu.edu/factsheet/SAG-16>
- Hoson, T., & Wakabayashi, K. (2015). Role of the plant cell wall in gravity resistance. *Phytochemistry*, 112, 84–90.
- Hua, D., Ma, C., Song, L., Lin, S., Zhang, Z., Deng, Z., & Xu, P. (2007). Enhanced vanillin production from ferulic acid using adsorbent resin. *Applied Microbiology and Biotechnology*, 74(4), 783–790.
- Huang, Y. C., Chen, Y. F., Chen, C. Y., Chen, W. L., Ciou, Y. P., Liu, W. H., & Yang, C. H. (2011). Production of ferulic acid from lignocellulolytic agricultural biomass by *Thermobifida fusca* thermostable esterase produced in *Yarrowia lipolytica* transformant. *Bioresource Technology*, 102, 8117–8122.
- Huang, C., Wu, H., Liu, Z.-J., Cai, J., Lou, W.-Y., & Zong, M.-H. (2012). Effect of organic acids on the growth and lipid accumulation of oleaginous yeast *Trichosporon fermentans*. *Biotechnology for Biofuels*, 5(1), 1-12.
- Humberstone, F. J., & Briggs, D. E. (2000). Extraction and assay of ferulic acid esterase from malted barley. *Journal of the Institute of Brewing*, 106(1), 21–30.
- Hussain, T., Roohi, A., Munir, S., Ahmed, I., Khan, J., Edel-Hermann, V., ... Anees, M. (2013). Biochemical characterization and identification of bacterial strains isolated from drinking water sources of Kohat, Pakistan. *African Journal of Microbiology Research*, 7(16), 1579-1590.
- Ibemhal, D. A., Majumder, P. B., Imotomba, R. K., & Laishram, J. M. (2015). Genomic DNA extraction from the black scented rice (Chakhao). *Indian Journal of Plant Sciences*, 4(1), 34–37.
- Idrees, M., Adnan, A., Malik, F., & Qureshi, F. A. (2013). Enzymatic saccharification and lactic acid production from banana pseudo-stem through optimized pretreatment at lowest catalyst concentration. *EXCLI Journal*, 12, 269–281.
- INIBAP. (2007). *Bananas International Network for the improvement of banana and plaintain*. Retrieved from <http://www.bananas.biodiversityinternational.org/files/files/pdf/publications/brochurebananas>

- Ismail, S. N., & Zainol, N. (2014a). Factorial analysis of ferulic acid extraction from banana stem waste. *International Journal of Applied Engineering Research*, 9, 6823-6833.
- Ismail, S. N., & Zainol, N. (2014b). Optimization of ferulic acid extraction from banana stem waste. *Asian Journal of Microbiology, Biotechnology and Environmental Sciences*, 16(3), 479-484.
- Irshad, M., Anwar, Z., But, H. I., Afroz, A., Ikram, N., & Rashid, U. (2013). The industrial applicability of purified cellulase complex indigenously produced by *Trichoderma viride* through solid-state bio-processing of agro-industrial and municipal paper wastes. *BioResources*, 8(1), 145–157.
- Ito, T., Suzuki, N., Nakayama, A., Ito, M., & Hashizume, K. (2014). Factors affecting phenolic acid liberation from rice grains in the sake brewing process. *Journal of Bioscience and Bioengineering*, 118(6), 640–645.
- Jabeen, H., Iqbal, S., Anwar, S., & Parales, R. E. (2015). Optimization of profenofos degradation by a novel bacterial consortium PBAC using response surface methodology. *International Biodeterioration and Biodegradation*, 100, 89–97.
- Janda, J. M., & Abbott, S. L. (2002). Bacterial identification for publication: when is enough enough?. *Journal of clinical microbiology*, 40(6), 1887-1891.
- Javed, M. M., Zahoor, S., Shafaat, S., Mehmooda, I., Gul, A., Rasheed, H., ... & Aftab, M. N. (2012). Wheat bran as a brown gold: Nutritious value and its biotechnological applications. *African Journal of Microbiology Research*, 6(4), 724-733.
- Jo, J. H., Lee, D. S., Park, D., Choe, W. S., & Park, J. M. (2008). Optimization of key process variables for enhanced hydrogen production by *Enterobacter aerogenes* using statistical methods. *Bioresource Technology*, 99(6), 2061–2066.
- Johnson, T. R., & Case, C. L. (2013). *Laboratory Experiments in Microbiology* (10th ed). USA: Pearson Education.
- Johnson, K. A., & Goody, R. S. (2011). The original Michaelis constant: translation of the 1913 Michaelis–Menten paper. *Biochemistry*, 50(39), 8264-8269.
- Kamalipourazad, M., Sharifi, M., Maivan, H. Z., Behmanesh, M., & Chashmi, N. A. (2016). Induction of aromatic amino acids and phenylpropanoid compounds in *Scrophularia striata* Boiss. cell culture in response to chitosan-induced oxidative stress. *Plant Physiology and Biochemistry*, 107, 374–384.

- Kaminari, N. M. S., Ponte, M. J. J. S., Ponte, H. A., & Neto, A. C. (2005). Study of the operational parameters involved in designing a particle bed reactor for the removal of lead from industrial wastewater -central composite design methodology. *Chemical Engineering Journal*, 105(3), 111–115.
- Karagöz, P., & Özkan, M. (2014). Ethanol production from wheat straw by *Saccharomyces cerevisiae* and *Scheffersomyces stipitis* co-culture in batch and continuous system. *Bioresource Technology*, 158, 286–293.
- Karazhiyan, H., Razavi, S. M., & Phillips, G. O. (2011). Extraction optimization of a hydrocolloid extract from cress seed (*Lepidium sativum*) using response surface methodology. *Food Hydrocolloids*, 25, 915-920.
- Kareem, S. A., Kefas, H. M., Chior, T. J., & Latinwo, G. K. (2011). Kinetics of fermentation by enzymes: a mathematical approach. *AU Journal of Technology*, 15(2), 109-114.
- Karunaratne, R., & Zhu, F. (2016). Physicochemical interactions of maize starch with ferulic acid. *Food Chemistry*, 199, 372–379.
- Kleerebezem, R., & van Loosdrecht, M. C. (2007). Mixed culture biotechnology for bioenergy production. *Current Opinion in Biotechnology*. 18(3), 207-212.
- Klimova, T., Esquivel, A., Reyes, J., Rubio, M., Bokhimi, X., & Aracil, J. (2006). Factorial design for the evaluation of the influence of synthesis parameters upon the textural and structural properties of SBA-15 ordered materials. *Microporous and Mesoporous Materials*, 93(1–3), 331–343.
- Kondo, T., & Kondo, M. (1996). Efficient production of acetic acid from glucose in a mixed culture of *Zymomonas mobilis* and *Acetobacter* sp. *Journal of Fermentation and Bioengineering*, 81, 42–46.
- Körbahti, B. K., & Rauf, M. A. (2008). Response surface methodology (RSM) analysis of photoinduced decoloration of toluidine blue. *Chemical Engineering Journal*, 136(1), 25–30.
- Koshijima, T., & Watanabe, T. (2013). *Association between lignin and carbohydrates in wood and other plant tissues*. New York: Springer Series in Wood Science.
- Kotay, S. M., & Das, D. (2007). Microbial hydrogen production with *Bacillus coagulans* IIT-BT S1 isolated from anaerobic sewage sludge. *Bioresource Technology*, 98(6), 1183-1190.

- Kovárová-Kovar, K., & Egli, T. (1998). Growth kinetics of suspended microbial cells: from single-substrate-controlled growth to mixed-substrate kinetics. *Microbiology and Molecular Biology Reviews*, 62(3), 646-666.
- Kroon, P. A, Faulds, C. B., Brézillon, C., & Williamson, G. (1997). Methyl phenylalkanoates as substrates to probe the active sites of esterases. *European Journal of Biochemistry / FEBS*, 248(1), 245-251.
- Kryndushkin, D. S., Alexandrov, I. M., Ter-Avanesyan, M. D., & Kushnirov, V. V. (2003). Yeast [PSI+] prion aggregates are formed by small Sup35 polymers fragmented by Hsp104. *Journal of Biological Chemistry*, 278(49), 49636-49643.
- Kühnel, S., Pouvreau, L., Appeldoorn, M. M., Hinz, S. W. A., Schols, H. A., & Gruppen, H. (2012). The ferulic acid esterases of *Chrysosporium lucknowense* C1: Purification, characterization and their potential application in biorefinery. *Enzyme and Microbial Technology*, 50(1), 77-85.
- Kumar, N., & Pruthi, V. (2014). Potential applications of ferulic acid from natural sources. *Biotechnology Reports*. 4, 86-93.
- Lee, K. E., Morad, N., Teng, T. T., & Poh, B. T. (2012). Factorial experimental design for reactive dye flocculation using inorganic-organic composite polymer. *APCBEE Procedia*, 1, 59-65.
- Lemos, M., Borges, A., Teodósio, J., Araújo, P., Mergulhão, F., Melo, L., & Simões, M. (2014). The effects of ferulic and salicylic acids on *Bacillus cereus* and *Pseudomonas fluorescens* single- and dual-species biofilms. *International Biodeterioration and Biodegradation*, 86, 42-51.
- Li, K., Fu, S., Zhan, H., Zhan, Y., & Lucia, L. A. (2010). Analysis of the chemical composition and morphological structure of banana pseudo-stem. *BioResources*, 5(2), 576-585.
- Lineweaver, H., & Burk, D. (1934). The determination of enzyme dissociation constants. *Journal of the American Chemical Society*, 56(3), 658-666.
- Linh, T. N., Fujita, H., & Sakoda, A. (2017). Release kinetics of esterified p-coumaric acid and ferulic acid from rice straw in mild alkaline solution. *Bioresource Technology*, 232, 192-203.
- Liyana-Pathirana, C., & Shahidi, F. (2005). Optimization of extraction of phenolic compounds from wheat using response surface methodology. *Food Chemistry*, 93(1), 47-56.

- Lomascolo, A., Uzan-Boukhris, E., Herpoël-Gimbert, I., Sigoillot, J. C., & Lesage-Meessen, L. (2011). Peculiarities of *Pycnopus* species for applications in biotechnology. *Applied Microbiology and Biotechnology*, 92(6), 1129-1149.
- Mahouachi, J., López-Climent, M. F., & Gómez-Cadenas, A. (2014). Hormonal and hydroxycinnamic acids profiles in banana leaves in response to various periods of water stress. *Scientific World Journal*, 2014, 1-9.
- Mandal, A., Kar, S., Mohapatra, P. K. D, Maity, C., Pati, B. R., & Mondal, K. C. (2012). Regulation of xylanase biosynthesis in *Bacillus cereus* BSA1. *Applied Biochemistry and Biotechnology*. 167(5), 1052–1060.
- Maoka, F. Tanimoto, M. Sano, K. Tsurukawa, T. Tsuno, S. Tsujiwaki, K., ... Takii, K. (2008). Effects of dietary supplementation of ferulic acid and  $\gamma$ -Oryzanol on integument color and suppression of oxidative stress in cultured Red Sea Bream, *Pagrus major*, *Journal of Oleo Science*, 57(2), 133-137.
- Marchetti, P., Jimenez Solomon, M. F., Szekely, G., & Livingston, A. G. (2014). Molecular separation with organic solvent nanofiltration: a critical review. *Chemical reviews*, 114(21), 10735-10806.
- Martendal, E., Budziak, D., & Carasek, E. (2007). Application of fractional factorial experimental and Box-Behnken designs for optimization of single-drop microextraction of 2, 4, 6-trichloroanisole and 2, 4, 6-tribromoanisole from wine samples. *Journal of chromatography A*, 1148(2), 131-136.
- Martinek, R. (1969). Practical clinical enzymology. *The American Journal of Medical Technology*, 31, 162.
- Masaji, Y. (1999). Preservation measures for food products. Preventive method of coloring agricultural products using ferulic acid. *Journal of Food Chemistry*, 8, 76–79.
- Masoumi, H. R. F., Kassim, A., Basri, M., & Abdullah, D. K. (2011). Determining optimum conditions for lipase-catalyzed synthesis of triethanolamine (TEA)-based esterquat cationic surfactant by a Taguchi robust design method. *Molecules*, 16, 4672-4680.
- Max, B., Salgado, J. M., Cortes, S., & Dominguez, J. M. (2010). Extraction of phenolic acids by alkaline hydrolysis from the solid residue obtained after prehydrolysis of trimming vine shoots. *Journal of Agricultural and Food Chemistry*, 58(3), 1909-1917.

- McAuley, K. E., Svendsen, A., Patkar, S. A., & Wilson, K. S. (2004). Structure of a feruloyl esterase from *Aspergillus niger*. *Acta Crystallographica Section D: Biological Crystallography*, 60(5), 878–887.
- McCleary, B., & Prosky, L. (Eds.). (2008). *Advanced dietary fibre technology*. USA: Blackwell Science Ltd.
- McMurrough, I., Madigan, D., Donnelly, D., Hurley, J., Doyle, A., Hennigan, G., ... Smyth, M. R. (1996). Control of ferulic acid and 4-vinyl guaiacol in brewing. *Journal of the Institute of Brewing*, 102(5), 327–332.
- Meehnian, H., Jana, A. K., & Jana, M. M. (2017). Pretreatment of cotton stalks by synergistic interaction of *Daedalea flavida* and *Phlebia radiata* in co-culture for improvement in delignification and saccharification. *International Biodeterioration & Biodegradation*, 117, 68-77.
- Mellefont, L. A., McMeehin, T. A., & Ross, T. 2008. Effect of relative inoculum concentration on *Listeria monocytogenes* growth in co-culture. *International Journal of Food Microbiology*, 121(2), 157-168.
- Micard, V., Renard, C. M. G. C., & Thibault, J. F. (1994). Studies on enzymic release of ferulic acid from sugar-beet pulp. *LWT - Food Science and Technology*, 27(1), 59–66.
- Michaevlevski, I., Kirshenbaum, N., & Sharon, M. (2010). T-wave ion mobility-mass spectrometry: basic experimental procedures for protein complex analysis. *Journal of Visualized Experiments*, 1–7.
- Michaelis, L., & Menten, M. (1913). Die kinetic der invertinwirkung/ The Kinetics of Invertase Action. *Biochemistry Zeilung*, 49, 333-69.
- Minton, A. P. (2001). The Influence of macromolecular crowding and macromolecular confinement on biochemical reactions in physiological media. *Journal of Biological Chemistry*. 276, 10577–10580.
- Mo, X. L., & Xu, Y. (2010). Ferulic acid release and 4-vinylguaiacol formation during chinese rice wine brewing and fermentation. *Journal of the Institute of Brewing*, 116(3), 304–311.

- Mohanasrinivasan, V., Diksha, D., Suganthi, V., Selvarajan, E., & Subathra Devi, C. (2014). Process optimization for enhanced production of  $\alpha$ -amylase by submerged (SmF) and solid state fermentation (SSF) using *Bacillus pumilus* VITMDS2. *Research Journal of Pharmaceutical, Biological and Chemical Sciences*, 5(2), 1784–1800.
- Mohapatra, D., Mishra, S., & Sutar, N. (2010). Banana and its by-product utilization: an overview. *Journal of Scientific and Industrial Research*, 69(5), 323-329.
- Moravec, T. L. (2012). Soil microbes. *Biotechnology and Bioengineering*, 27(10), 1482-1489.
- Morgan, M. C., Boyette, M., Goforth, C., Sperry, K.V., & Greene, S. R. (2009). Comparison of the Biolog OmniLog Identification System and 16S ribosomal RNA gene sequencing for accuracy in identification of a typical bacteria of clinical origin. *Journal of Microbiological Methods*, 79, 336-343.
- Mrwebi, M. (2004). *Testing Monod: growth rate as a function of glucose concentration in Saccharomyces cerevisiae* (Master's thesis, University of Stellenbosch). Retrieved from [https://scholar.sun.ac.za/bitstream/handle/10019.1/16398/mrwebi\\_testing\\_2004.pdf?sequence=1](https://scholar.sun.ac.za/bitstream/handle/10019.1/16398/mrwebi_testing_2004.pdf?sequence=1)
- Mukherjee, G., Singh, R. K., Mitra, A., & Sen, S. K. (2007). Ferulic acid esterase production by *Streptomyces* sp. *Bioresource Technology*, 98, 211–213.
- Müller, V. (2001). Bacterial fermentation. *Encyclopedia of Life Science*, 1–7.
- Mussatto, S. I., Dragone, G., & Roberto, I. C. (2007). Ferulic and p-coumaric acids extraction by alkaline hydrolysis of brewer's spent grain. *Industrial Crops and Products*, 25(2), 231–237.
- Myburg, A. A., Lev-Yadun, S., & Sederoff, R. R. (2013). Xylem structure and function. USA: John Wiley & Sons.
- Myers, R. H., & Montgomery, D. C. (2002). *Response surface methodology: Process and product optimization using designed experiments*, USA: John Wiley & Sons.
- Nakamura, S., Wakabayashi, K., Nakai, R., Aono, R., & Horikoshi, K. (1993). Production of alkaline xylanase by a newly isolated alkaliphilic *Bacillus* sp. strain 41M-1. *World Journal of Microbiology and Biotechnology*, 9(2), 221–224.

- Nemeckova, I., Solichova, K., Roubal, P., Uhrova, B., & Svirakove, E. (2011). Methods for detection of *Bacillus* sp., *B. cereus*, and *B. licheniformis* in raw milk. *Czech Journal of Food Sciences*, 29, S55–S60.
- Nienow, A. W., McLeod, G., & Hewitt, C. J. (2010). Studies supporting the use of mechanical mixing in large scale beer fermentations. *Biotechnology Letters*, 32(5), 623-633.
- Noor Hasyierah, M. S. N., Zulkali, M. M. D., & Ku Syahidah, K. I. (2008). Ferulic acid from lignocellulosic biomass: review. *Proceeding of MUCET2008*. pp. 1-8
- Odabaş, H. İ., & Koca, I. (2016). Application of response surface methodology for optimizing the recovery of phenolic compounds from hazelnut skin using different extraction methods. *Industrial Crops and Products*, 91, 114–124.
- Oh, S. E., Kim, K. S., Choi, H. C., Cho, J., & Kim, I. S. (2000). Kinetics and physiological characteristics of autotrophic denitrification by denitrifying sulfur bacteria. *Water science and technology*, 42(3-4), 59-68.
- Okada, T., Nakagawa, K., & Yamaguchi, N. (1982). Antioxidative activities of amino compounds on fats and oils: VIII. Antioxidative activity of ferulate and the synergistic effect of amino compounds. *Journal of the Japanese Society for Food Science and Technology-Nippon Shokuhin Kogyo Gakkaishi*, 29(5), 305–309.
- Okpokwasili, G. C., & Nweke, C. O. (2005). Microbial growth and substrate utilization kinetics. *African Journal of Biotechnology*, 5(4), 305-317.
- Oliveira, L., Cordeiro, N., Evtuguin, D. V., Torres, I. C., & Silvestre, A. J. D. (2007). Chemical composition of different morphological parts from “Dwarf Cavendish” banana plant and their potential as a non-wood renewable source of natural products. *Industrial Crops and Products*, 26(2), 163–172.
- Oliveira, L., Freire, C. S. R., Silvestre, A. J. D., Cordeiro, N., Torres, I. C., & Evtuguin, D. (2006). Lipophilic extractives from different morphological parts of banana plant “Dwarf Cavendish.” *Industrial Crops and Products*, 23(2), 201–211.
- Olkowski, A. A., Amarowicz, R., Peiquiang, Y., McKinnon, J. J., & Maenz, D. D. (2003). A rapid HPLC method for determination of major phenolic acids in plant material. *Polish journal of food and nutrition sciences*, 12(1), 53-58.
- Olmos, J. C., & Hansen, M. Z. (2012). Enzymatic depolymerization of sugar beet pulp: Production and characterization of pectin and pectic-oligosaccharides as a potential source for functional carbohydrates. *Chemical Engineering Journal*, 192, 29-36.

- Omoniyi, T. E., & Olorunnisola, A. O., (2014). Experimental characterisation of bagasse biomass material for energy production. *International Journal of Engineering and Technology*, 4(10), 582-589.
- Onsekizoglu, P., Bahceci, K. S., & Acar, J. (2010). The use of factorial design for modeling membrane distillation. *Journal of Membrane Science*, 349, 225-230.
- Ou, S., & Kwok, K. C. (2004). Ferulic acid: Pharmaceutical functions, preparation and applications in foods. *Journal of the Science of Food and Agriculture*. 84(11), 1261–1269.
- Ou, S., Zhang, J., Wang, Y., & Zhang, N. (2011). Production of feruloyl esterase from *Aspergillus niger* by solid-state fermentation on different carbon sources. *Enzyme Research*, 2011, 1-4.
- Papagianni, M., Nokes, S. E., & Filer, K. (2001). Submerged and solid-state phytase fermentation by *Aspergillus niger*: Effects of agitation and medium viscosity on phytase production, fungal morphology and inoculum performance. *Food Technology and Biotechnology*, 39(4), 319–326.
- Pappu, A., Patil, V., Jain, S., Mahindrakar, A., Haque, R., & Thakur, V. K. (2015). Advances in industrial prospective of cellulosic macromolecules enriched banana biofibre resources: A review. *International Journal of Biological Macromolecules*. 79, 449-458.
- Price, N. C. (1985). The determination of  $K_m$  values from lineweaver-burk plots. *Biochemical Education*, 13(2), 81-81.
- Pasangulapati, V., Ramachandriya, K. D., Kumar, A., Wilkins, M. R., Jones, C. L., & Huhnke, R. L. (2012). Effects of cellulose, hemicellulose and lignin on thermochemical conversion characteristics of the selected biomass. *Bioresource Technology*, 114, 663–669.
- Pavan, F. A., Gushikem, Y., Mazzocato, A. C., Dias, S. L. P., & Lima, E. C. (2007). Statistical design of experiments as a tool for optimizing the batch conditions to methylene blue biosorption on yellow passion fruit and mandarin peels. *Dyes and Pigments*, 72(2), 256–266.
- Pérez-Rodríguez, N., Moreira, C. D., Torrado Agrasar, A., & Domínguez, J. M. (2016). Feruloyl esterase production by *Aspergillus terreus* CECT 2808 and subsequent application to enzymatic hydrolysis. *Enzyme and Microbial Technology*, 91, 52–58.

- Pettit, R. K. (2009). Mixed fermentation for natural product drug discovery. *Applied Microbiology and Biotechnology*, 83(1), 19-25.
- Pires, E. J., Ruiz, H. A., Teixeira, J. A., & Vicente, A. A. (2012). A new approach on brewer's spent grains treatment and potential use as lignocellulosic yeast cells carriers. *Journal of Agricultural and Food Chemistry*, 60(23), 5994–5999.
- Prasad, K. N., Kong, K. W., Ramanan, R. N., Azlan, A., & Ismail, A. (2012). Determination and optimization of flavonoid and extract yield from brown mango using response surface methodology. *Separation Science and Technology*, 47(1), 73–80.
- Prasad, S., Singh, A., & Joshi, H. C. (2007). Ethanol as an alternative fuel from agricultural, industrial and urban residues. *Resources, Conservation and Recycling*, 50(1), 1-39.
- Ragauskas, A. J., Williams, C. K., Davison, B. H., Britovsek, G., Cairney, J., Eckert, C. A., ... Tschaplinski, T. (2006). The path forward for biofuels and biomaterials. *Science (New York, N.Y.)*, 311(5760), 484–489.
- Rajasekar, V., Murty, R. V., & Muthukumar, C. (2015). Development of a simple kinetic model and parameter estimation for biomass and nattoxinase production by *Bacillus subtilis* 1A752. *Austin Journal of Biotechnology and Bioengineering*, 2(1), 1-5.
- Rangel-Vazquez, N. A., Hernandez-Montoya, V., & Bonilla-Petriciolet, A. (2014). Spectroscopy analysis and applications of rice husk and gluten husk using computational chemistry. In V. K. Thakur (Ed.). *Lignocellulosic polymer composites: Processing, characterization, and properties*. USA: John Wiley & Sons.
- Rao, R. S. P., & Muralikrishna, G. (2006). Water soluble feruloyl arabinoxylans from rice and ragi: Changes upon malting and their consequence on antioxidant activity. *Phytochemistry*, 67(1), 91–99.
- Razak, M. I. M., Abas, N. M., Yaacob, N. J. A., Rodzi, S. N. A. M., Yusof, N. M., & Azidin, R. A. (2015). An overview of primary sector in Malaysia. *International Journal of Economics, Commerce and Management*, 3(2), 13.
- Rinne, M., Jaakkola, S., & Huhtanen, P. (1997). Grass maturity effects on cattle fed silage-based diets. 1. Organic matter digestion, rumen fermentation and nitrogen utilization. *Animal Feed Science and Technology*, 67(1), 1-17.

- Riser-Roberts, E. (1998). *Remediation of petroleum contaminated soils: biological, physical, and chemical processes*. Boca Raton: CRC Press.
- Robertson, J. A., P'Anson, K. J. A., Treimo, J., Faulds, C. B., Brocklehurst, T. F., Eijssink, V. G. H., & Waldron, K. W. (2010). Profiling brewers' spent grain for composition and microbial ecology at the site of production. *LWT - Food Science and Technology*, 43(6), 890–896.
- Rodríguez Porcel, E. M., Casas López, J. L., Sánchez Pérez, J. A., Fernández Sevilla, J. M., & Chisti, Y. (2005). Effects of pellet morphology on broth rheology in fermentations of *Aspergillus terreus*. *Biochemical Engineering Journal*, 26(2–3), 139–144.
- Rosa, N. N., Dufour, C., Lullien-Pellerin, V., & Micard, V. (2013). Exposure or release of ferulic acid from wheat aleurone: impact on its antioxidant capacity. *Food chemistry*, 141(3), 2355-2362.
- Russell, W. R., Labat, A., Scobbie, L., Duncan, G. J., & Duthie, G. G. (2009). Phenolic acid content of fruits commonly consumed and locally produced in Scotland. *Food Chemistry*, 115(1), 100–104.
- Khan, T. S., & Mubeen, U. (2012). Wheat straw: A pragmatic overview. *Current Research Journal of Biological Sciences*, 4(6), 673–675.
- Salleh, N. H. M., Daud, M. Z. M., Arbain, D., Ahmad, M. S., & Ismail, K. S. K. (2011). Optimization of alkaline hydrolysis of paddy straw for ferulic acid extraction. *Industrial Crops and Products*, 34(3), 1635–1640.
- Santos, L. O., Gonzales, T. A., Ubeda, B. T., & Monte Alegre, R. (2007). Influence of culture conditions on glutathione production by *Saccharomyces cerevisiae*. *Applied Microbiology and Biotechnology*, 77, 763–769.
- Saqib, M., Mumtaz, M. W., Mahmood, A., & Abdullah, M. I. (2012). Optimized biodiesel production and environmental assessment of produced biodiesel. *Biotechnology and Bioprocess Engineering*, 17(3), 617–623.
- Saraiva, A. B., Pacheco, E. B. A. V., Visconte, L. L. Y., & Pereira, E. (2012). Potentials for utilization of post-fiber extraction waste from tropical fruit production in Brazil – the example of banana. *International Journal of Environment and Bioenergy*, 4(2), 101–119.
- Sarangi, P. K., & Sahoo, H.P. (2010). Ferulic Acid production from wheat bran using *Staphylococcus aureus*. *New York Science Journal*, 4, 79-81.

- Saravanan, K., & Aradhya, S. M. (2011). Polyphenols of pseudostem of different banana cultivars and their antioxidant activities. *Journal of Agricultural and Food Chemistry*, 59(8), 3613–3623.
- Saulnier, L., & Thibault, J. F. (1999). Ferulic acid and diferulic acids as components of sugar-beet pectins and maize bran heteroxylans. *Journal of the Science of Food and Agriculture*. 79(3), 396–402.
- Shah, M., & Pathak, K., (2010). Development and statistical optimization of solid lipid nanoparticles of simvastatin by using 2<sup>3</sup> full-factorial design. *AAPS PharmSci Tech.*, 11, 489-496.
- Shahidi, F., & Naczk, M. (2006). *Phenolics in food and nutraceuticals*. New York: CRC Press.
- Shahzadi, T., Anwar, Z., Iqbal, Z., Anjum, A., Aqil, T., Afzal, A., ... Irshad, M. (2014a). Induced production of exoglucanase, and  $\beta$ -Glucosidase from fungal co-culture of *T. viride* and *G. lucidum*. *Advances in Bioscience and Biotechnology*, 5(5), 426-433.
- Shahzadi, T., Mehmood, S., Irshad, M., Anwar, Z., Afroz, A., Zeeshan, N., ... Sughra, K. (2014b). Advances in lignocellulosic biotechnology: A brief review on lignocellulosic biomass and cellulases. *Advances in Bioscience & Biotechnology*, 5, 246–251.
- Shalin, T., Sindhu, R., Binod, P., Soccol, C. R., & Pandey, A. (2014). Mixed cultures fermentation for the production of poly- $\beta$ -hydroxybutyrate. *Brazilian Archives of Biology and Technology*, 57(5), 644-652.
- Sharp, R. J., Riley, P. W., & White, D. (1992). Heterotrophic thermophilic *Bacilli*. In J. K. Kristjansson (Ed.). *Thermophilic bacteria*. Boca Raton: CRC Press.
- Shimizu, H., Mizuguchi, T., Tanaka, E., & Shioya, S. (1999). Nisin production by a mixed-culture system consisting of *Lactococcus lactis* and *Kluyveromyces marxianus*. *Applied and Environmental Microbiology*, 65(7), 3134-3141.
- Shin, H. D., & Chen, R. R. (2006). Production and characterization of a type B feruloyl esterase from *Fusarium proliferatum* NRRL 26517. *Enzyme and Microbial Technology*, 38(3–4), 478–485.
- Shuler, M. L., & Kargi, F. (2010). *Bioprocess engineering basic concepts*. United Sate, Westford: Pearson Education Publisher.

- Silhavy, T. J., Kahne, D., & Walker, S. (2010). The bacterial cell envelope. *Cold Spring Harbor perspectives in biology*, 2(5), 1-16.
- Silva, J. P. A., Mussatto, S. I., & Roberto, I. C. (2010). The influence of initial xylose concentration, agitation, and aeration on ethanol production by *Pichia stipitis* from rice straw hemicellulosic hydrolysate. *Applied Biochemistry and Biotechnology*, 162(5), 1306–1315.
- Silveira, M. L. L., Furlan, S. A., & Ninow, J. L. (2008). Development of an alternative technology for the oyster mushroom production using liquid inoculum. *Food Science and Technology (Campinas)*, 28(4), 858-862.
- Silveira, J. E., Zazo, J. A., Pliego, G., Bidóia, E. D., & Moraes, P. B. (2014). Electrochemical oxidation of landfill leachate in a flow reactor: optimization using response surface methodology. *Environmental Science and Pollution Research*, 22(8), 5831–5841.
- Singh, B., Singh, J. P., Kaur, A., & Singh, N. (2016). Bioactive compounds in banana and their associated health benefits - A review. *Food Chemistry*. 206, 1-11.
- Srinivasan, M., Sudheer, A. R., & Menon, V. P. (2007). Ferulic acid: Therapeutic potential through its antioxidant property. *Journal of Clinical Biochemistry and Nutrition*, 40(2), 92-100.
- Stager, C. E., & Davis, J. (1992). Automated systems for identification of microorganisms. *Clinical Microbiology Review*, 5, 302-327.
- Strigul, N., Dette, H., & Melas, V. B. (2009). Environmental modelling & software a practical guide for optimal designs of experiments in the Monod model. *Environmental Modelling and Software*, 24, 1019–1026.
- Suhaila, Y. N., Ramanan, R. N., Rosfarizan, M., Latif, I. A., & Ariff, A. B. (2013). Optimization of parameters for improvement of phenol degradation by *Rhodococcus* UKMP-5M using response surface methodology. *Annals of microbiology*, 63(2), 513-521.
- Suli, E. (2014). *Numerical solution of ordinary differential equations*. Retrieved from <https://people.maths.ox.ac.uk/suli/nsodes.pdf>
- Süli, E., & Mayers, D. F. (2003). *An introduction to numerical analysis*. UK: Cambridge University Press.

- Sun, R., Sun, X. F., Wang, S. Q., Zhu, W., & Wang, X. Y. (2002). Ester and ether linkages between hydroxycinnamic acids and lignins from wheat, rice, rye, and barley straws, maize stems, and fast-growing poplar wood. *Industrial Crops and Products*, *15*(3), 179-188.
- Sutton, S. (2006). The gram stain. Retrieved from the Microbiology Network, Inc. website: <http://www.microbiol.org/resources/monographswhite-papers/the-gram-stain/>
- Suzuki, N., Ito, T., Hiroshima, K., Tokiwano, T., & Hashizume, K. (2016). Formation of ethyl ferulate from feruloylated oligosaccharide by transesterification of rice koji enzyme under sake mash conditions. *Journal of Bioscience and Bioengineering*, *121*(3), 281–285.
- Szwajgier, D., & Jakubczyk, A. (2011). Production of extracellular ferulic acid esterases by *Lactobacillus* strains using natural and synthetic carbon sources. *Acta Scientiarum Polonorum, Technologia Alimentaria*, *10*(3), 287–302.
- Tabaraki, R., & Nateghi, A. (2011). Optimization of ultrasonic-assisted extraction of natural antioxidants from rice bran using response surface methodology. *Ultrasonics Sonochemistry*, *18*(6), 1279–1286.
- Tesfaw, A., & Assefa, F. (2014). Co-culture: A great promising method in single cell protein production. *Biotechnology and Molecular Biology Reviews*, *9*(2), 12-20.
- Tilay, A., Bule, M., Kishenkumar, J., & Annapure, U. (2008). Preparation of ferulic acid from agricultural wastes: Its improved extraction and purification. *Journal of Agricultural and Food Chemistry*, *56*(17), 7644–7648.
- Torre, P., Aliakbarian, B., Rivas, B., Dominguez, J. M., & Converti, A. (2008). Release of ferulic acid from corn cobs by alkaline hydrolysis. *Biochemical Engineering Journal*, *40*(3), 500–506.
- Tripathi, S., Singh, S., Gangwar, A., Mishra, O. P., Chakrabarti, S. K., Bhardwaj, N. K., & Varadhan, R. (2013). Blending of banana stem with wheat straw and bagasse to enhance physical strength properties of paper. *Ippta*, *25*(2), 121–125.
- Tsuchiya, T., & Takasawa, M. (1975). Oryzanol, ferulic acid, and their derivatives as preservatives. *Japan. Kokai Tokkyo Koho JP*, *7*, 518–521.

- Ueno, A., Ito, Y., Yumoto, I., & Okuyama, H. (2007). Isolation and characterization of bacteria from soil contaminated with diesel oil and the possible use of these in autochthonous bioaugmentation. *World Journal of Microbiology and Biotechnology*, 23(12), 1739-1745.
- Uraji, M., Kimura, M., Inoue, Y., Kawakami, K., Kumagai, Y., Harazono, K., & Hatanaka, T. (2013). Enzymatic production of ferulic acid from defatted rice bran by using a combination of bacterial enzymes. *Applied Biochemistry and Biotechnology*, 171(5), 1085–1093.
- Van Soest, P. V., Robertson, J. B., & Lewis, B. A. (1991). Methods for dietary fiber, neutral detergent fiber, and nonstarch polysaccharides in relation to animal nutrition. *Journal of dairy science*, 74(10), 3583-3597.
- Vieira, F. C. S., & Nahas, E. (2005). Comparison of microbial numbers in soils by using various culture media and temperatures. *Microbiological Research*, 160(2), 197–202.
- Volm, T. G. 1997. *Isolation, purification, characterization of ferulic acid decarboxylase from Bacillus pumilus*. (Doctoral dissertation). Available from ProQuest Dissertations and Theses Global data base. (UMI No. 9731883)
- Walter, R. H. (1991). *The chemistry and technology of pectin*. Food Sci. Technol. San Diego, California: Academic Press, Inc.
- Wang, N. S. (2007). Experiment No. 9C–Measurements of cell biomass concentration. Retrieved from University of Maryland website: <http://terpconnect.umd.edu/~nsw/ench485/lab9c.htm>
- Wang, J., Sun, B., Cao, Y., Tian, Y., & Li, X. (2008). Optimisation of ultrasound-assisted extraction of phenolic compounds from wheat bran. *Food Chemistry*, 106(2), 804–810.
- Widdel, F. (2007). Theory and measurement of bacterial growth. *Grundpraktikum Mikrobiologie*, 4, 1-11.
- Williamson, K., & McCarty, P. L. (1976). A model of substrate utilization by bacterial films. *Journal (Water Pollution Control Federation)*, 48(1), 9-24.
- Witek-Krowiak, A., Chojnacka, K., Podstawczyk, D., Dawiec, A., & Pokomeda, K. (2014). Application of response surface methodology and artificial neural network methods in modelling and optimization of biosorption process. *Bioresource Technology*, 160, 150–160.

- Wragg, P., Randall, L., & Whatmore, A. M. (2014). Comparison of Biolog GEN III MicroStation semi-automated bacterial identification system with matrix-assisted laser desorption ionization-time of flight mass spectrometry and 16S ribosomal RNA gene sequencing for the identification of bacteria of veterinary interest. *Journal of microbiological methods*, *105*, 16-21.
- Wu, H., Li, H., Xue, Y., Luo, G., Gan, L., Liu, J., ... & Long, M. (2017). High efficiency co-production of ferulic acid and xylooligosaccharides from wheat bran by recombinant xylanase and feruloyl esterase. *Biochemical Engineering Journal*.
- Xiangli, F., Wei, W., Chen, Y., Jin, W., & Xu, N. (2008). Optimization of preparation conditions for polydimethylsiloxane (PDMS)/ceramic composite pervaporation membranes using response surface methodology. *Journal of Membrane Science*, *311*, 23-33.
- Xie, C. Y., Gu, Z. X., You, X., Liu, G., Tan, Y., & Zhang, H. (2010). Screening of edible mushrooms for release of ferulic acid from wheat bran by fermentation. *Enzyme and Microbial Technology*, *46*(2), 125–128.
- Xiros, C., Moukouli, M, Topakas, E., & Christakopoulos, P. (2009). Factors affecting ferulic acid release from brewer's spent grain by *Fusarium oxysporum* enzymatic system. *Bioresource Technology*, *100*(23), 5917–5921.
- Xue, Y., Wang, R., Zhang, J., Xu, C., & Sun, H. (2012). Production and some properties of the thermostable feruloyl esterase and xylanase from *Bacillus pumilus*. *African Journal of Biotechnology*. *11*(15), 3617-3622.
- Yagi, K., & Ohishi, N. (1979). Action of ferulic acid and its derivatives as antioxidants. *Journal of nutritional science and vitaminology*, *25*(2), 127–130.
- Yahiaoui, I., & Aissani-Benissad, F. (2010). Experimental design for copper cementation process in fixed bed reactor using two-level factorial design. *Arabian Journal of Chemistry*, *3*(3), 187–190.
- Yang, W., Meng, F., Peng, J., Han, P., Fang, F., Ma, L., & Cao, B. (2014). Isolation and identification of a cellulolytic bacterium from the tibetan pig's intestine and investigation of its cellulase production. *Electronic Journal of Biotechnology*, *17*(6), 262–267.
- Yao, J., Chen, Q. L., Shen, A. X., Cao, W., & Liu, Y. H. (2013). A novel feruloyl esterase from a soil metagenomic library with tannase activity. *Journal of Molecular Catalysis B: Enzymatic*, *95*, 55-61.

- Yi, S., Su, Y., Qi, B., Su, Z., & Wan, Y. (2010). Application of response surface methodology and central composite rotatable design in optimizing the preparation conditions of vinyltriethoxysilane modified silicalite/polydimethylsiloxane hybrid pervaporation membranes. *Separation and Purification Technology*, 71(2), 252–262.
- Yoshizawa, K., Komatsu, S., Takahashi, I., & Otsuka, K. (1970). Phenolic compounds in the fermented products: part I. Origin of ferulic acid in saké. *Agricultural and Biological Chemistry*, 34(2), 170-180.
- Yu, P., McKinnon, J. J., & Christensen, D. A. (2005a). Hydroxycinnamic acids and ferulic acid esterase in relation to biodegradation of complex plant cell walls. *Canadian journal of animal science*, 85(3), 255-267.
- Yu, P., McKinnon, J. J., & Christensen, D. A. (2005b). Improving the nutritional value of oat hulls for ruminant animals with pretreatment of a multienzyme cocktail: In vitro studies. *Journal of Animal Science*, 83(5), 1133–1141.
- Yuan, X., Liu, J., Zeng, G., Shi, J., Tong, J., & Huang, G. (2008). Optimization of conversion of waste rapeseed oil with high FFA to biodiesel using response surface methodology. *Renewable Energy*, 33, 1678-1684.
- Alam, M. Z., Fakhru'l-Razi, A., Abd-Aziz, S., & Molla, A. H. (2003). Optimization of compatible mixed cultures for liquid state bioconversion of municipal wastewater sludge. *Water, Air, and Soil Pollution*, 149(1), 113–126.
- Zhang, Z. Y., Pan, L. P., & Li, H. H. (2010). Isolation, identification and characterization of soil microbes which degrade phenolic allelochemicals. *Journal of Applied Microbiology*, 108(5), 1839–1849.
- Zhang, S. B., Wang, L., Liu, Y., Zhai, H. C., Cai, J. P., & Hu, Y.S. (2015). Expression of feruloyl esterase A from *Aspergillus terreus* and its application in biomass degradation. *Protein expression and purification*, 115, 153-157.
- Zhao, Z., & Moghadasian, M. H. (2008). Chemistry, natural sources, dietary intake and pharmacokinetic properties of ferulic acid: A review. *Food Chemistry*, 109(4), 691-702.
- Zhao, S., Yao, S., Ou, S., Lin, J., Wang, Y., Peng, X., ... Yu, B. (2014). Preparation of ferulic acid from corn bran: Its improved extraction and purification by membrane separation. *Food and Bioproducts Processing*, 92(3), 309–313.

- Zheng, D. Q., Wu, X. C., Tao, X. L., Wang, P. M., Li, P., Chi, X. Q., ... Zhao, Y. H. (2011). Screening and construction of *Saccharomyces cerevisiae* strains with improved multi-tolerance and bioethanol fermentation performance. *Bioresource Technology*, *102*(3), 3020–3027.
- Zheng, Y., Zhang, H., Wang, D., Gao, P., & Shan, A. (2014). Strain development and optimized fermentation conditions for blood meal using *Aspergillus niger* and *Aspergillus oryzae*. *Journal of Microbiological Methods*, *101*(1), 70–80.
- Zywicka, A., Peitler, D., Rakoczy, R., Konopacki, M., Kordas, M. & Fijałkowski, K. (2015). The effect of different agitation modes on bacterial cellulose synthesis by *Gluconacetobacter xylinus* strains. *Acta Scientiarum Polonorum Zootechnica*, *14*(1), 137-150.