

Estimation of Plasma Insulin and Endogenous Insulin Secretion in Critically Ill Patients Using ICING Model

Ummu K. Jamaludin¹, Fatanah M. Suhaimi², Normy Norfiza A. Razak³, Azrina M. Ralib⁴, Mohd Basri Mat Nor⁴

¹Human Engineering Focus Group (HEG), Universiti Malaysia Pahang, 26600 Pekan, Pahang, Malaysia

²Advanced Medical and Dental Institute, Universiti Sains Malaysia, Bertam, 13200 Kepala Batas, Pulau Pinang, Malaysia

³College of Engineering, UNITEN, KM7, Jalan Ikram UNITEN, 43000 Kajang, Selangor, Malaysia

⁴Department of Anaesthesiology & Intensive Care, Kuliyyah of Medicine, International Islamic University Malaysia, Bandar Indera Mahkota Campus, Jalan Sultan Ahmad Sha, 25200 Kuantan, Pahang, Malaysia

ABSTRACT

The objective of this study is to estimate total plasma insulin level and endogenous insulin secretion by using Intensive Control Insulin-Nutrition-Glucose (ICING) model and 90 critically ill patients' data from Hospital Tengku Ampuan Afzan, Kuantan. Integral-based method was applied to solve mathematical equations defined in ICING model to find critical parameters of insulin sensitivity (SI) and results of total endogenous insulin secretion and total plasma insulin level were presented in median and 95% confidence interval (CI). It is reported that the total median plasma insulin is 1.35×10^6 mU while $(6.59 \times 10^5, 2.79 \times 10^6)$ mU is in 95% CI, and the total median endogenous insulin secretion is 12.9% from the total median plasma insulin. The results elucidated the effectiveness of current practice via Intensive Insulin Infusion Therapy (IIT) and also suggest a further study on investigating the incretin mechanism which is strongly believed to contribute to the total plasma insulin level and help to simulate endogenous insulin secretion.

KEYWORDS: ICING Model, Endogenous Insulin, Exogenous Insulin, Plasma Insulin, Critically Ill