

REFERENCES

- Abdul Hamid, N. A. (2013). *Natural dye products business plan*, (Master's thesis). Universiti Malaysia PAhang, Gampang.
- Aljeboree, A. M., Alshirifi, A. N., & Alkaim, A. F. (2014). Kinetics and equilibrium study for the adsorption of textile dyes on coconut shell activated carbon. *Arabian Journal of Chemistry*, 1-13.
- Azami, M., Bahram, M., & Nouri, S. (2013). Central composite design for the optimization of removal of the azo dye, Methyl Red, from waste water using Fenton reaction. *Current Chemistry Letters*, 2(2): 57–68.
- Azeredo, H. M. C. (2009). Betalains: properties, sources, applications, and stability. *International Journal of Food Science & Technology*, 44(12): 2365–2376.
- Azimah, S. (2014). *Separation of Betacyanin from Hylocerus Polyrhizus peel using ultrafiltration membrane*, (Master's thesis). Universiti Malaysia Pahang.
- Baishya, D., Talukdar, J., & Sandhya, S. (2012). Cotton dyeing with natural dye extracted from flower of Bottlebrush (*Callistemon citrinus*). *Universal Journal of Environmental Research and Technology*, 2(5): 377–382.
- Beeby, J. L. (2006). The theory of desorption. *Critical Reviews in Solid State and Materials Science*, 7(2): 153–166.
- Bhute, K. H. P. & A. S. (2012). Plant based dyes and mordant : A Review. *Journal of Natural Products and Plant Resources*, 6(November): 649–664.
- Blainski, A., Lopes, G. C., & de Mello, J. C. P. (2013). Application and analysis of the folin ciocalteu method for the determination of the total phenolic content from *Limonium Brasiliense* L. *Molecules*, 18: 6852–6864.
- Bogdanov, M. G., & Svinyarov, I. (2013). Ionic liquid-supported solid – liquid extraction of bioactive alkaloids. II. Kinetics, modeling and mechanism of glaucine extraction from *Glaucium flavum*. *Separation and Purification Technology*, 103: 279–288.
- Boonsong, P., Laohakunjit, N., Kerdchoechuen, O., Thonburi, T., & Matta, F. B. (2011). Detection of pigments and natural colorants from Thai herbal plants for possible use as coloring dyes. *Hort Science*, 46(2): 265–272.

- Branch, B. (2010). Functional dyeing of wool with natural dye extracted from *Berberis vulgaris* wood and *Rumex Hymenosepolus* root as biomordant. *Iran Journal Chemical Chemical Engineering*, 29(3): 55–60.
- Camenzind, M., Freitag, R., & Furter, R. (2013). *Raw Silk and Spun Silk Quality and Measurement*.
- Cao, S., Liu, T., Jiang, Y., He, S., Harrison, D. K., & Joyce, D. C. (2012). The effects of host defence elicitors on betacyanin accumulation in *Amaranthus mangostanus* seedlings. *Food Chemistry*, 134(4): 1715–1718.
- Chan, L. S., Cheung, W. H., Allen, S. J., & McKay, G. (2012). Error analysis of adsorption isotherm models for acid dyes onto bamboo derived activated carbon. *Chinese Journal of Chemical Engineering*, 20(3): 535–542.
- Chandrasekhar, J., Sonika, G., Madhusudhan, M. C., & Raghavarao, K. S. M. S. (2015). Differential partitioning of betacyanins and betaxanthins employing aqueous two phase extraction. *Journal of Food Engineering*, 144: 156–163.
- Chavan, R. (2001). Indian textile industry-environmental issues. *Indian Journal of Fibre and Textile Research*, 26: 11–21.
- Chemat, F., Vian, M. A., & Cravoto, G. (2016). Green extraction of natural products : Concept and principles. *International Journal of Molecular Science*, 13(7): 8615–8627.
- Chowdry, S., Chakraborty, S., & Das Saha, P. (2013). Response surface optimization of a dynamic dye adsorption process: a case study of crystal violet adsorption onto NaOH-modified rice husk. 20: 1698–1705. doi: 10.1007/s11356-012-0989-7
- Chung, H., Kim, W., Park, J., Cho, J., Jeong, T., & Park, P. (2015). Application of Langmuir and Freundlich isotherms to predict adsorbate removal efficiency or required amount of adsorbent. *Journal of Industrial and Engineering Chemistry*, 28: 241–246.
- Cordeiro de Azeredo, H. M., Pereira, A. C., Rodrigues de Souza, A. C., Gouveia, S. T., & Barbosa Mendes, K. C. (2009). Study on efficiency of betacyanin extraction from red beetroots. *International Journal of Food Science & Technology*, 44(12): 2464–2469.
- Deka, A., Chougule, B. S., Parveen, A., & Lahan, J. P. (2014). Natural pigment betacyanin as tracking dye for gel electrophoresis. *Indian Journal of Natural Product and Resources*, 6(March): 23–26.

- Delgado-Vargas, F., Jimenez, A. R., & Paredes-Lopez, O. (2014). Natural pigments: carotenoids, anthocyanins, and betalains — Characteristics, biosynthesis, processing, and stability. *Critical Reviews in Food Science and Nutrition*, 40(3):173-289. doi: 10.1080/10408690091189257
- Denni, M., & Rt, S. (2012). Identification and characterization of a betacyanin pigment from the semi-arid plant *Aerva lanata* juss ex . Schultes. *International Journal of Research in Pharmaceutical and Biomedical Sciences*, 3(1): 212–214.
- Desta, M. B. (2013). Batch Sorption Experiments : Langmuir and Freundlich Isotherm studies for the adsorption of textile metal ions onto Teff Straw (*Eragrostis tef*) Agricultural Waste. *Journal of Thermodynamics*, 2013: 1-6. doi: 10.1155/2013/375830
- Dikshith, T. S. S. (2008). Tin and its compounds. *Hazardous Chemicals: Safety Management and Global Regulations* (pp. 388–390).New York. CRC Press.
- Elmorsi, T. M. (2011). Equilibrium isotherms and kinetic studies of removal of methylene blue dye by adsorption onto miswak leaves as a natural adsorbent. *Journal of Environmental Protection*, 2011(August): 817–827. doi: 10.4236/jep.2011.26093
- Fouda, I. M., Seisa, E. a., & Eid, F. M. (2015). Characterizations of the optical and structure properties of rayon acetate fibers due to different thermal conditions. *Optik - International Journal for Light and Electron Optics*, 126(2): 258–264. doi: 10.1016/j.ijleo.2014.08.169
- Gamal, A. M., Farha, S. A. A., Sallam, H. B., Mahmoud, G. E. A., & Ismail, L. F. M. (2010). Kinetic Study and Equilibrium Isotherm Analysis of Reactive Dyes Adsorption onto Cotton Fiber. *Nature and Science*, 8(11): 95–110.
- Garcia, E. R., Medina, R. L., Lozano, M. M., Perez, I. H., Valero, M. J., & Franco, M. M. (2014). Adsorption of azo-dye orange II from aqueous solutions using a metal-organic framework material: iron- benzenetricarboxylate. *Materials*,7: 8037–8057. doi:10.3390/ma7128037
- Gaurav, P., Kumar, J. N., Narendra, N., & Chatap, V. K. (2010). Bougainvillea glabra – A Natural Indicator. *Pharmacognosy Journal*, 2(5): 25–28.
- Ghasemi, E., Raofie, F., & Najafi, N. M. (2011). Application of response surface methodology and central composite design for the optimisation of supercritical fluid extraction of essential oils from *Myrtus communis* L . leaves. *Food Chemistry*, 126(3): 1449–1453.

- Ghosh, R. (2006). Overview of bioseparation engineering. *Principles of bioseparations engineering* (pp. 1-12). Singapore: World Scientific Publishing Co. Pte. Ltd.
- Grifoni, D., Bacci, L., Zipoli, G., Albanese, L., & Sabatini, F. (2011). The role of natural dyes in the UV protection of fabrics made of vegetable fibres. *Dyes and Pigments*, *91*(3): 279–285.
- Gupta, V., George, M., Joseph, L., Singhal, M., & Singh, H. P. (2009). Evaluation of antibacterial activity of *Bougainvillea glabra* 'snow white' and *Bougainvillea glabra* 'choicy'. *Journal of Chemical and Pharmaceutical Research*, *1*(1): 233–237. Retrieved from <http://www.jocpr.com/articles/evaluation-of-antibacterial-activity-of-bougainvillea-glabra-snow-white-and-bougainvillea-glabra-choicy.pdf>
- Ha, G., & Kim, J. (2016). Kinetic and thermodynamic characteristics of ultrasound-assisted extraction for recovery of paclitaxel from biomass. *Process Biochemistry*, *51*(2016): 1664–1673. doi: 10.1016/j.procbio.2016.08.012
- Habip, D., Dilek, K., Nigar, M., & Seyda, C. (2015). The effect of dyeing properties of fixing agent and plasma treatment on silk fabric dyed with natural dye extract obtained from *Sambucus Ebulus* L. plant. *Procedia- Social and Behavioral Sciences*, *195*(2015): 1609–1617. doi: 10.1016/j.sbspro.2015.06.201
- Haji, A. (2012). Antibacterial dyeing of wool with natural cationic dye using metal mordants. *Materials Science*, *18*(3): 267–270. doi: 10.5755/j01.ms.18.3.2437
- Hamid, N., Abdul Munaim, M. S., & Abu Seman, M. N. (2016). Regression analysis for the adsorption isotherms of betacyanin extracts from the dragon fruit peel onto the spun silk yarn. *International Journal of Engineering Technology and Sciences*, *6*(1): 44–53. doi: 10.15282/ijets.6.2016.1.7.1057
- Haque, N., Hannan, M. A., & Rana, M. (2015). Compatibility analysis of reactive dyes by exhaustion-fixation and adsorption isotherm on knitted cotton fabric. *Fashion and Textiles*, *2*(3): 1-12. doi: 10.1186/s40691-015-0026-9
- Hasan, M., Nayem, K. A., Yousuf, A., Anwarul, M., & Ghosh, N. C. (2015). Application of purified lawsone as natural dye on cotton and silk fabric. *Journal of Textiles*, *2015*: 1–8. doi: 10.1155/2015/932627
- Hassan, S. S. M., Awwad, N. S., & Aboterika, A. H. A. (2009). Removal of synthetic reactive dyes from textile wastewater by Sorel's cement. *Journal of Hazardous Materials*, *162*: 994–999. doi: 10.1016/j.jhazmat.2008.05.138
- Hemmatzadeh, R., & Mohammadi, A. (2013). Improving optical absorptivity of natural dyes for fabrication of efficient dye-sensitized solar cells. *Journal of Theoretical and Applied Physics*, *7*(57): 1-7.

- Hernandez, A. R., Estevez, M., Vargas, S., Quintanilla, F., & Rodriguez, R. (2011). New dye-sensitized solar cells obtained from extracted bracts of *Bougainvillea glabra* and *spectabilis* betalain pigments by different purification processes. *International Journal of Molecular Sciences*, *12*(9): 5565–76.
- Hosseini, M., Montazer, M., & Damerchely, R. (2013). Enhancing Dye-ability and Antibacterial Features of Silk through Pre-treatment with Chitosan. *Journal of Engineered Fibers and Fabrics*, *8*(3): 102–111.
- Hou, X., Chen, X., Cheng, Y., Xu, H., Chen, L., & Yang, Y. (2013). Dyeing and UV-protection properties of water extracts from orange peel. *Journal of Cleaner Production*, *52*(2013): 410–419.
- Ibrahim, U. K., Muhammad, I. I., & Salleh, R. M. (2011). The effect of pH on color behavior of *Brassica oleracea* anthocyanin. *Journal of Applied Sciences*, *11*(13): 2406–2410.
- Jerz, G., Wybraniec, S., Gebers, N., & Winterhalter, P. (2010). Target-guided separation of *Bougainvillea glabra* betacyanins by direct coupling of preparative ion-pair high-speed countercurrent chromatography and electrospray ionization mass-spectrometry. *Journal of Chromatography. A*, *1217*(27): 4544–4554.
- Jothi, D. (2008). Extraction of natural dyes from african marigold flower (*Tagetes Erecta* L). *AUTEX Research Journal*, *8*(2): 49–53. Retrieved from http://www.autexrj.com/cms/zalaczone_pliki/4-08-2.pdf
- Jung, J. Y., Choi, M. S., & Yang, J. K. (2013). Optimization of concentrated acid hydrolysis of waste paper using response surface methodology. *Journal of Korean Wood Science and Technology*, *41*(2): 87–99. doi: 10.5658/WOOD.2013.41.2.87
- Jyothi, S. N. & S. (2012). A comparative phytochemical analysis of *Bougainvillea glabra* choisy and calforina gold. *International Journal of Pharma and Bio Sciences* *3*(3): 247–250.
- Kalirajan, A., Mariselvam, R., Michael, J. S., Narayanan, K. R., Narayanan, G. A., & Singh, A. J. A. R. (2012). Utilization of *Bougainvillea glabra* for prepared natural colouring agent and biopesticides. *International Journal of Current Research*, *4*(9): 9–11.
- Kaur, M. (2014). *Adsorption study of betacyanin of fibre*. URP Thesis. Universiti Malaysia Pahang. Gambang.
- Kechi, A., Chavan, R. B., & Moeckel, R. (2013). Dye yield, color strength and dyeing properties of natural dyes extracted from ethiopian dye plants. *Textiles and Light Industrial Science and Technology*, *2*(3): 137–145.

- Khataee, A. R., Zarei, M., & Moradkhannejhad, L. (2010). Application of response surface methodology for optimization of azo dye removal by oxalate catalyzed photoelectro-Fenton process using carbon nanotube-PTFE cathode. *Desalination*, 258(2010): 112–119.
- Killedar, S., Pawar, A., Nadaf, S., Nale, A., Tamboli, U., & Pishawikar, S. (2014). Novel analytical method development for some amide group containing drugs using *Bougainvillea spectabilis* bract extracts. *Asian Pacific Journal of Tropical Medicine*, 7(Suppl 1): S560–S567.
- Kiurski, J., Adamovic, S., & Krstic, J. (2011). Adsorption efficiency of low-cost materials in the removal of Zn (II) ions from printing developer. *ACTA Technica Corviniensis - Bulletin of Engineering*, 4(2011): 61–66.
- Koleva, V., & Simeonov, E. (2014). Solid liquid extraction of phenolic and flavonoid compounds from *Cotinus coggygia* and concentration by nanofiltration. *Chemical, Biochemical Engineering*, 28(4): 545–551. doi: 10.15255/CABEQ.2014.2006
- Kreuzer, H. J. (1988). Theory of Desorption Kinetics. *Chemistry and Physic of Solid Surfaces VII*, 259–260.
- Kugler, F., Stintzing, F. C., & Carle, R. (2007). Characterisation of betalain patterns of differently coloured inflorescences from *Gomphrena globosa* L. and *Bougainvillea* sp. by HPLC-DAD-ESI-MSⁿ. *Analytical and Bioanalytical Chemistry*, 387(2): 637–648.
- Lee, J., Kang, M. H., Lee, K., & Lee, Y. (2013). Characterization of natural dyes and traditional korean silk fabric by surface analytical techniques. *Materials*, 2013(6) 2007–2025. doi: 10.3390/ma6052007
- Li, F., Yang, Q., Hong, C., & Zhong, Z. (2010). Color analysis on natural polyphenolic dyed cotton cellulose fibers. *Journal of Fiber Bioengineering and Informatics* 3(2): 100–105. doi:10.3993/jfbi09201007
- Mahmoud, A. S., Ghaly, A. E., & Brooks, S. L. (2007). Influence of temperature and pH on the stability and colorimetric measurement of textile dyes. *American Journal of Biotechnology and Biochemistry*, 3(1): 33–41.
- Mai, S. D., Thuan, T. N., Trinh, L. T. N., Trinh, T. T. N., & Ngo, T. X. (2012). Survey the Betacyanin extraction from the flower of purple. *Journal Food Science and Engineering*, 2:(2012): 254

- Manjunath, M. S., & Sannappa, J. (2011). Variation of amino acids in some biological and pharmaceutical sample using FT-IR study. *International Journal of Pharmacy and Technology*, 3(1): 1946–1951.
- Maran, J. P., Priya, B., & Nivetha, C. V. (2015). Optimization of ultrasound-assisted extraction of natural pigments from *Bougainvillea glabra* flowers. *Industrial Crops and Products*, 63(2015): 182–189. doi: 10.1016/j.indcrop.2014.09.059
- Martinez, M. P., & Tena, M. T. (2011). Focused ultrasound solid-liquid extraction and selective pressurised liquid extraction to determine bisphenol A and alkylphenols in sewage sludge by gas chromatography-mass spectrometry. *Journal of Separation Science*, 34: 2513-2522. doi: 10.1002/jssc.201100159
- Miraboutalebi, S. M., Nikouzad, S. K., Peydayesh, M., Allahgholi, N., Vafajoo, L., & McKay, G. (2017). Methylene blue adsorption via maize silk powder: Kinetic, equilibrium, thermodynamic studies and residual error analysis. *Process Safety and Environmental Protection*, 106: 191–202.
- Mirjalili, M., & Karimi, L. (2013). Extraction and characterization of natural dye from green walnut shells and its use in dyeing polyamide: focus on antibacterial properties. *Journal of Chemistry*, 2013:1-9
- Mishra, N., Joshi, S., Tandon, V. L., & Munjal, A. (2009). Evaluation of anti-fertility potential of aqueous extract of *Bougainvillea* in swiss albino mice. *International Journal of Pharmaceutical Sciences and Drug Research*, 1(1): 19-23
- Moghaddam, S. S., Moghaddam, M. R. A., & Arami, M. (2010). Coagulation / flocculation process for dye removal using sludge from water treatment plant: Optimization through response surface methodology. *Journal of Hazardous Materials*, 175(2010): 651–657. doi: 10.1016/j.jhazmat.2009.10.058
- Munipan, M. A. (2013). *Effect Of Enzyme Concentration And Temperature On Viscosity And Betacyanin Content From Pitaya Waste Extract*. URP Thesis. Universiti Malaysia Pahang. Gambang.
- Naderi, N., Ghazali, H. M., Shobirin, A., & Hussin, M. (2012). Characterization and quantification of dragon fruit (*Hylocereus polyrhizus*) betacyanin pigments extracted by two procedures. *Pertanika Journal of Tropical Agricultural Science* 35(1): 33–40.
- Narayan, S., Kumar, A., Ritesh, S. K., Sharmila, G., & Muthukumar, C. (2013). Extraction optimization and characterization of water soluble red purple pigment from floral bracts of *Bougainvillea glabra*. *Arabian Journal of Chemistry*. doi: 10.1016/j.arabjc.2013.07.047 (in press)

- Noda, N., & Adachi, T. (2000). Induction of betacyanin synthesis and pigment accumulation in cell suspension cultures of *Portulaca*. *Plant Biotechnology*, 17(1): 27-34.doi: 10.5511/plantbiotechnology.17.27
- Norasiha, H. (2011). *Inclusion complex formation between natural dye extracted from pitaya fruit skin and beta -cyclodextrin:kinetic and thermodynamic studies*. Universiti Malaysia Pahang. Gambang.
- Orsini, F., Pasquier, A. Du, Beaudoin, B., & Tarascon, J. M. (1998). In situ scanning electron microscopy (SEM) observation of interfaces within plastic lithium batteries. *Journal of Power Sources*, 76(1998): 19–29.
- Owen, T. (2000). Fundamentals of modern UV-visible spectroscopy. Retrieved from https://www.agilent.com/cs/library/primers/Public/59801397_020660.pdf
- Padamwar, M. N., & Pawar, A. P. (2004). Silk sericin and its applications : A review. *Journal of Scientific and Industrial Research*, 63(April): 323–329.
- Patil, P. D., Rao, C. R., Wasif, A. I., & Nagla, J. R. (2014). Taguchi optimization for efficient extraction of a natural dye from *Bougainvillea glabra* bracts. *International Journal of Research in Engineering and Technology*, 3(4): 23–30.
- Phebe, D., Chew, M. K., Suraini, A. A., Lai, O. M., & Janna, O. A. (2009). Red-fleshed pitaya (*Hylocereus polyrhizus*) fruit colour and betacyanin content depend on maturity. *International Food Research Journal*, 242(16): 233–242.
- Pisitsak, P., Hutakamol, J., Jeenapak, S., & Wanmanee, P. (2016). Natural dyeing of cotton with *Xylocarpus granatum* bark extract : dyeing , fastness , and ultraviolet protection properties. *Fibers and Polymers 2016*, 17(4): 560–568.
- Pradesh, A. (2014). Mass transfer enhancement through optimized extraction of a natural dye from *Bougainvillea glabra* Juss . bracts. *Indian Journal of Natural Products and Resources*, 5(4): 332–337.
- Prusty, A. K., Das, T., Nayak, A., & Das, N. B. (2010). Colourimetric analysis and antimicrobial study of natural dyes and dyed silk. *Journal of Cleaner Production*, 18(16–17): 1750–1756.
- Radojkovic, M., Zekovic, Z., Jokic, S., Vidovic, S., Lepojevi, Z., & Milosevic, S. (2012). Optimization of Solid-liquid extraction of antioxidants from black Mulberry leaves by response surface methodology. *Food Technology Biotechnology*, 50(2): 167-176.

- Rajendran, R., Balakumar, C., Kalaivani, J., & Sivakumar, R. (2011). Dyeability and antimicrobial properties of cotton fabrics finished with *Punica Granatum* extracts. *Journal of Textile and Apparel, Technology and Management*, 7(2): 1–12.
- Raymond H. M., Douglas C. M., & Christine, M. A. C. (2009). *Response Surface Methodology: Process and Product Optimization Using Designed Experiments, 3rd Edition*. John Wiley and Sons.
- Reshmi, S. K., Aravindhana, K. M., & Devi, P. S. (2012). The effect of light, temperature, pH on stability of betacyanin pigments in *Basella Alba* fruit. *Asian Journal of Pharmaceutical and Clinical Research*, 5(4): 5–8.
- Ristić, N., & Ristić, I. (2012). Cationic modification of cotton fabrics and reactive dyeing characteristics. *Journal of Engineered Fibers and Fabrics*, 7(4): 113–121.
- Rossi, T., Silva, P. M. S., Moura, L. F. De, Araújo, M. C., Brito, J. O., & Freeman, H. S. (2016). Waste from eucalyptus wood steaming as a natural dye source for textile fibers. *Journal of Cleaner Production*, 1–8. doi:10.1016/j.jclepro.2016.12.109
- Rungruangkitkrai, N., & Mongkholrattanasit, R. (2012). Eco-Friendly of textiles dyeing and printing with natural dyes. *Proceedings of International Conference: Textiles & Fashion 2012*, Thailand, 1–17. Retrieved from <http://textileconference.rmutp.ac.th/wp-content/uploads/2012/10/010-Eco-Friendly-of-Textiles-Dyeing-and-Print-with-Natural-Dyes.pdf>
- Sahraei, R., Farmany, A., & Mortazavi, S. S. (2013). A nanosilver-based spectrophotometry method for sensitive determination of tartrazine in food samples. *Food Chemistry Journal*, 138(2013): 1239–1242.
- Samanta, A. K., & Agarwal, P. (2009). Application of natural dyes on textiles. *Indian Journal of Fibre & Textile Research*, 34(December): 384–399.
- Saxena, S., & Raja, A. S. M. (2014). *Natural dyes: sources, chemistry, application and sustainability issues*. (S. S. Muthu, Ed.). Singapore: Springer Singapore.
- Shah, B., Shah, A., & Shah, P. (2011). Sorption isotherms and column separation of Cu (II) And Zn (II) using ortho substituted benzoic acid chelating resins. *Archives of Applied Science Research*, 3(3): 327–341.
- Shahid, M., Ull Islam, S., & Mohammad, F. (2013). Recent advancements in natural dye applications : a review. *Journal of Cleaner Production*, 53: 310–331.

- Shoba, V., Krishnapriya, K., Boopathy Raja, A., Elanchezhiyan, C., & S. (2013). Antimicrobial activity of *Bougainvillea glabra* flower extract doped mgs nanoparticles on *Escherichia coli*, *Salmonella typhi* and *Pseudomonas aeruginosa*. (1991): 2006–2009.
- Singh, H. B., & Bharati, K. A. (2015). *Handbook of Natural Dyes and Pigments* (pp. 1–3).
- Singh, R., Jain, A., Panwar, S., Gupta, D., & Khare, S. K. (2004). Antimicrobial activity of some natural dyes. *Dyes and Pigments*, 5–8.(in press) Retrieved from <https://pdfs.semanticscholar.org/73eb/f013f994d2956bb72d53b4004da5b7e838c2.pdf>
- Sinha, K., Chowdhury, S., Das, P., & Datta, S. (2013). Modeling of microwave-assisted extraction of natural dye from seeds of *Bixa orellana* (Annatto) using response surface methodology (RSM) and artificial neural network (ANN). *Industrial Crops & Products*, 41(2013): 165–171. doi: 10.1016/j.indcrop.2012.04.004
- Sinha, K., Das, P., & Datta, S. (2012). Dyes and pigments extraction of natural dye from petals of flame of forest (*Butea monosperma*) flower : Process optimization using response surface methodology (RSM). *Dyes and Pigments*, 94(2): 212–216.
- Siva, R. (2015). Status of natural dyes and dye-yielding plants in India Status of natural dyes and dye-yielding plants in India. *Current Science*, 92(7): 916–925.
- Sreekanth, D., Arunasree, M. K., Roy, K. R., Chandramohan Reddy, T., Reddy, G. V., & Reddanna, P. (2007). Betanin a betacyanin pigment purified from fruits of *Opuntia ficus-indica* induces apoptosis in human chronic myeloid leukemia Cell line-K562. *Phytomedicine: International Journal of Phytotherapy and Phytopharmacology*, 14(11): 739–46.
- Stamatopoulos, K., Chatzilazarou, A., & Katsoyannos, E. (2014). Optimization of multistage extraction of olive leaves for recovery of phenolic compounds at moderated temperatures and short extraction times. *Foods*, 3: 66–81. doi: 10.3390/foods3010066
- Stintzing, F. C., & Schieber, A. (2003). Evaluation of colour properties and chemical quality parameters of cactus juices. *European Food Research and Technology*, (216): 303–311. doi: 10.1007/s00217-002-0657-0
- Strack, D., Vogt, T., & Schliemann, W. (2003). Recent advances in betalain research. *Phytochemistry* (pp. 247–269). doi: 10.1016/S0031-9422(02)00564-2
- Vankar, P. S. (2000). *Chemistry of Natural Dyes*, (October): 73–80.

- Vinod, K. N., & Puttaswamy. (2010). Kinetic and adsorption studies of Indian siris (*Albizia lebbek*) natural dye on silk. *Indian Journal of Fibre & Textile Research*, 35(June): 159–163.
- Wang, X., Wu, Q., Wu, Y., Chen, G., Yue, W., & Liang, Q. (2012). Response surface optimized ultrasonic-assisted extraction of flavonoids from *Sparganii Rhizoma* and evaluation of their in vitro antioxidant activities. *Molecules*, 17: 6769–6783. doi: 10.3390/molecules17066769
- Wanyama, P. A. G., Kiremire, B. T., Ogowok, P., & Murumu, J. S. (2010). The effect of different mordants on strength and stability of colour produced from selected dye-yielding plants in Uganda. *International Archieve of Applied Sciences and Technology*, 1(2): 81–92.
- Wen, X., Yang, Q., Yan, Z., & Deng, Q. (2011). Determination of cadmium and copper in water and food samples by dispersive liquid – liquid microextraction combined with UV – vis spectrophotometry. *Microchemical Journal journal*, 97: 249–254.
- Wybraniec, S., Jerz, G., Gebers, N., & Winterhalter, P. (2010). Ion-pair high-speed countercurrent chromatography in fractionation of a high-molecular weight variation of acyl-oligosaccharide linked betacyanins from purple bracts of *Bougainvillea glabra*. *Journal of Chromatography. B, Analytical Technologies in the Biomedical and Life Sciences*, 878(5–6): 538–50.
- Wybraniec, S., Stalica, P., Jerz, G., Klose, B., Gebers, N., Winterhalter, P., Sporna, A., Szaleniec, M., & Mizrahi, Y. (2009). Separation of polar betalain pigments from cacti fruits of *Hylocereus polyrhizus* by ion-pair high-speed countercurrent chromatography. *Journal of Chromatography A*, 1216(41): 6890–6899. doi: 10.1016/j.chroma.2009.08.035
- Xu, Q., Shen, Y., Wang, H., Zhang, N., Xu, S., & Zhang, L. (2013). Application of response surface methodology to optimise extraction of flavonoids from *fructus sophorae*. *Food Chemistry*, 138(4): 2122–2129. doi: 10.1016/j.foodchem.2012.11.099
- Zhang, H., Low, W. P., & Lee, H. K. (2012). Evaluation of sulfonated graphene sheets as sorbent for micro - solid - phase extraction combined with gas chromatography – mass spectrometry. *Journal of Chromatography A*, 1233: 16–21.
- Zheng, Y., & Chow, A. H. L. (2009). Production and characterization of a spray-dried complex Spray-dried hydroxypropyl- β -cyclodextrin/quercetin complex. *Drug Development And Industrial Pharmacy*, 35(6): 727-734. doi: 10.1080/03639040802526805

- Zhu, Q. Y., Zhang, Q. Y., & Cao, J. (2016). Cyclodextrin - assisted liquid - solid extraction for determination of the composition of jujube fruit using ultrahigh performance liquid chromatography with electrochemical detection and quadrupole time-of- flight tandem mass spectrometry. *Food Chemistry*, 213(2016): 485–493.
- Zhu, Q. yao, Zhang, Q. yun, Cao, J., Cao, W., Xu, J. jing, & Peng, L. qing. (2016). Cyclodextrin-assisted liquid-solid extraction for determination of the composition of jujube fruit using ultrahigh performance liquid chromatography with electrochemical detection and quadrupole time-of-flight tandem mass spectrometry. *Food Chemistry*, 213: 485–493.
- Zularisam, A. W., Ismail, A. F., Salim, M. R., Sakinah, M., & Matsuura, T. (2009). Application of coagulation – ultrafiltration hybrid process for drinking water treatment: Optimization of operating conditions using experimental design. *Separation and Purification Technology*, 65: 193–210.