## Biosorption of Chromium (VI) Ions using Sustainable Eggshell Impregnated Pandanus amaryllifolius Roxb. Adsorbent

## Noor Yahida Yahya 1, Chan Jia Xin 2, Norzita Ngadi 2,\*

 Faculty of Civil Engineering Technology, Universiti Malaysia Pahang, Lebuharya Tun Razak, 26300 Gambang Kuantan, Pahang Darul Makmur, Malaysia
School of Chemical and Energy Engineering, Faculty of Engineering, Universiti Teknologi Malaysia, 81310 UTM Skudai, Johor Darul Takzim, Malaysia
\*Corresponding Author's email: <u>norzita@cheme.utm.my</u>

## Abstract:

The conventional methods used nowadays is cost inefficient and non-ecofriendly. Thus, biosorption using waste was proposed in this study. The aim of this study is to remove chromium (VI) ions using raw eggshell and hybrid eggshell-pandan biosorbent. The result obtained showed that hybrid sorbent had higher efficiency in removal of chromium (VI) ions. The experimental results had indicated that the optimum conditions for the raw eggshell sorbent to achieve the optimum removal percentage of 17.78% were 140 minutes of contact time, 0.5 g of sorbent, room temperature, pH 5 and 0.8 mg/L solution. While for hybrid sorbent, the optimum removal percentage achieved 37.88% when the contact time is 140 minutes, with 0.5 g dosage, 50°C, pH 4 and 0.8 mg/L solution. The study showed that the adsorption of Cr (VI) ions onto the hybrid sorbent follows pseudo-second-order kinetic model of Legergren, Freundlich isotherm model and the thermodynamic study showed negative values of Gibb's energy, positive value of enthalpy (16.68 KJ/mol) and entropy change (0.0473 KJ/mol). In short, hybrid eggshell-pandan sorbent is a good biosorbent to removal Cr (VI) ions from solution.

Keywords: Eggshell; Pandan; adsorption; Cr (VI) Ions; Biosorbent

## Acknowledgement

This study was supported by FRGS Research Grant received by one of the author; Norzita Ngadi, Universiti Teknologi Malaysia and RDU Research Grant received by Noor Yahida Yahya, Universiti Malaysia Pahang No. vot RDU 1803167.