

Simultaneous removal of toxic ammonia and lettuce cultivation in aquaponic system using microwave pyrolysis biochar

Man Huan Su^{a,b,1}, Elfini Azwar^{b,1}, Ya Feng Yang^{a,1}, Christian Sonne^{c,1}, Peter Nai Yuh Yek^{b,d}, Rock Keey Liew^e, Chin Kui Cheng^f, Pau Loke Show^g, Su Shiung Lam^{b,a}

^a Henan Province Engineering Research Center for Biomass Value-added Products, School of Forestry, Henan Agricultural University, Zhengzhou, 450002, China

^b Pyrolysis Technology Research Group, Institute of Tropical Aquaculture and Fisheries (AKUATROP), Universiti Malaysia Terengganu, 21030, Kuala Nerus, Terengganu, Malaysia

^c Aarhus University, Department of Bioscience, Arctic Research Centre (ARC), Frederiksborgvej 399, PO Box 358, DK-4000, Roskilde, Denmark

^d School of Engineering and Technology, University College of Technology Sarawak, Lot 88, Persiaran Brooke, 96000 Sibu, Sarawak, Malaysia

^e NV WESTERN PLT, No. 208B, Jalan Macalister, Georgetown, Pulau Pinang 10400, Malaysia

^f Faculty of Chemical & Natural Resources Engineering, Lebuhraya Tun Razak, Universiti Malaysia Pahang, Gambang Kuantan, Pahang 26300, Malaysia

^g Department of Chemical and Environmental Engineering, Faculty of Science and Engineering, University of Nottingham Malaysia, Jalan Broga, 43500 Semenyih, Selangor Darul Ehsan, Malaysia

ABSTRACT

This study examined an aquaponic approach of circulating water containing ammonia excretions from African catfish grown in an aquaculture tank for bacterial conversion into nitrates, which then acted as a nutrient substance to cultivate lettuce in hydroponic tank. We found that microwave pyrolysis biochar (450 g) having microporous (1.803 nm) and high BET surface area (419 m²/g) was suitable for use as biological carrier to grow nitrifying bacteria (63 g of biofilm mass) that treated the water quality through removing the ammonia (67%) and total suspended solids (68%), resulting in low concentration of remaining ammonia (0.42 mg/L) and total suspended solid (59.40 mg/L). It also increased the pH (6.8), converted the ammonia into nitrate (29.7 mg/L), and increased the nitrogen uptake by the lettuce (110 mg of nitrogen per plant), resulting in higher growth in lettuce (0.0562 %/day) while maintaining BOD₅ level (3.94 mg/L) at acceptable level and 100% of catfish survival rate. Our results demonstrated that microwave pyrolysis biochar can be a promising solution for growing nitrifying bacteria in aquaponic system for simultaneous toxic ammonia remediation and generation of nitrate for growing vegetable in aquaculture industry.

KEYWORDS

Aquaponics; Pyrolysis; Biochar; Ammonia; Lettuce

ACKNOWLEDGEMENT

The authors thank the financial and technical support provided by Henan Agricultural University, Universiti Malaysia Terengganu and University College of Technology Sarawak. The authors acknowledge the Ministry of Education Malaysia for the financial support under Fundamental Research Grant Scheme (Vot 59512), and Universiti Malaysia Terengganu under Golden Goose Research Grant Scheme (Vot 55191).