

**SOLID STATE FERMENTATION OF *Monascus purpureus* IN STIRRED DRUM BIOREACTOR FOR PRODUCTION OF RED PIGMENT FROM OIL PALM FROND**

**MOHAMAD AL AAMIN BIN RAZALI**

**MASTER OF ENGINEERING (BIO-PROCESS)**

**UNIVERSITI MALAYSIA PAHANG**



### **SUPERVISOR'S DECLARATION**

I hereby declare that I have checked this thesis and in my opinion, this thesis is adequate in terms of scope and quality for the award of the degree of Master of Engineering (Bio-Process).

---

Full Name : DR FARHAN BINTI MOHD SAID

Position : SENIOR LECTURER

Date :



## STUDENT'S DECLARATION

I hereby declare that the work in this thesis is based on my original work except for quotations and citation which have been duly acknowledged. I also declare that it has not been previously or concurrently submitted for any other degree at Universiti Malaysia Pahang or any other institutions.

---

Full Name : MOHAMAD AL AAMIN BIN RAZALI

ID Number : MKB 13005

Date :

SOLID STATE FERMENTATION OF *Monascus purpureus* IN  
STIRRED DRUM BIOREACTOR FOR PRODUCTION OF  
RED PIGMENT FROM OIL PALM FROND

MOHAMAD AL AAMIN BIN RAZALI

Thesis submitted in fulfilment of the requirements  
for the award of the degree of  
Master of Engineering  
(Bio-Process)

Faculty of Chemical and Natural Resources Engineering  
UNIVERSITI MALAYSIA PAHANG

SEPTEMBER 2019

## **ACKNOWLEDGEMENT**

All praise to Allah S.W.T who has showed me with kindness and affection during the course of my study that I cannot adequately thank for. His endless grace and love have provided me with the strength to finish the study.

I would like to express my greatest gratitude to my supervisor, Dr Farhan binti Mohd Said for her higher degree of patience, ideas, encouragement, guidance, suggestions, and support in accomplishing this research. She constantly captivated me with her outstanding professional conduct and her strong conviction for science. I am very appreciating her consistent support from the first day I applied to graduate programme to these concluding moments. I am truly grateful for her advanced vision about my training in science, her commitment to my forthcoming career and her tolerance of my countless mistakes. Thank you for not give up with me.

My sincere thanks to go to all my lab mates especially Dr. Abu Hannifa and members of the staff of Faculty of Chemical & Natural Resources Engineering who helped me in many ways and made my day at UMP comfortable and unforgettable.

Finally, a special appreciation and deepest gratitude to my lovely superhero wife, Nor Farhana, for her love, dream and sacrifice throughout this journey. Without her hard work and sacrifice, I can say that the existence of this thesis is impossible at all. Last but not least, thanks to my parents, Che Jah Shafie and Razali MD. Salleh, family members and friends who shared their support. I cannot find the appropriate words that could properly describe my appreciation for their devotion, support and faith in me to successfully completing this study.

## **ABSTRAK**

Disebabkan terdapat pelbagai keprihatinan mengenai kesan pewarna sintetik, pengeluaran pewarna semulajadi secara besar-besaran menjadi alternatif yang berpotensi dalam memenuhi permintaan pewarna dalam industri makanan. Kajian ini memberi tumpuan kepada keupayaan dram bioreaktor untuk menghasilkan pewarna merah menggunakan *Monascus purpureus* FTC 5357 dan pelepas kelapa sawit (OPF) sebagai substrat, melalui proses fermentasi secara pejal. Terdapat tujuh parameter yang terlibat iaitu kandungan lembapan awal (% v/w), saiz inokulum (spora/mL), kepekatan peptone (% w/w), pH awal, kadar pengudaraan (vvm) program agitasi (kitaran/hari) dan kapasiti muatan (% v/v) telah diuji untuk menentukan kadar penghasilan pigmen merah, biomas dan penggunaan glukosa. Tiga respon, iaitu pengeluaran pigmen merah, biomas, dan penggunaan glukosa, untuk mengenal pasti faktor-faktor penting yang mempengaruhi proses fermentasi secara pejal. Teknik satu faktor pada satu masa (OFAT) dan kajian saringan faktorial rekaan 2 peringkat telah digunakan dalam kajian ini. Dalam kajian OFAT, pengeluaran pigmen tertinggi ditunjukkan pada kadar pengudaraan 1 vvm (6.09 AU/g) dan pengeluaran terendah menunjukkan pada 45% kandungan lembapan awal (0.1 AU/g). Dalam dua peringkat reka bentuk faktorial, pengeluaran maksimum pigmen adalah 71.86 AU/g dengan menggabungkan 75% v/w kandungan lembapan awal,  $10^8$  spora/mL saiz inokulum, pH awal 4, kadar pengudaraan 1.21 vvm, kapasiti muatan 35% v/v dan program agitasi adalah 6 kitaran/hari. Faktor yang paling penting yang menyumbang kepada pengeluaran pigmen merah ialah kadar kelembapan awal, dengan nilai F dan nilai p pada 46.54 dan 0.0002, masing-masing diikuti dengan saiz inokulum, dengan nilai F dan nilai p ialah 14.12 dan 0.0071. Pemerhatian yang sama juga terdapat pada biomass, di mana kelembapan awal menjadi penyumbang tertinggi dengan nilai F dan p adalah 55.07 dan 0.0051, diikuti dengan kepekatan peptone dengan nilai F dan p adalah 20.47 dan 0.0202. Kesimpulannya, kajian ini telah membuktikan potensi penggunaan dram bioreaktor dan substrat OPF dalam pengeluaran pigmen merah yang lebih tinggi (71.86 AU/g) berbanding dengan penapaian secara di dalam tabung balang (2.3 AU/g).

## ABSTRACT

Various concern have been reported regarding the effect of existing synthetic colorant, hence, mass production of natural colorant seems to be a potential alternative in fulfilling the demand of colorant in industries. This research has emphasized on the process capability of stirred drum bioreactor on performing solid state fermentation (SSF) of *Monascus purpureus* FTC 5357 by using oil palm frond (OPF) for production of red pigment. Seven parameters, namely, initial moisture content (% v/w), inoculum size (spores/mL), peptone concentration (% w/w), initial pH, aeration rate (vvm), loading capacity (% v/v) and agitation programme (cycles/day) were evaluated for red pigment production, biomass, and glucose consumption. One Factor at A Time (OFAT) and two level factorial screening designs were used in the study. Three responses, which were red pigment production, biomass, and glucose consumption have been used as responses to identify the significant factors affecting the SSF process. In OFAT studies, the highest pigment production was recorded at aeration rate of 1 vvm (6.09 AU/g) and the lowest production was obtained at 45% of initial moisture content (0.1 AU/g). In two level factorial design experimental studies, the highest pigment production was 71.86 AU/g by incorporating 75% v/w of initial moisture content,  $10^8$  spores/mL of inoculum size, initial pH of 4, aeration rate of 1.21 vvm, loading capacity of 35 % v/v and agitation programme of 6 cycles/day. The most significant factor contributing to the red pigment production was initial moisture content, with F-value and p-value of 46.54 and 0.0002, respectively, followed by inoculum size, with F-value of 14.12 and p-value of 0.0071, respectively. Similar observation also occurred for biomass, shows that highest contribution from initial moisture content with F-value of 55.07 and p-value of 0.0051, followed by peptone with F-value of 20.47 and p-value of 0.0202, respectively. In conclusion, this work demonstrated the potential of using stirred drum bioreactor and OPF substrate in producing higher yield of red pigment production (71.86 AU/g) compared to shake flask fermentation (2.3 AU/g).

## TABLE OF CONTENT

### **DECLARATION**

### **TITLE PAGE**

<b>ACKNOWLEDGEMENT</b>	<b>ii</b>
------------------------	-----------

<b>ABSTRAK</b>	<b>iii</b>
----------------	------------

<b>ABSTRACT</b>	<b>iv</b>
-----------------	-----------

<b>TABLE OF CONTENT</b>	<b>v</b>
-------------------------	----------

<b>LIST OF TABLES</b>	<b>viii</b>
-----------------------	-------------

<b>LIST OF FIGURES</b>	<b>ix</b>
------------------------	-----------

<b>LIST OF ABBREVIATIONS</b>	<b>xi</b>
------------------------------	-----------

<b>CHAPTER 1</b>	<b>INTRODUCTION</b>	<b>1</b>
------------------	---------------------	----------

1.1	Background	1
1.2	Problem Statement	2
1.3	Objectives	4
1.4	Scopes	4

<b>CHAPTER 2</b>	<b>LITERATURE REVIEW</b>	<b>5</b>
------------------	--------------------------	----------

2.1	Colorants	5
2.1.1	Natural Colorants	5
2.1.2	Synthetic colorants	6
2.2	<i>Monascus sp.</i>	7
2.2.1	Monascus's red pigment	8
2.3	Oil palm fronds	9
2.3.1	Nutritional aspects of oil palm frond	10
2.3.2	OPF as the substrate for fermentation	12
2.4	Solid state fermentation	12
2.4.1	Environmental factors in SSF	13
2.4.2	Nutritional factors	15

2.5	SSF fermenter	17
	2.5.1 Static flask fermenter	17
	2.5.2 SSF bioreactor	17
2.6	Experimental design	22
	2.6.1 One factor at A Time (OFAT)	22
	2.6.2 2-level factorial design	22
<b>CHAPTER 3                  MATERIALS AND METHOD</b>		<b>24</b>
3.1	Introduction	24
3.2	Chemicals	26
3.3	Preparation	26
	3.3.1 Microorganism	26
	3.3.2 Preparation of <i>Monascus purpureus</i> culture	27
	3.3.3 Inoculum preparation	27
	3.3.4 Substrate preparation	27
3.4	Fermentation	28
	3.4.1 Stirred drum bioreactor study	28
	3.4.2 Preliminary experiment	29
	3.4.3 One Factor at A Time (OFAT)	30
3.5	Analysis	31
	3.5.1 Pigment extraction and analysis	31
	3.5.2 Glucose consumption	31
	3.5.3 Biomass	33
3.6	Experimental design	34
	3.6.1 Two-level factorial design	34
<b>CHAPTER 4                  RESULTS &amp; DISCUSSION</b>		<b>36</b>
4.1	Preliminary result	36
	4.1.1 Viability of red pigment production	36
	4.1.2 Temperature of fermentation	37
	4.1.3 Stirred drum bioreactor	39
4.2	Fermentation profile	39
4.3	One Factor at A Time experiment on red pigment production	41

4.3.1	Effect of initial moisture content	41
4.3.2	Effect of inoculum size	42
4.3.3	Effect of aeration rate	43
4.3.4	Effect of agitation programme	45
4.3.5	Effect of loading capacity content	46
4.3.6	Effect of peptone concentration	48
4.3.7	Effect of initial pH	49
4.4	Two level factorial study	50
4.4.1	Pigment production	52
4.4.2	Biomass	57
4.4.3	Glucose concentration	60
<b>CHAPTER 5</b>	<b>CONCLUSION AND RECOMMENDATIONS</b>	<b>69</b>
5.1	Conclusion	68
5.2	Recommendation	69
<b>REFERENCES</b>		<b>71</b>

## LIST OF TABLES

Table 2.1	Pigment production from other microorganism	6
Table 2.2	Natural and synthetic colorant in market	7
Table 2.3	The analysis of main mineral compositions in OPF	11
Table 3.1	Chemicals and manufacturers	26
Table 3.2	OPF substrate with different condition	29
Table 3.3	Experimental design of the independent parameters	35
Table 4.1	Red pigment production on different nutritional supplement	37
Table 4.2	Pigment production on different incubation environment	38
Table 4.3	Red pigment production at 3 different positions of drum	39
Table 4.4	Range of values of independent variable	50
Table 4.5	Experimental design and results	51
Table 4.6	Analysis of Variance (ANOVA) for red ragment production	54
Table 4.7	Analysis of Variance (ANOVA) for biomass	59
Table 4.8	Analysis of Variance (ANOVA) for glucose	63

## LIST OF FIGURES

Figure 2.1	Anatomy of an oil palm tree and oil palm frond	10
Figure 2.2	Types of SSF bioreactor	18
Figure 2.3	Diagram of the basic bioreactor	20
Figure 2.4	Temperature dynamics of the substrate regarding their positions in the bioreactor	22
Figure 3.1	Overview process of this research	25
Figure 3.2	Major component of the stirred drum bioreactor	28
Figure 3.3	Sampling position on stirred drum bioreactor	30
Figure 3.4	Standard curve for glucose determination	32
Figure 3.5	Standard curve for biomass	34
Figure 4.1	Monascus sp. Culture	38
Figure 4.2	Pigment production and glucose reduction profile of <i>Monascus purpureus</i> FTC 5357 in stirred drum bioreactor	40
Figure 4.3	Effect of initial moisture content	42
Figure 4.4	Effect of inoculum size	43
Figure 4.5	Effect of aeration rate	44
Figure 4.6	Effect of agitation programme	46
Figure 4.7	Effect of loading capacity	47
Figure 4.8	Effect of peptone concentration	48
Figure 4.9	Effect of initial pH	49
Figure 4.10	Pareto chart for red pigment production	53
Figure 4.11	Interaction between initial moisture content and aeration rate on red pigment production	56
Figure 4.12	Interaction between initial moisture content and inoculum size on red pigment production	57
Figure 4.13	Pareto Chart of biomass production	58
Figure 4.14	Interaction between initial moisture content and inoculum size for biomass	60
Figure 4.15	Pareto chart of glucose concentration	61
Figure 4.16	Interaction between initial moisture content and peptone concentration for glucose	64
Figure 4.17	Interaction between initial moisture content and aeration rate for glucose	65
Figure 4.18	Interaction between initial moisture content and inoculum size for glucose	66

- Figure 4.19 Interaction of initial moisture content and inoculum size on lower pH 67
- Figure 4.20 Interaction of initial moisture content and inoculum size on higher pH 68

## LIST OF ABBREVIATIONS

Abs	Absorbance
ANOVA	Analysis of variance
CCD	Central composite design
cm	Centimetre
df	Dilution factor
DM	Dry matter
DNS	Dinitrosalicylic acid
EFB	Empty fruit bunch
g	Gram
hrs	Hour
L	Litre
Mm	Milimetre
MPOB	Malaysian Palm Oil Board
nm	Nanometre
OD	Optical density
OFAT	One factor at One Time
OPB	Oil palm biomass
OPF	Oil palm frond
OPK	Oil palm kernel
OPS	Oil palm shell
OPT	Oil palm trunk
SmF	Submerge fermentation
SSF	Solid state fermentation

## REFERENCES

- Abu Hassan, O., Ishida, M., Mohammed Shukri, I. & Tajuddin, Z. A. (1996). Oil-palm fronds as a roughage feed source for ruminants in Malaysia. Extension Bulletin - ASPAC, Food & Fertilizer Technology Center, (420): 8.
- Aim-oeb, J., Harnbunchong, A., Insung, O.A. & Engkagul, A. (2008). Chemical composition and ruminal dry matter and organic matter degradability of oil palm fronds. *Proceedings of the 46th Kasetsart University Annual Conference,: Animals & Veterinary Medicine*, pp. 46-55.
- Akbari, S., Mahmood, S.M., Tan, I.M., Bharadwaj, A.M. & Hematpour, H. (2017). Experimental investigation of the effect of different process variables on the viscosity of sulfonated polyacrylamide copolymers. *Journal of Petroleum Exploration and Production Technology*, Vol 7, issue 1, pp 87-101.
- Alam, M.Z., Mamun, A.A., Qudsieh, I., Muyibi, S., Salleh, H.M. & Omar, M.N. (2009). Solid state bioconversion of oil palm empty fruit bunches for cellulase enzyme production using a rotary drum bioreactor. *Biochemical Engineering Journal*. 46. 61-64. 10.1016/j.bej.2009.03.010.
- Ali, H.K.Q. & Zulkali, M.M.D. (2011). Design aspects of bioreactors for solid state fermentation: A review. *Chemical and Biochemical Engineering Quarterly*, 25(2): 255-266.
- Asagbra, A.E., Sanni, A.I. & Oyewole, O.B. (2005). Solid-state fermentation production of tetracycline by *Streptomyces* strains using some agricultural wastes as substrate. *World Journal of Microbiology and Biotechnology*, 21(2), 107-114.
- Astimar, A.A., Mohamad, H. & Anis, M. (2002). Preparation of cellulose from oil palm empty fruit bunches via ethanol digestion: effect of acid and alkali catalysts. *Journal of Oil Palm Research*, 14(1): 9-14.
- Babitha, S., Soccol, C.R. & Pandey, A. (2006). Jackfruit seed- a novel substrate for the production of *Monascus* pigments through solid-state fermentation. *Journal of Food Technology & Biotechnology*, 44(4): 465-471.
- Babitha, S., Soccol, C.R. & Pandey, A. (2007). Solid-state fermentation for the production of *Monascus* pigments from jackfruit seed. *Bioresource Technology*, 98(8), 1554-1560.

- Bachalla, N. (2016). Identification of synthetic food colors adulteration by paper chromatography and spectrophotometric methods. *International Archives of Integrated Medicine*, 3(6): 182-191.
- Banerjee, R., Mukherjee, G. & Patra, K.C. (2005). Microbial transformation of tannin-rich substrate to gallic acid through co-culture method. *Bioresource Technology*, 96 (8): 949–953.
- Bau, Y.S. & Wong, H.C. (1979). Zinc effects on growth, pigmentation and antibacterial activity of *Monascus purpureus*. *Physiologia Plantarum*, 46(1), 63-67.
- Bolwerk, A., Lagopodi, A.L., Wijfjes, A.H.M., Lamers, G.E.M., Thomas, F.C., Woeng, C.A., Lugtenberg, B.J.J. & Bloemberg, G.V. (2003). Interactions in the tomato rhizosphere of two pseudomonas biocontrol strains with the phytopathogenic fungus *Fusarium oxysporum*. *Radicis-Lycopersici*. MPMI, 16(11), 983–993.
- Broder, C.U. & Koehler, P.E. (1980). Pigments produced by *Monascus purpureus* with regard to quality and quantity. *Journal of Food Science*, 45(3), 567-569.
- Calvo, A.M., Wilson, R.A., Bok, J.W. & Keller, N.P. (2002). Relationship between secondary metabolism and fungal development. *Microbiology Molecular Biology*, R66: 447–459.
- Carels, M. & Shepherd, D. (1979). The effect of changes in pH on phosphate and potassium uptake by *Monascus rubiginosus* ATCC 16367 in submerged shaken culture. *Canada Journal Microbiology*, 25(12): 1484-1488.
- Carels, M. & Shepherd, D. (1977). The effect of different nitrogen sources on pigment production and sporulation of *Monascus* species in submerged, shaken culture. *Canadian Journal of Microbiology*, 23(10): 1360-1372.
- Carocho, M., Barreiro, M.F., Morales, P. & Ferreira, I.C.F.R. (2014). Adding molecules to food, pros and cons: A review on synthetic and natural food additives. *Comprehensive Reviews in Food Science and Food Safety*. Vol. 13. doi: 10.1111/1541-4337.12065
- Carvalho, J.C., Oishi, B.O., Woiciechowski, A.L., Pandey, A., Babitha, S. & Soccol, C. R. (2007). Effect of substrates on the production of *Monascus* biopigments by solid-state fermentation and pigment extraction using different solvents. *Indian Journal of Biotechnology*, 6(2): 194.
- Carvalho, J.C.D., Pandey, A., Oishi, B.O., Brand, D., Leon, J.A.R. & Soccol, C.R. (2006). Relation between growth, respirometric analysis and biopigments production from

*Monascus* by solid-state fermentation. *Biochemical Engineering Journal*, 29(3): 262-269.

Chen, M.H. & Johns, M.R. (1993). Effect of pH and nitrogen source on pigment production by *Monascus purpureus*. *Applied Microbiology and Biotechnology*, 40(1):132-138.

Chen, M.H. & Johns, M.R. (1994). Effect of carbon source on ethanol and pigment production by *Monascus purpureus*. *Enzyme and Microbial Technology*, 16(7), 584-590.

Chysirichote, T. (2013). Kinetics of growth and red pigment production of the fungus *Monascus ruber* in solid surface cultures. Ph.D. Thesis. The graduate school of science and engineering, Tokyo Institute of Technology, Tokyo, Japan.

Chysirichote, T. (2016). Effect of aeration mode on the red pigment production by *Monascus ruber* on rice during fermenting in the packed bed bioreactor. *KKU Resources Journal*, 21 (2), 176-184.

CIRAD, 1991. *Laboratory data 1963-1991*. France: Centre de coopération Internationale en Recherche Agronomique pour le Développement.

Collins, L. M., Dziak, J. J., Kugler, K. C., & Trail, J. B. (2014). Factorial experiments: efficient tools for evaluation of intervention components. *American journal of preventive medicine*, 47(4), 498-504.

Cortez, R., Luna-Vital, D.A., Margulis, D. & Mejia, E.G. (2017). Natural pigments: Stabilization methods of anthocyanins for food applications. *Comprehensive Reviews in Food Science and Food Safety*, Vol.16, doi: 10.1111/1541-4337.12244

Costa, F.T.M., Justo, G.Z., Dura'n, N., Nogueira, P.A. & Lopes, S.C.P.: The use of violacein in its free and encapsulated form in polymeric systems against malaria. *Brazilian Patent PIBr*, 056399-0

Czitrom, V. (1999). One Factor at A Time versus Designed experiment. *The American Statistician*, Vol 53, No 2.

Dahlan, I. (2000). Oil palm frond, a feed for herbivores. *Asian-Australia Journal Animal Science*, 300-303.

Danuri, H. (2008). Optimizing angkak pigments and lovastatin production by *Monascus purpureus*. *Journal of Biosciences*, Vol 15, No 2.

Davami, F., Baldi, L., Rajendra, Y., & M. Wurm, F. (2014). Peptone supplementation of culture medium has variable effects on the productivity of CHO Cells. *International Journal of Molecular and Cellular Medicine*, 3(3), 146–156.

Di Mascio, P., Kaiser, S. & Sies, H. (1989). Lycopene as the most efficient biological carotenoid singlet oxygen quencher. *Arch Biochemistry Biophysic*, 274(2):532–538. doi: 10.1016/0003-9861(89)90467-0.

Dufossé, L. (2016). Current and potential natural pigments from microorganisms (bacteria, yeasts, fungi, microalgae). In: Carle, R., Schweiggert, R. (Eds.), *Handbook on Natural Pigments in Food and Beverages. Industrial Applications for Improving Food Colour*, first ed. Sawston, Cambridge: Woodhead Publishing (Elsevier group), pp. 337–354.

Dufossé, L. (2009). Microbial and microalgal carotenoids as colourants and supplements. *In Carotenoids Birkhäuser Basel*, 83–98.

Durand, A. (2003). Bioreactor designs for solid state fermentation. *Biochemical Engineering Journal*, 13(3):113-125.

Fannin, T. E., Marcus, M. D., Anderson, D. A., & Bergman, H. L. (1981). Use of a fractional factorial design to evaluate interactions of environmental factors affecting biodegradation rates. *Applied and environmental microbiology*, 42(6), 936-43.

Florencio, J.A., Soccol, C.R., Furlanetto, L.F., Bonfim, T.M.B., Krieger, N., Baron, M. & Fontana, J.D. (1998). A factorial approach for a sugarcane juice-based low cost culture medium: increasing the astaxanthin production by the red yeast *Phaffia rhodozyma*. *Bioprocess Engineering*, 19:161–164.

Flores-Cotera, L.B. & Sanchez, S. (2001). Copper but not iron limitation increases astaxanthin production by *Phaffia rhodozyma* in a chemically defined medium. *Biotechnology Letter*, 23:793–797. doi: 10.1023/A:1010358517806.

Future Market Insight (FMI). (2014). Natural food colours market: Global industry-analysis and opportunity assessment 2014-2020, available from <http://www.futuremarketinsights.com/reports/details/global-natural-food-colours-market>, accessed 9 November 2017.

Gabiatti, J.R., Vendruscolo, C.F., Piaia, J.C.Z., Rodrigues, R.C., Durrant, L.R. & Costa, J.A.V. (2006). Radial growth rate as a tool for the selection of filamentous fungi

for use in bioremediation. *Brazilian Archives of Biology and Technology*, 49: 29-34.

Ghani, A.A.A., Rusli, N.D., Shahudin, M.S., Goh, Y., Saad, M., Hafandi, A. & Hassim, H.A. (2017). Utilisation of oil palm fronds as ruminant feed and its effect on fatty acid metabolism. *Pertanika Journal of Tropical Agricultural Science*. 40. 215-224.

Giovannucci, E., Rimm, E.B., Liu, Y., Stampfer, M.J. & Willett, W.C. (2002). A prospective study of tomato products, lycopene, and prostate cancer risk. *Journal National Cancer Institute*. 94(5):391–398. doi: 10.1093/jnci/94.5.391

Gmoser, R., Ferreira, J.A., Lundin, M., Taherzadeh, M.J. & Lennartsson, P.R. (2018). Pigment production by the edible filamentous fungus *Neurospora Intermedia*. *Fermentation*, 4,11; doi:10.390

Gonzalez, J.B., Fernandez, F.J. & Tomasini, A. (2003). Microbial secondary metabolite production and strain improvement. *Indian Journal Biotechnology*. 2(3): 322-333.

Grimm, L., Kelly, S., Krull, R. & Hempel. D. (2005). Morphology and productivity of filamentous fungi. *Applied Microbiology and Biotechnology*, 69(4): 375-384.

Gross, S. & Robbins, E. (2000). Acidophilic and acid-tolerant fungi and yeasts. *Hydrobiologia*. 433. 91-109. 10.1023/A:1004014603333

Gulrajani, M.L. (2001). Present status of natural dyes. *Indian Journal of Fibre and Textile Research*, Vol 26, pp- 191-201.

Gutierrez-Rojas, M., Hosn, S., Auria, R., Revah, S. & Favela-Torres, E. (1996). Heat transfer in citric acid production by solid state fermentation. *Process Biochemistry*, 31(4): 363-369.

Hajjaj, H., Klaebe, A., Goma, G., Blanc, P.J. & Barbier, E. (2000). Mediumchain fatty acids affect citrinin production in the filamentous fungus *Monascus ruber*. *Applied Environmental Microbiology*, 66(3):1120-1125.

Hamid, N.F. & Said, F.M. (2016).Factorial design screening for the red pigment production by *Monascus purpureus* FTC 5356. *Jurnal Teknologi*, Vol 78.

Hamid, N.F. (2017). Production of red pigment from *Monascus purpureus* FTC 5356 using oil palm frond in solid state fermentation. Master Thesis. Universiti Malaysia Pahang. Malaysia.

Hamid, N.F. & Said, F.M. (2018). Optimization of red pigment production by *Monascus purpureus* FTC 5356 using response surface methodology, *IIUM Engineering Journal*, 19 (1), pp 34-47.

Han, O. & Mudgett, R.E. (1992). Effects of oxygen and carbon dioxide partial pressures on *Monascus* growth and pigment production in solid state fermentations. *Biotechnology Progress*, 8, 5-10.

Hassan, O.A., Ishida, M., Shukri Mohd, I. & Tajuddin, Z.A. (1994). Oil-palm fronds as a roughage feed source for ruminants in Malaysia. ASPAC, Food & Fertilizer Technology Center.

Heber, D., Yip, I., Ashley, J.M., Elashoff, D.A., Elashoff, R.M. & Go, V.L. (1999). Cholesterol-lowering effects of a proprietary Chinese red-yeast-rice dietary supplement. *American Journal of Clinical Nutrition*, 69(2): 231-236.

Hong, L. S., Ibrahim, D. & Omar, I. (2012). Microscopic studies of oil palm frond during processing for saccharification. *The Internet Journal of Bioengineering*, 4(2).

Hossain, M.A., Laxmi, J., Ganesan, P., Sahu, J., Ramesh, S. & Poh, S.C. (2016). Microwave pyrolysis of oil palm fiber (OPF) for hydrogen production: Parametric investigation. *Energy Conversion and Management*. 115. 232-243. 10.1016/j.enconman.2016.02.058.

Ibrahim, Z., Aziz, A.A., RamLi, R., Mokhtar, A., Omar, R. & Lee, S. (2014). Production of medium density fibreboard (MDF) from oil palm trunk (OPT). *Journal of Applied Sciences*. 14. 1174-1179. 10.3923/jas.2014.1174.1179

Inayat, A., Ahmad, M.M., Mutalib, M. I. A., Yusup, S. & Khan, Z. (2017). Economic analysis and optimization for bio-hydrogen production from oil palm waste via steam gasification, *Energy Sources, Part B: Economics, Planning, and Policy*, 12:2, 158-165, DOI: 10.1080/15567249.2014.937881

Institute of Food Technologists. (2016). Coloring Foods & Beverages. Available from Available from: <http://s36.a2zinc.net/clients/IFT/IFT16/Public>, Accessed 2016 July 20.

Ishida, M. & Abu Hassan, O. (1997). Utilization of oil palm frond as cattle feed. *JARQ, Japan Agricultural Research Quarterly*, 31(1): 41-47.

Islam, M., Dahlan, I., Rajion, M.A. & Jelan, Z.A. (2000). Productivity and nutritive values of different fractions of oil palm (*Elaeis guineensis*) frond. *Asian-Australasian Journal of Animal Sciences*, 13(8): 1113-1120.

Jahromi, M.F., Liang, J.B., Ho, Y.W., Mohamad, R., Goh, Y.M. & Shokryazdan, P. (2012). Lovastatin production by *Aspergillus terreus* using agro-biomass as substrate in solid state fermentation. *Journal of Biomedicine and Biotechnology*, 2012.

Jirasatid, S. & Nopharatana, M. (2017). Improvement of pigments production of rice bran fermented by *Monascus purpureus*. The 6<sup>th</sup> Burapha University International Conference 2017.

Johns, M.R. & Stuart, D.M. (1991). Production of pigments by *Monascus purpureus* in solid culture. *Journal of Industrial Microbiology*, 8: 23-28.

Johnson, G.T. & Mchan, F. (1975). Some effects of zinc on the utilization of carbon sources by *Monascus purpureus*. *Mycologia*, 67(4): 806-16.

Kaimanen, M. (2014). Stability of natural colorants of plant origin. PhD Thesis. University of Turku. Finland.

Khalil, H. P. S., Jawaid, M., Hassan, A., Paridah, M. T. & Zaidon, A. (2012). Oil palm biomass fibres and recent advancement in oil palm biomass fibres based hybrid biocomposites. Intech Open, pp. 187-220.

Khamseekhiew, B., Liang, J.B., Wong, C.C. & Jalan, Z.A. (2001). Ruminal and intestinal digestibility of some tropical legume forages. *Asian Australasian Journal of Animal Sciences*, 14(3): 321-325.

Kim, C.H., Kim, S.W. & Hong, S.I. (1999). An integrated fermentation-separation process for the production of red pigment by *Serratia* sp. KH-95. *Process biochemistry*, 35(5): 485-490.

Kosugi, A., Tanaka, R., Magara, K., Murata, Y., Arai, T., Sulaiman, O. & Mori, Y. (2010). Ethanol and lactic acid production using sap squeezed from old oil palm trunks felled for replanting. *Journal of Bioscience and Bioengineering*, 110(3): 322-325.

Krasniewski, I., Molimard, P., Feron, G., Vergoignan, C., Durand, A. & Cavin, J. F. (2006). Impact of solid medium composition on the conidiation in *Penicillium camemberti*. *Process Biochemistry*, 41, 1318–1324.

Kristiani, A., Abimanyu, H., Setiawan, A.H., Sudiyarmanto, & Aulia, F. (2013). Effect of pretreatment process by using diluted acid to characteristic of oil palm frond. *International conference on Sustainable Energy Engineering and Application*. 1876-6102.

Kusdiyantini, E., Nugraha, S. & Lunggani, A. (2017). Pigment production of *Monascus* sp. Isolated from angkak in Semarang region, Central Java, Indonesia. *Indonesian Food and Nutrition Progress*, Vol 14 (1), doi: 10.22146/ifnp.24256

Lee, B.K., Park, N.H., Piao, H.Y. & Chung, W.J. (2001). Production of red pigments by *Monascus purpureus* in submerged culture. *Biotechnology and Bioprocess Engineering*, 6(5): 341-346.

Lee, B.K., Piao, H.Y. & Chung, W.J. (2002). Production of red pigments by *Monascus purpureus* in solid state culture. *Biotechnology Bioprocess Engineering*, 7:21-25.

Li, X., Xu, J., Tang, X., Liu, Y., Yu, X., Wang, Z. & Liu, W. (2016a). Anthocyanins inhibit trastuzumab resistant breast cancer in vitro and in vivo. *Molecular Medical Report*, 13:4007–13.

Lin, T.F. & Demain, A.L. (1991). Effect of nutrition of *Monascus* sp. on formation of red pigments. *Applied Microbial Biotechnology*, 36, 70-75.

Lonsane, B.K. & Ramesh, M.V. (1992). Production of bacterial thermostable enzymes by solid state fermentation: a potential tool for achieving economy in enzyme production. *Advances in Applied Microbiology*, 15: 1-48.

Lonsane, B.K., Ghildyal, N.P., Budiatman, S. & Ramakrishna, S.V. (1985). Engineering aspects of solid state fermentation. *Enzyme and Microbial Technology*, 7(6): 258-265.

Ma, J., Li, Y., Ye, Q., Li, J., Hua, Y., Ju, D., Zhang, D., Cooper, R. & Chang, M. (2000). Constituents of red yeast rice, a traditional Chinese food and medicine. *Journal Agricultural Food Chemistry*. 48: 5220-5225.

Manan, M.A. & Webb, C. (2017). Design aspects of solid state fermentation as applied to microbial bioprocessing. *Journal Applied Biotechnology Bioengineering*, 4(1): 00094. doi: 10.15406/jabb.2017.04.00094

Manan, M.A. (2014). Design aspects of solid state fermentation. PhD Thesis. The University of Manchester, United Kingdom.

Manan, M.A., Mohamad, R. & Ariff, A. (2017). *Monascus* sp.: A source of natural microbial colour through fungal biofermentation. *Journal of Microbiology and Experimentation*. 5(3): 00148.

Mandal, B.K., Mathiyalagan, S., Dalavai, R., & Ling, Y.C. (2017). Simultaneous identification of synthetic and natural dyes in different food samples by UPLC-MS. *IOP Conference Series: Materials Science and Engineering*, 263 (2017) 022011 doi:10.1088/1757-899X/263/2/022011

Mapari, S.A., Thrane, U. & Meyer, A.S. (2010). Fungal polyketide azaphilone pigments as future natural food colorants? *Trends in biotechnology*, 28(6): 300-307.

Martins, S., Mussatto, S.I., Martínez-Avila, G., Montanez-Saenz, J., Aguilar, C.N. & Teixeira, J.A. (2011). Bioactive phenolic compounds: production and extraction by solid-state fermentation. A review. *Biotechnology Advance* 29, 365–373.

Mazaheri, D., Shojaosadati, S.A., Mousavi, S.M., Hejazi, P. & Saharkhiz, S. (2012). Bioethanol production from carob pods by solid-state fermentation with *Zymomonas mobilis*. *Applied Energy*.

Mchan, F. & Johnson, G.T. (1970). Zinc and amino acids: Important components of a medium promoting growth of *Monascus purpureus*. *Mycologia* 62: 1018-1031.

Meyer. H.G. (1990).The effect of nitrogen and phosphate in the formation of pigment in *Monascus purpureus* Went DSM 1379, Thesis, Department of Microbiology, University of Saarland, Saarbrücken. England. 446p.

Miller, G.L. (1959). Use of dinitrosalicylic acid reagent for determination of reducing sugar. *Analytical Chemistry*, 31, 426-428.

Mitchell, D. A., Berovic, M., & Krieger, N. (2006). Solid state fermentation bioreactors: fundamentals of design and operations. *Springer*. Verlag Berlin Heidelberg.

Mitchell, D.A., Lonsane, B.K., Durand, A., Renaud, R., Almanza, S., Maratary, J., Desgranges, C., Crooke, P.S., Hong, K., Tanner, R.D. & Malaney, G.W. (1992). General principles of reactor design and operation for SSC p115-136In: Solid-substrate Cultivation, Doelle HW (ed.), Elsevier, Barking, Essex.

Mortensen, A. (2006). Carotenoids and other pigments as natural colorants. *Pure Applied Chemistry*, Vol 78, No 8, pp. 1477-1491. doi:10.1351/pac200678081477

Mostafa, M.E. & Abbady, M.S. (2014). Secondary metabolites and bioactivity of the *Monascus* pigments review article. *Global Journal of Biotechnology & Biochemistry*, 9 (1): 01-13, doi: 10.5829/idosi.gjbb.2014.9.1.8268

Mpountouka, P., Panatzaki, A., Kostareli, E., Christo-doulou, P., Kareli, D., Poliliou, S., Mourelatos, C., Lam-bropoulou, V. & Laliaris, T. A. (2010). Cytogenetic

evaluation and DNA interaction studies of the food colorants amaranth, erythrosine and tartrazine. *Food and Chemical Toxicology*, 4810, 2934-2944.

Mudgett, R.E. (1986). Solid state fermentation. In: Demain, A.L. & Solomon, N.A. (Eds.) *Manusla of Industrial Microbiology and Biotechnology*, American Society for Microbiology, Washington, USA, pp 66-83.

Mukherjee, G. & Singh, S.K. (2011). Purification and characterization of a new red pigment from *Monascus purpureus* in submerged fermentation. *Process Biochemistry*. 46: 188-192.

Rao, M. P., Xiao, M., & Li, W. J. (2017). Fungal and bacterial pigments: secondary metabolites with wide applications. *Frontiers in Microbiology*, 8, 1113. <http://doi.org/10.3389/fmicb.2017.01113>

National Cancer Institute (NCI). (2015). Benzidine. Available from <https://www.cancer.gov/about-cancer/causes-prevention/risk/substances/benzidine>, accessed 9<sup>th</sup> April 2018.

Nimnoi, P. & Lumyoung, S. (2009). Improving solid state fermentation of *Monascus purpureus* on agricultural products for pigment production. *Food & Bioprocess Technology*, doi: 10.1007/s11947-009-0233-8.

Onken, U. & Liefke, E. (1989). Effect of total and partial pressure (oxygen and carbon dioxide) on aerobic microbial processes. *Advances in Biochemistry, Engineering and Biotechnology*, 40, 137–169.

Oostra, J., Tamper, J. & Rinzema, A. (2000). Model-base bioreactor selection or large-scale solid state cultivation of *Coniothyrium minitans* spores on oats. *Enzyme and Microbial Technology*, 27, 652-663.

Orozco, S.F.B. & Kilikian, B. (2008). Effect of pH on citrinin and red pigments production by *Monascus purpureus* CCT3802. *World Journal of Microbiology and Biotechnology*. 24. 263-268. 10.1007/s11274-007-9465-9.

Pandey, A. (2003). Solid-state fermentation. *Biochemistry Engineering Journal*, 13(2), 81–84.

Pandey, A., Soccol, C.R., and Mitchell, D. (2000) New developments in solid state fermentation:I-bioprocesses and products. *Process Biochemistry*, 35, 1153–1169.

Pandey, A., Soccol, C.R. & Larroche, C. (2007) Current Developments in Solid-state Fermentation,in: A.Pandey,C.R.Soccol,C.Larroche(Eds.),Current Developments in

Solid-state Fermentation, Springer Science/Asiatech Publishers, Inc., New York, USA/New Delhi, India, pp. 13–25

- Pattanagul, P., Pinthong, R., Phianmongkhol, A. & Leksawasdi, N. (2007). Review of angkak production (*Monascus purpureus*). *Chiang Mai Journal of Science*, 34, 319–328.
- Pengnoi, P., Mahawan, R., Khanongnuch, C. & Lumyong, S. (2017). Antioxidant properties and production of Monacolin K, citrinin, and red pigments during solid state fermentation of purple rice (*Oryzae sativa*) varieties by *Monascus purpureus*. *Czech Journal of Food Sciences*, 35. 10.17221/154/2016-CJFS.
- Rafidah, D., Ainun, Z. M.A., Hazwani, H.A., Rushdan, I., Luqman, C.A., Sharmiza, A., Paridah, M.T. & Jalaluddin, H. (2017). Characterisation of pulp and paper manufactured from oil palm empty fruit bunches and kenaf fibres. *Pertanika Journal Tropical Agricultural Science*, 40 (3): 449 – 458.
- Raimbault, M. & Alazard, D. (1980). Culture method to study fungal growth in solid fermentation. *European Journal Application Microbiology Biotechnology*, 9, 199–209.
- Ramirez, I., Nunez, M.L. & Valdivia, R. (2000). Increased astaxanthin production by a *Phaffia rhodozyma* mutant grown on date juice from *Yucca fillifera*. *Journal Industrial Microbiology Biotechnology*, 24:187–190. doi: 10.1038/sj.jim.2900792.
- Rasli, S.R.A.M., Ishak, A., Lazim, A.M. & Hamzah, A. (2017). Extraction and characterization of cellulose from agricultural residue-oil palm fronds. *Malaysian Journal of Analytical Sciences*, Vol 21, No 5: 1065 – 1073.
- Razali, M.AA & Said, F.M. (2017). Red pigment production by *Monascus purpureus* in stirred-drum bioreactor. *Galeri Warisan Sains*, 1 (1) (2017), pp. 13-15.
- Rondle, C.J.M. & Morgan, W.T.J. (1955). The determination of glucosamine and galactosamine. *Biochemistry Journal*, 61, 586.
- Roopesh, K., Ramachandran, S., Nampoothiri, K.M., Szakacs, G. & Pandey, A. (2006). Comparison of phytase production on wheat bran and oilcakes in solid state fermentation by *Mucor racemosus*. *Bioresource Technology*, 97, 506-511.
- Said, F.M. (2010) *Monascus ruber* ICMP 15220 fermentation for the production of pigments. Ph. D Thesis, Massey University, New Zealand.

- Saithi, S. & Tongta, A. (2016). Phytase production of *Aspergillus niger* on soybean meal by solid state fermentation using rotating drum bioreactor. *Agricultural Science Procedia*, 11, 25-30.
- Sakurai, Y., Lee, T.H. & Shiota, H. (1977). On the convenient method for glucosamine estimation in koji. *Agricultural and Biological Chemistry*, 41, 619-624.
- Santos-Ebinuma, V. C., Roberto, I. C., Teixeira, M. F. S. & Pessoa, A. (2014). Improvement of submerged culture conditions to produce colorants by *Penicillium purpurogenum*. *Brazilian Journal of Microbiology*, 45(2), 731–742.
- Seal, H.P., Ali, M.A., Ali, M.U., Akhter, M.H. & Sultana, F. (2012). Thin layer chromatographic analysis of food colorants from three morphotypes of annato (*Bixa orellana L.*). *International Journal Agriculture Resources Innovation and Technology*, 2(1): 2224-0616.
- Selamat, M.D. (2010). Extraction of green pigment from *Pandanus odoratus*. Degree Thesis. Universiti Malaysia Pahang. Malaysia.
- Shuit, S.H., Tan, K.T., Lee, K.T. & Kamaruddin, A.H. (2009). Oil palm biomass as a sustainable energy source: A Malaysian case study. *Energy*, Vol. 34, No. 9, pp. 1225-1235.
- Sigurdson, G., Tang, P. & Giusti, M.M. (2017). Natural colorants: Food colorants from natural sources. *Annual Review Food Science Technology*, 8: 261- 280.
- Singhania, R.R., Patel, A.K., Soccol, C.R. & Pandey, A. (2009). Recent advances in solid-state fermentation. *Biochemical Engineering Journal*, 44, 13–18.
- Skromne, I., Sanchez, O. & Aguirre, J. (1995). Starvation stress modulates the expression of the *Aspergillus nidulans* brlA regulatory gene. *Microbiology*, 141: 21–28.
- Soccol, C.R. & Vandenberghe, L.P.S. (2003). Overview of applied solid state fermentation in Brazil. *Biochemical Engineering Journal*, 13, 205-218.
- Soccol, C.R., Costa, E., Scopel, Letti, L.A.J., Karp S.G., Woiciechowski A.L. & Vandenberghe L.P.S. (2017). Recent developments and innovations in solid state fermentation. *Biotechnology Research and Innovation*, 1, 52-71. doi.org/10.1016/j.biori.2017.01.002
- Srianta, I., Hendrawan, B., Kusumawati, N. & Blanc, P.J. (2012). Study on durian seed as a new substrate for angkak production. *International Food Research Journal*, 19 (3): 941-945.

Su, Y.C. (1978). The production of *Monascus* pigments (in Chinese). *Food Science*. 5:4A-17A.

Sudha, T., Divya, G., Sujaritha, J. & Duraimurugan, P. (2017). Review of experimental design in analytical chemistry. *Indo American Journal of Pharmaceutical Research*, 7 (08).

Swift, M.J. (1973). The estimation of mycelial biomass by determination of the hexosamine content of wood tissue decayed by fungi. *Soil Biology Biochemistry*, 55,321-332.

Temudo, M.F., Mato, T., Kleerebezem, R. & Mark, C.M. (2009). Xylose anaerobic conversion by open-mixed cultures. *Applied Microbiology and Biotechnology*. 82: 231.

Tengerdy, R.P. (1985). Solid substrate fermentation. *Trends Biotechnology*, 3:96–99.

Tseng, K.C, Fang, T.J., Chiang, S.S., Liu C.F., Wu C.L. & Pan, T.M. (2012). *Journal Science Foz2od Agriculture*, 92(7):1483-9.

Umsza-Guez, M.A., Diaz, A.B., de Ory, I., Blansino, A., Gomes, E. & Caro, I. (2011). Xylanase production by *Aspergillus awamori* under solid state fermentation conditions on tomato pomace. *Brazillian Journal of Microbiology* ,42(4): 1585-1597.

Van Den Oever, M. J. A., Boeriu, C. G., Blaauw, R., & Van Haveren, J. (2004). Colorants based on renewable resources and food grade colorants for application in thermoplastics. *Journal of Applied Polymer Science*, 92, 2961–2969.

Velmurugan, P., Hur, H., Balachandar, V., Kannan, S.K., Lee, K.J., Lee, S.M., Chae, J.C., Shea, P.J. & Oh, B.T. (2011). *Monascus* pigment production by solid-state ferment6+ation with corn cob substrate. *Journal of Bioscience and Bioengineering*, 112:6, pp.590-594; doi: 10.1016

Vidyalakshmi, R., Paranthaman, R., Murugesh, S. & Singaravel, K. (2009). Stimulation of *Monascus* pigments by intervention of different nitrogen sources. *Global Journal of Biotechnology and Biochemistry*, 4: 25-28.

von Meien, O.F., Luz, L.F.L., Mitchell, D.A., Pérez-Correa, J.R., Agosin, E., Fernández, M.F. & Arcas, J. (2004). Control strategies for intermittently mixed, forcefully aerated solid-state fermentation bioreactors based on the analysis of a distributed parameter model. *Chemical Engineering Science*. 59. 4493-4504. 10.1016/j.ces.2004.06.027.

- Walker, G.M. & White, N.A. (2005). Introduction to fungal physiology. McGraw-Hill Publishers London: 10. Volume 99, pp 372-378, doi.org/10.1016/j.apenergy.2012.05.045.
- Wan Zahari, M., Sato, J., Furuichi, S., Azizan, A.R. & Yunus, M. (2004). Commercial processing of oil palm fronds feed in Malaysia. Proceeding of 8th Meeting of Regional Working Group on Grazing and Feed Resources for South-East Asia, MARDI.
- Wang, J., Shi, H. & Yi, Q. (2000). Wastewater treatment in a hybrid biological reactor (HBR): Effect of organic loading rates. *hy.* 36. 297-303. 10.1016/S0032-9592(00)00153-9.
- Weiland, P. (1988). Principles of solid state fermentation, In : Treatment of lignocellulosics with white rot fungi. Zadrazil, F. and Reniger, P. (Eds.). Elsevier, London. pp 64-67.
- Weiss, B. (2012). Synthetic food colors and neurobehavioral hazards: The view from environmental health research. *Environmental Health Perspective.* 120(1): 1–5.
- Wong; H. C., Lin Y. C. & Koehler, P. E. (1981). Regulation of growth and pigmentation of *Monascus purpureus* by carbon and nitrogenconcentration. *Mycologia,* 73(4):649-654.
- Wongjewboot, I. & Kongruang, S. (2011). pH stability of ultrasonic Thai isolated *Monascus purpureus* pigments. *International Journal of Bioscience, Biochemistry and Bioinformatics,* Vol. 1, No, 1.
- Wongjewboot, I., Bowdang, T. & Kongruang, S. (2012). Stability of colorants from ultrasonic induced *Monascus purpureus*. *3rd International Conference on Biology, Environment and Chemistry,* Vol 46. doi: 10.7763/IPCBEE. 2012. V46. 26
- Xie, G., (2006) Ph.D. Thesis. University of South Dakota, USA.
- Yongsmit, B., Kitprechavanich, V., Chitradon, L. & Budda, N. (2000). Color mutants of *Monascus* sp. KB9 and their comparative glucoamylases on rice solid culture. *Journal of Molecular Catalysis B Enzymatic,* 10(1-3): 263-272.
- Yongsmit, B., Tabloka, W., Yongmanitchai, W., and Bavavoda, R. (1993). Culture conditions for yellow pigment formation by *Monascus* sp. KB 10 grown on cassava medium, *World Journal Microbiology Biotechnology.* 9, 85–90.

Yoshimura, M., Yamanada, S., Mitsugi, K. & Hirose, Y. (1975). Production of *Monascus* pigment in a submerged culture. *Agrotechnology Biology Chemistry*, 39:1789-1795.

Zahari M. A. K. M., Zakaria, M. R., Ariffin, H., Mokhtar, M. N., Sa-lihon, J., Shirai, Y. & Hassan, M. A. (2012). *Bioresource Technology*. 110: 566-571.

Zhou, B., Pu, Y.W., Zhu, M.J. & Liang, S.Z. (2008). Effects of nitrogen sources on *Monascus* yellow pigment production by *Monascus* mutant. *Modern Food Science and Technology* 2:123-127.