

A review on taste masked multiparticulate dosage forms for paediatric for tenacity and toughness improvement

AL-Japairai, Khater^a; Hamed Almurisi, Samah^b; Almonem Doolaanea, Abd^c; Mahmood, Syed^d; Alheibshy, Fawaz^{e, f}; Alobaida, Ahmed^e; Abdul-Halim, Nadiya^a; Chatterjee, Bappaditya

^a Department of Pharmaceutical Engineering, Faculty of Chemical and Process Engineering Technology, Universiti Malaysia Pahang, Gambang, 26300, Malaysia

^b Department of Pharmaceutical Technology, Kulliyah of Pharmacy, International Islamic University Malaysia, Kuantan, 25200, Malaysia

^c Department of Pharmaceutical Technology, Faculty of Pharmacy, Kolej Universiti Antarabangsa Maiwp (UCMI), Taman Batu Muda, Batu Caves, Kuala Lumpur, 68100, Malaysia

^d Department of Pharmaceutical Technology, Faculty of Pharmacy, Universiti Malaya, Kuala Lumpur, 50603, Malaysia

^e Department of Pharmaceutics, College of Pharmacy, University of Hail, Hail, 2240, Saudi Arabia

^f Department of Pharmaceutics, College of Pharmacy, Aden University, Aden, 6075, Yemen

^g SPP School of Pharmacy and Technology Management, SVKM's NMIMS, Mumbai, India

ABSTRACT

Taste refers to those sensations perceived through taste buds on the tongue and oral cavity. The unpleasant taste of drugs leads to the refusal of taking the medicine in the paediatric population. It is widely known that a pharmaceutical product's general acceptability is the result of numerous contributing components such as swallowability, palatability (taste, flavour, texture, and mouthfeel), appearance, ease of administration, and patient characteristics. Multiparticulate as a dosage form is a platform technology for overcoming paediatrics' incapacity to swallow monolithic dosage forms, masking many medications' inherent nasty taste, and overcoming the obstacles of manufacturing a commercially taste masked dosage form. This review will discuss the considerations that must be taken into account to prepare taste masked multiparticulate dosage forms in the best way for paediatric use.

KEYWORDS

Multiparticulate system; Paediatric; Taste masking; Unpleasant drug

ACKNOWLEDGEMENTS

The authors are thankful to the universiti Malaysia Pahang for providing support in the form of an internal grant (RDU220308).