

## Research Article

# Lipid Enhancement in Oleaginous *Nannochloropsis* sp. under Nitrate Limitation for Future Bioenergy Production

Karthick Murugan Palanisamy <sup>1</sup>, Prakash Bhuyar <sup>2</sup>, MohdHasbi Ab. Rahim <sup>1</sup>,  
Ashiwin Vadiveloo <sup>3</sup>, Naif Abdullah Al-Dhabi <sup>4</sup>, Natanamurugaraj Govindan <sup>5</sup>,  
and GaantyPragas Maniam <sup>1</sup>

<sup>1</sup>Algae Culture Collection Center & Laboratory, Faculty of Industrial Sciences and Technology, University of Malaysia Pahang Al-Sultan Abdullah, Lbh Persiaran Tun Khalil Yaakob, Kampung Melayu Gambang, 26300 Kuantan, Pahang, Malaysia

<sup>2</sup>International Industry and Agriculture Innovation Research Center (IIAR), International College, Maejo University, Nong Han, San Sai, Chiang Mai 50290, Thailand

<sup>3</sup>Algae R&D Centre, School of Biological Sciences & Biotechnology, Murdoch University, Murdoch 6150, Western Australia, Australia

<sup>4</sup>Department of Botany and Microbiology, College of Science, King Saud University, P.O. Box 2455, Riyadh 11451, Saudi Arabia

<sup>5</sup>Department of Botany and Microbiology, Hemvati Nandan Bahuguna Garhwal University, Chauras Campus Srinagar Garhwal, 249161 Uttarakhand, India

Correspondence should be addressed to GaantyPragas Maniam; [gaanty@umpsa.edu.my](mailto:gaanty@umpsa.edu.my)

Received 12 July 2023; Revised 7 October 2023; Accepted 25 November 2023; Published 14 December 2023

Academic Editor: Kiran Babu Uppuluri

Copyright © 2023 Karthick Murugan Palanisamy et al. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Microalgae possess high oil content, exhibit rapid growth rates and biomass productivity, and leave a minimal environmental footprint, making them highly attractive for biofuel applications. Consequently, this present study is aimed at evaluating the impact of nitrogen starvation on the marine microalgae *Nannochloropsis* sp. to enhance lipid accumulation. The microalga culture was grown under an irradiance of  $120 \mu\text{mol photon m}^{-2} \text{s}^{-1}$ , temperature  $25 \pm 1^\circ\text{C}$ , continuous air-CO<sub>2</sub>, and with different concentrations of sodium nitrites over a 14-day cultivation period. From the results, it was found that the growth of the strain decreased under limited nitrogen conditions. After harvesting, the cells in the supernatant were regenerated by cultivating in double concentration of media. The results showed that on the 12<sup>th</sup> day, the highest lipid content was achieved, reaching  $68.0 \pm 2.3 \text{ wt.}\%$  of the dry weight under nitrogen limitation. The composition of the obtained lipid consisted of 18.0 wt.% saturated fatty acids, 72.38 wt.% monounsaturated fatty acids, and 8.95 wt.% polyunsaturated fatty acids. The lipid composition was dominated by monounsaturated oleic acid ( $70.97 \pm 4.6 \text{ wt.}\%$ ), saturated palmitic acid ( $13.68 \pm 1.0 \text{ wt.}\%$ ), and polyunsaturated linoleic acid ( $8.41 \pm 0.4 \text{ wt.}\%$ ). The findings of this study demonstrated that limiting nitrogen availability enhanced lipid accumulation in *Nannochloropsis* sp. Despite experiencing inhibited growth, this microalga species shows great potential for the large-scale cultivation of lipids and essential fatty acids. These results contribute to understanding the relationship between nitrogen availability and lipid metabolism in microalgae, providing valuable insights for optimizing lipid production in the context of biofuel and biotechnology applications.

## 1. Introduction

Microalgae offer a significant advantage over edible crops by mitigating the adverse effects of human food consumption and providing a sustainable and carbon-neutral alternative to petroleum fuels. While biodiesel derived from edible

crops falls short in meeting the current demand for transportation fuels due to direct competition with food resources, microalgae present a promising solution as they can be grown on nonarable land [1, 2]. Microalgal biochemical compositions and growth are greatly influenced by the availability of nutrients such as nitrogen and phosphorus,

by which the limitation and depletion of nitrogen in the cultivation media can limit its growth with concurrent increases in lipid accumulation [3]. *Nannochloropsis* sp. is capable of accumulating triacylglycerols when exposed to nutrient limitation and responses to carbon dioxide fluctuation [4] particularly nitrate (N) and is therefore considered a promising organism for essential oil [5]. Moreover, *Nannochloropsis* sp. exhibits high biomass and lipid production potential under various stress conditions [6] and has been shown to be capable of growth in palm oil mill effluent [7–9].

Various species of *Nannochloropsis* have been extensively studied for lipid production due to their high lipid content of 37–64% [8, 10–14]. The interconnectedness between the environment and the activity of microalgae and cyanobacteria is crucial for environmental sustainability. Over the years, algae and their immediate descendants have played a significant role in sequestering vast amounts of atmospheric carbon in fossil deposits [1, 15]. Microalgae removed huge tonnages of carbon from the atmosphere to sequester it in mineral deposits and fossilized carbon [16]. These renewable biomass resources serve as valuable inputs to produce various goods and services such as biofuel production, aquafeed, and high-value products [17, 18]. A continuous flow process is developed for the recovery of the biomass of the marine microalga *Nannochloropsis salina* using flocculation methods and recovering more than 85% of the biomass [19]. *Nannochloropsis* species have drawn attention in lipid and biofuel research, due to their rapid growth in open ponds or photobioreactors and their ability to grow in seawater with high lipid yields (up to 60% of dry weight) [20]. Raceway ponds can also produce large-scale production of algal biomass [21]. Several microalga strains, including *Chlorella vulgaris*, *Spirulina* sp., *Chlamydomonas reinhardtii*, *Dunaliella* sp., *Scenedesmus obliquus*, and *Coelastrum* sp., could accumulate significant amounts of protein, lipids, and carbohydrates, making them ideal feedstocks for the production of biofuel [22].

In a continuous culture system operating at a steady state with a consistent biomass concentration, it is crucial for the dilution rate to match the specific growth rate [2]. *Nannochloropsis* species can accumulate more than 48% of their biomass as lipids in nitrogen-limiting conditions [13, 23]. This enhancement and lipid accumulation is consistent with predictions of the metabolic model of the microalgae [23–26]. Therefore, the focus of this study is to investigate the ability of *Nannochloropsis* sp. which can accumulate lipid components and their survivability culture under different concentrations of nitrogen environment. Additionally, an unconventional approach was carried out where the supernatants were subjected to repleted nitrogen media and then provided 24 hours of continuous light and aeration to initiate new growth. Besides that, the composition of lipids was analyzed and quantified for biodiesel production through gas chromatography-mass spectroscopy.

## 2. Materials and Methods

**2.1. Strain and Culture Medium.** *Nannochloropsis* sp. was obtained from the Algae Culture Collection Center and

Laboratory, Faculty of Industrial Sciences and Technology, Universiti Malaysia Pahang Al-Sultan Abdullah, Pahang, Malaysia. The stock culture was maintained under sterile condition in 1 L Erlenmeyer flask with 500 mL standard f/2 medium with composition of  $\text{NaNO}_3$  ( $8.82 \times 10^{-4}$  M),  $\text{NaH}_2\text{PO}_4 \cdot \text{H}_2\text{O}$  ( $3.62 \times 10^{-5}$  M),  $\text{Na}_2\text{CO}_3$  ( $1.06 \times 10^{-4}$  M),  $\text{FeCl}_3 \cdot 6(\text{H}_2\text{O})$  ( $1.17 \times 10^{-5}$  M),  $\text{Na}_2(\text{EDTA}) \cdot 2(\text{H}_2\text{O})$  ( $1.17 \times 10^{-5}$  M),  $\text{CuSO}_4 \cdot 5(\text{H}_2\text{O})$  ( $3.93 \times 10^{-8}$  M),  $\text{Na}_2\text{MoO}_4$  ( $2.60 \times 10^{-8}$  M),  $\text{ZnSO}_4 \cdot 7(\text{H}_2\text{O})$  ( $7.65 \times 10^{-8}$  M),  $\text{CoCl}_2 \cdot 6(\text{H}_2\text{O})$  ( $4.20 \times 10^{-8}$  M),  $\text{MnCl}_2 \cdot 4(\text{H}_2\text{O})$  ( $9.10 \times 10^{-7}$  M), thiamine HCl ( $2.96 \times 10^{-7}$  M), biotin ( $2.05 \times 10^{-9}$  M), and cyanocobalamin ( $3.69 \times 10^{-10}$  M). The culture was grown under a light intensity of  $120 \mu\text{mol m}^{-2} \text{s}^{-1}$  and with a photoperiod of 16:8 h light-to-dark cycle, optimum temperature ( $25 \pm 1^\circ\text{C}$ ), and provided agitation with filtered air through an air pump to prevent sediment. For nitrate starvation experiments, various concentrations of  $\text{NaNO}_3$  were used, in addition to other standard compounds (0 M,  $4.41 \times 10^{-4}$ ,  $8.82 \times 10^{-4}$ , and  $1.76 \times 10^{-3}$  M for zero nitrate, depleted nitrate, standard nitrate, and repleted nitrate, respectively).

**2.2. Experimental Conditions.** To determine the effect of nitrogen concentration on the growth, biomass, and lipid productivity of *Nannochloropsis* sp., cultures were grown for 14 days in batch mode. The f/2 medium was prepared with different concentrations of nitrate, for repleted experiments  $1.76 \times 10^{-3}$  M (double the amount as standard), for standard experiments  $8.82 \times 10^{-4}$  M, for limited experiments  $4.41 \times 10^{-4}$  M (half the amount as standard), and for N-free experiments zero nitrate, while other nutrients have remained in the concentration as stated above. The stock *Nannochloropsis* sp. culture was diluted with distilled water before being inoculated with each specific formulated f/2 medium at a 1:3 ratio ( $v v^{-1}$ ) for the experiment. The initial cell concentration of each treatment was approximately  $2.5 \times 10^5$  microalgal cells per milliliter. The environmental conditions such as light, agitation, and temperature were provided for the culture same as mentioned above.

**2.3. Growth Measurement.** Cell counts were determined using a hemocytometer, while optical density values at a wavelength of 680 nm were measured every two days using a Genesis UV-VIS spectrophotometer. After a 14-day period, the biomass was harvested by centrifugation at  $2500 \times g$  for 10 minutes. The wet biomass was then rinsed with distilled water and subjected to another round of centrifugation. Before and after drying the biomass at  $70^\circ\text{C}$  for 24 hours, its weight was measured to ensure a constant weight. After the harvesting process, an unconventional approach was carried out, in which the supernatants were subjected to repleted nitrogen media and then provided 24 hours of continuous light and aeration to initiate the growth cycle of *Nannochloropsis* sp. The next growth of the cycle was monitored and studied to regenerate cells' growth rate for continuous biomass production. In this approach, continuous cycles of cultivation can be carried out, with less biomass wastage.

**2.4. Lipid Extraction.** Lipids were extracted using methods described in Bligh and Dyer [27] and Smedes and Thomasen assisted with an ultrasound technique [28–30]. Dry biomass was weighed and dissolved in 50 mL of hexane (1:8,  $w v^{-1}$ ) solution in a Falcon tube. The mixture was vortexed for 30 sec. The tube was placed in a water bath at  $70 \pm 2^\circ\text{C}$  for 60 min to disrupt the cells. The mixture was then centrifuged at  $3293 \times g$ ,  $10^\circ\text{C}$  for 5 min, and the supernatant was collected. This step was repeated by adding 50 mL of hexane solvent to the biomass until the solution became colorless [29]. The supernatant containing the extracted lipid was collected after filtering through  $0.45 \mu\text{m}$  PTFE disk filters. The hexane solvent was separated from the mixture, and the extract was evaporated at  $50^\circ\text{C}$ ,  $269 \times g$  for 15 min by using a rotary evaporator. The separated lipid was weighed and collected in 2 mL of the centrifuged tube. The extracted lipid was transesterified by using hexane/methanolic-KOH (2:1, 1%) [31]. The mixture was heated at  $70^\circ\text{C}$  for 10 min and vortex for 30 sec. Two phases were formed in the mixture and the top phase by pipetting in a vial [32].

**2.5. Fatty Acid Methyl Ester Analysis.** Gas chromatography-mass spectrometry was used to detect fatty acid methyl esters content. Fatty acid methyl esters (FAMES) were measured and analyzed in an Agilent 7890A gas chromatography (GC) system equipped with a capillary mega wax MS column (30 m length  $\times$  0.32 mm diameter  $\times$  0.50  $\mu\text{m}$  film thickness). The mass spectrometer detector used helium as the carrier gas at  $1.0 \text{ mL min}^{-1}$ . The oven is programmed with the following time-temperature program:  $190^\circ\text{C}$  (2 min),  $190\text{--}230^\circ\text{C}$  ( $5^\circ\text{C min}^{-1}$ ), and  $230^\circ\text{C}$  (2 min). The mass spectra were recorded at 70 eV. Mass range was 40–250  $\text{mz}^{-1}$ . About 40 mg of fatty acid methyl ester sample was weighed in the vial; 1000  $\mu\text{L}$  of internal standard dilute on octane and  $15 \text{ mg mL}^{-1}$  concentration were added. In homogenous mixture form, sample (1  $\mu\text{L}$ ) was injected into GC. The inlet temperature was maintained at  $230^\circ\text{C}$  with a split ratio of 50:1 selected. The individual peaks of FAMES were identified through the chromatographic peaks by comparing retention times with those of standards [33].

**2.6. Statistical Analysis.** All the experiments were carried out in triplicate ( $n = 3$ ), and the observed standard deviations were reported accordingly.

### 3. Results and Discussions

**3.1. Effect of Nitrate Concentration.** The effect of varying nitrate concentration on the growth of *Nannochloropsis* sp. (Figure 1) monitored through cell counts and OD at 680 nm is illustrated in Figure 2. The growth of cells over the day period was higher in the standard concentration of  $f/2$  media when compared to the other nitrate treatments. The growth curve exhibited a lag phase on the first day, followed by an exponential phase starting from the second day, during which cell multiplication was rapid until day 12. Subsequently, the growth entered a stationary phase and showed a rapid decline on the 13<sup>th</sup> and 14<sup>th</sup> days in both the standard and repleted medium cultures. Both cell counts

and optical density (OD) measurements confirmed that the standard nitrate concentration resulted in the highest growth, followed by repleted nitrate, limited nitrate, and finally nitrate-free media.

On the other hand, for a continuous cycle of cultivation, the supernatant was aerated and kept at the same temperature and light environment. The result found that the growth of the cell was still present, and it was boosted by adding double the concentration of media. The minimum amount of nutrients present in the supernatant initiates the growth cell. However, the addition of media increases biomass productivity at optimum conditions. This helps to minimize the usage of water reduce the loss of nutrients and reduce the cost of cultivation for biomass production.

Based on Figure 3, the highest dry biomass yield was obtained from standard media ( $1234 \pm 35.6 \text{ mg L}^{-1}$ ) followed by repleted-N ( $1011 \pm 32.1 \text{ mg L}^{-1}$ ), limited-N ( $850 \pm 25.4 \text{ mg L}^{-1}$ ), and free-N ( $765 \pm 35.6 \text{ mg L}^{-1}$ ) media after 14-day cultivation period. On the other hand, the lipid productivity trend differed in the order of limited-N ( $68.0 \pm 2.3 \text{ wt.}\%$ ), then free-N ( $62.3 \pm 2.9 \text{ wt.}\%$ ), standard-N ( $51.2 \pm 2.5 \text{ wt.}\%$ ), and finally repleted-N ( $48.1 \pm 2.8 \text{ wt.}\%$ ). The optimum productivity of biomass and lipids is shown in Table 1. The results obtained in this study can be compared with previous studies mentioned in Table 2. *Nannochloropsis* sp. showed the least growth and biomass production in limited-N media, likely due to a reduction in their light harvesting efficiency and energy transduction, leading to decreased photosynthetic efficacy. Nitrogen deficiency in microalgae causes the loss of their photosynthetic efficacy of antenna, chlorophyll a, and photosystem II completely [34]. In response to nitrogen deficiency, microalgae can boost light intake and divert carbon metabolism toward lipid synthesis and carbohydrate storage. Intracellular nitrogen is re-assigned through reducing biosynthetic pathways under sustained low nitrogen stress. Under nitrogen deficiency, protein synthesis and cell development both slowed [35, 36]. A similar result mentioned that the cellular density reduced when the  $\text{NaNO}_3$  was reduced, where the photosynthetic efficiency was inhibited due to the response of microalgae to nitrogen limitation. The reason behind this condition is the relationship between pigment synthesis and nitrogen metabolism. However, increasing the  $\text{NaNO}_3$  enhances the biomass dry weight cell of *Chlorella* sp. [37]. Nitrogen in the forms of nitrate, nitrite, urea, and ammonium can be assimilated by microalgae. Green microalgae favor ammonium because they can easily use it from a metabolic standpoint. Indeed, a series of enzymatic pathways is required to transform other forms [38]. A study [39] investigated the nitrogen limitation on *Nannochloropsis oceanica*, *Isochrysis aff. galbana clone T-Iso*, *Rhodomonas baltica*, and *Phaeodactylum tricornerutum*. As a result of response to severe nitrogen limitation, all four species accumulated lipids, primarily in the form of TAG. In moderate nitrogen limitation, *N. oceanica* accumulated 51% of the dry weight as a lipid, up to 87% of the fatty acids in TAG. The only species where the fraction of polyunsaturated fatty acids, particularly the fraction of docosahexaenoic acid, was found was *Isochrysis aff. galbana clone T-Iso*. Another study by [40] found that under






FIGURE 1: *Nannochloropsis* sp. photomicrographs under a fluorescence microscope ( $\times 1000$ ) (a) and image under a field emission scanning electron microscope ( $\times 5500$ ) (b).



(a)



(b)

FIGURE 2: Growth curve (a) by cell counts and optical density (at 680 nm) (b) of *Nannochloropsis* sp. cultivated under various concentrations of nitrate.






FIGURE 3: The final dry biomass (mg L<sup>-1</sup>) and lipid productivity (%) of *Nannochloropsis* sp.

TABLE 1: Biomass and lipid outputs of *Nannochloropsis* sp. under N-starvation ( $4.41 \times 10^{-4}$  M) condition. Conditions: light intensity  $120 \mu\text{mol photon m}^{-2} \text{s}^{-1}$ , temperature  $25 \pm 2^\circ\text{C}$ , and standard *f/2* media with shaking 80 rpm.

Parameter	Value
Biomass concentration (mg L <sup>-1</sup> )	$850.3 \pm 25.4$
Biomass productivity (mg L <sup>-1</sup> d <sup>-1</sup> )	$60.73 \pm 0.9$
Lipid in biomass (% <i>w w</i> <sup>-1</sup> )	$68.0 \pm 2.3$

lower light and limited nitrogen conditions, the starch content of *Pseudochlorococcum* increased in the first 4 days and then began to decrease slightly on day 10. However, neutral lipid content begins to accumulate at day 10. Under high light with limited nitrogen, *Pseudochlorococcum* showed rapid growth and high nitrate reductase activity. Therefore, starch acts as the main carbon storage which can indicate that *Pseudochlorococcum* applies starch as the primary carbon storage product when it has grown under sufficient nitrogen conditions [40]. Therefore, the rate at which nitrogen (supplied as  $\text{NaNO}_3$ ) is consumed during the synthesis of starch and lipids was monitored by measuring the concentration of nitrogen in the growth medium and the activity of nitrate reductase because the ability to take up and assimilate nitrogen frequently determines algal growth and basal metabolism.

**3.2. Transesterification and Methyl Ester Analysis.** Figure 4 depicts the GC chromatogram of transesterified lipid that is extracted from *Nannochloropsis* sp. The extraction and transesterification processes showed high efficiency and yielded the successful recovery of fatty acids. Microalgae have emerged as a possible source of highly unsaturated fatty acids, and *Nannochloropsis* sp. is observed as one of the most promising microalgae due to its fatty acid profile. Under nitrate limitation, the highest fatty acid content (oleic acid) from the lower biomass specific was obtained. The current work demonstrates that the fatty acyl compositions of the various major lipid classes are also distinctive. Fatty acid

methyl esters are the main component of biodiesel, a sustainable kind of transportation fuel. The size distribution and degree of unsaturation in FAME profiles, in particular, may have a major impact on the physical and chemical characteristics of biodiesel [45].

Table 3 compares the result of detected fatty acids of C16:0, C18:0, C20:0, C16:1, C18:1, C20:1, C18:2, and C18:3 under various concentrations of nitrite media. Oleic acid (C18:1,  $70.97 \pm 4.6\%$ ), palmitic fatty acid (C16:0,  $13.68 \pm 1.0\%$ ), and followed by linoleic acid (C18:2,  $8.41 \pm 0.4\%$ ) are the most dominant fatty acids in each subgroup of fatty acids. Wang et al. [39] found that high percentage of oleic acid (C18:1) and other dominating fatty acids such as 14:0, 16:0, 16:1, and 20:5 in the fatty acid profile of *Nannochloropsis oceanica*. Hindarso et al. [49] reported that methyl palmitic 35.21% and methyl  $\gamma$  15.98% weight as the highest composition of methyl ester from the *Nannochloropsis*. Since oleic acid (39.51%) and palmitoleic acid (31.24%) are the dominant components in the extracted oil, these unsaturated fatty acids will improve the cold performance of biodiesel. In turn, the produced biodiesel will be comparatively more suitable for export to cold climate regions. According to earlier research, the best biodiesel would have relatively low quantities of saturated fatty acids and polyunsaturated fatty acids to reduce oxidative stability and cold flow issues [45].


Biodiesel quantification was done according to EN14103 in which methyl heptadecanoate ( $\text{C}_{17}$ ) was used as an internal standard. Peaks of methyl esters were identified by comparing them to their respective standards.

$$\text{Methyl ester content (wt.\%)} = \frac{(\sum A_i) - A_{IS}}{A_{IS}} \times \frac{C_{IS} - V_{IS}}{m} \times 100, \quad (1)$$

where  $\sum A_i$  represents the sum of methyl ester peak area,  $A_{IS}$  is the area of internal standard,  $C_{IS}$  is the concentration of the internal standard in heptane (mg/mL),  $V_{IS}$  is the volume of internal standard (mL), and  $m$  is the weight of the sample. Accordingly, GC results confirmed that the transesterification reaction has yielded methyl ester conversion of  $98.7 \pm$

TABLE 2: The biomass and lipid productivities of various strains of *Nannochloropsis* sp. from previous work.

Species	Maximum biomass (mg L <sup>-1</sup> )	Lipid productivity (mg L <sup>-1</sup> d <sup>-1</sup> )	References
<i>Nannochloropsis</i> sp.	360	48	This study
<i>Nannochloropsis gaditana</i>	185	185	[41]
<i>Nannochloropsis oculata</i>	497	151	[42]
<i>Nannochloropsis salina</i>	610	228	[43]
<i>Nannochloropsis</i> sp.	440	108	[44]
<i>Nannochloropsis oceanica</i> IMET1	301 ± 0.02	158 ± 0.013	[43]
<i>Nannochloropsis granulate</i> CCMP525	216 ± 0.01	130 ± 0.037	[45]
<i>Nannochloropsis oceanica</i> NIOF15/001	75	28	[46]
<i>Nannochloropsis limnetica</i>	640		
	490 ± 0.01	173	[47]
<i>Nannochloropsis gaditana</i> 1049	540 ± 0.013	289 ± 0.02	[48]
<i>N. oculata</i> NCTU-3	480 ± 0.029	142 ± 0.049	[42]

FIGURE 4: GC chromatogram depicting fatty acid composition of *Nannochloropsis* sp.TABLE 3: Fatty acid quantification (%  $w w^{-1}$ ) of lipid extracted from *Nannochloropsis* sp.

Fatty acids	Amount (% $w w^{-1}$ )			
	Nitrogen limited	Nitrogen repleted	Standard nitrogen	Nitrogen free
<i>Saturated fatty acid</i>				
Palmitic acid (C16:0)	13.68 ± 0.14	15.73 ± 0.32	15.44 ± 0.12	14.85 ± 0.22
Stearic acid (C18:0)	4.90 ± 0.30	3.72 ± 0.1	3.34 ± 0.01	3.76 ± 0.32
Eicosanoic acid (C20:0)	0.45 ± 0.21	0.47 ± 0.12	0.76 ± 0.13	0.69 ± 0.01
<i>Monounsaturated fatty acid</i>				
Palmitoleic acid (C16:1)	31.24 ± 0.3	32.75 ± 0.21	33.38 ± 0.35	33.54 ± 0.54
Oleic acid (C18:1)	39.51 ± 0.6	37.81 ± 0.3	35.95 ± 0.25	36.71 ± 0.64
Eicosenoic acid (C20:1)	0.27 ± 0.2	0.22 ± 0.20	0.18 ± 0.12	0.21 ± 0.30
<i>Polyunsaturated fatty acid</i>				
Linoleic acid (C18:2)	9.41 ± 0.4	8.83 ± 0.51	10.53 ± 0.32	9.73 ± 0.43
Linolenic acid (C18:3)	0.54 ± 0.01	0.47 ± 0.21	0.42 ± 0.52	0.51 ± 0.34

Data reported as mean ± SD.  $n = 3$  repetitions.

0.4 wt.%, surplus well the minimum requirements of 96.5 wt.% methyl ester content as per EN14103 standard.

#### 4. Conclusion

This work evaluated the locally isolated oleaginous *Nannochloropsis* sp. by imposing nitrate starvation. Results show that N-starvation gives a positive response in accumulating lipids, especially the content of oleic acid (~71 wt.%). After optimization (light intensity  $120 \mu\text{mol photon m}^{-2} \text{s}^{-1}$ , temperature  $25 \pm 2^\circ\text{C}$ , and standard f/2 media with continuous air-CO<sub>2</sub> while shaking at 80 rpm), in 14-day cycle cultivations, the highest lipid in biomass was  $68.0 \pm 2.3 \text{ wt.}\%$ , while maximum biomass concentration was  $850.3 \pm 25.4 \text{ mg L}^{-1}$  and biomass productivity stands at  $60.73 \pm 0.9 \text{ mg L}^{-1} \text{ d}^{-1}$ . The work proves that although lower biomass is produced under N-limitation, lipid content was considerably higher. It was found that nitrogen starvation has enhanced lipid production; however, it reduces the growth rate (total biomass) obtained. In order to further the next cultivation cycle, a higher (three times higher) concentration of standard media needs to be added to the culture for continuous growth. Therefore, *Nannochloropsis* sp. has great potential for biomass for biodiesel production, which facilitates exporting the produced biodiesel to cold climate regions for better cold performance due to higher unsaturated fatty acids.

#### Data Availability

The research data used to support the findings of this study are included in the article.

#### Conflicts of Interest

The authors declare that they have no conflicts of interest.

#### Acknowledgments

The authors gratefully acknowledge University of Malaysia Pahang Al Sultan Abdullah (UMPSA) for financial support through Flagship Research Grant (RDU182205), Internal Research Grant (RDU2203101), and Postgraduate Research Grants Scheme (PGRS220376).

#### References

- [1] Y. Chisti, "Biodiesel from microalgae," *Biotechnology Advances*, vol. 25, no. 3, pp. 294–306, 2007.
- [2] Y. Chisti, "Fermentation technology," in *Industrial biotechnology: sustainable growth and economic success*, W. Soetaert and E. J. Vandamme, Eds., pp. 149–171, Wiley-VCH, New York, 2010.
- [3] M. A. Yaakob, R. M. S. R. Mohamed, A. Al-Gheethi, R. A. Gokare, and R. R. Ambati, "Influence of nitrogen and phosphorus on microalgal growth, biomass, lipid, and fatty acid production: an overview," *Cells*, vol. 10, no. 2, p. 393, 2021.
- [4] L. Wei, B. Liu, D. Liu, Z. Xu, R. Wang, and W. Zhang, "Identification and expression analysis of genome-wide long non-coding RNA responsive CO<sub>2</sub> fluctuated environment in marine microalga *Nannochloropsis oceanica*," *Marine Pollution Bulletin*, vol. 176, no. 113419, 2022.
- [5] T. Castillo, D. Ramos, T. García-Beltrán, M. Brito-Bazan, and E. Galindo, "Mixotrophic cultivation of microalgae: an alternative to produce high-value metabolites," *Biochemical Engineering Journal*, vol. 176, p. 108183, 2021.
- [6] N. Govindan, G. P. Maniam, A. Z. Sulaiman, A. Ajit, T. Chatsungnoen, and Y. Chisti, "Production of renewable lipids by the diatom *Amphora* copulate," *Fermentation*, vol. 7, no. 1, p. 37, 2021.
- [7] Y. Chisti, "Constraints to commercialization of algal fuels," *Journal of Biotechnology*, vol. 167, no. 3, pp. 201–214, 2013.
- [8] P. Paramasivam, K. Kanagesan, P. Bhuyar, N. Govindan, and G. P. Maniam, "Biomass and lipid production from indigenous *Nannochloropsis* sp. by employing stress factors for improved biodiesel production," *Environment, Development and Sustainability*, pp. 1–15, 2021.
- [9] P. Paramasivam, K. M. Palanisamy, S. Jayakumar, N. Govindan, M. H. A. Rahim, and G. P. Maniam, "Lipid production from *Nannochloropsis* sp. grown in palm oil mill effluent," *In IOP Conference Series: Earth and Environmental Science*, vol. 641, no. 1, p. 12021, 2021.
- [10] R. Kandilian, J. Pruvost, J. Legrand, and L. Pilon, "Influence of light absorption rate by *Nannochloropsis oculata* on triglyceride production during nitrogen starvation," *Bioresource Technology*, vol. 163, pp. 308–319, 2014.
- [11] X. N. Ma, T. P. Chen, B. Yang, J. Liu, and F. Chen, "Lipid production from *Nannochloropsis*," *Marine drugs*, vol. 14, no. 4, p. 61, 2016.
- [12] M. G. Sung, J. I. Han, B. Lee, and Y. K. Chang, "Wavelength shift strategy to enhance lipid productivity of *Nannochloropsis gaditana*," *Biotechnology for Biofuels*, vol. 11, no. 1, pp. 1–12, 2018.
- [13] T. Menegol, G. I. Romero-Villegas, M. López-Rodríguez et al., "Mixotrophic production of polyunsaturated fatty acids and carotenoids by the microalga *Nannochloropsis gaditana*," *Journal of Applied Phycology*, vol. 31, no. 5, pp. 2823–2832, 2019.
- [14] J. Y. Schambach, A. M. Finck, P. Kitin et al., "Growth, total lipid, and omega-3 fatty acid production by *Nannochloropsis* spp. cultivated with raw plant substrate," *Algal Research*, vol. 51, article 102041, 2020.
- [15] Y. Chisti, *Microalgae biotechnology: a brief introduction*, Academic Press, 2020.
- [16] Y. Chisti, *Society and microalgae: understanding the past and present*, Academic Press, 2018.
- [17] Y. Chisti, *Biorefinery: Integrated Sustainable Processes for Biomass Conversion to Biomaterials, Biofuels, and Fertilizers*, J.-R. Bastidas-Oyanedel and J. E. Schmidt, Eds., Springer, Cham, Switzerland, 2019.
- [18] Y. Chisti, *Introduction to algal fuels*, Elsevier, 2019.
- [19] T. Chatsungnoen and Y. Chisti, "Continuous flocculation-sedimentation for harvesting *Nannochloropsis salina* biomass," *Journal of Biotechnology*, vol. 222, pp. 94–103, 2016.
- [20] Z. Y. Du, J. Alvaro, B. Hyden et al., "Enhancing oil production and harvest by combining the marine alga *Nannochloropsis oceanica* and the oleaginous fungus *Mortierella elongata*," *Biotechnology for Biofuels*, vol. 11, no. 1, pp. 1–16, 2018.
- [21] Y. Chisti, "Large-scale production of algal biomass: raceway ponds," in *In Algal biotechnology*, pp. 21–40, Springer, Cham, 2016.

- [22] S. Khan, P. Das, M. A. Quadir et al., "Microalgal feedstock for biofuel production: recent advances, challenges, and future perspective," *Fermentation*, vol. 9, no. 3, p. 281, 2023.
- [23] Y. Chisti, "Raceways-based production of algal crude oil," *Green*, vol. 3, no. 3-4, pp. 195–216, 2013.
- [24] P. Bhuyar, S. Sundararaju, M. H. A. Rahim, R. Ramaraj, G. P. Maniam, and N. Govindan, "Microalgae cultivation using palm oil mill effluent as growth medium for lipid production with the effect of CO<sub>2</sub> supply and light intensity," *Biomass Conversion and Biorefinery*, vol. 11, no. 5, pp. 1555–1563, 2021.
- [25] T. C. Pedersen, R. D. Gardner, R. Gerlach, and B. M. Peyton, "Assessment of *Nannochloropsis gaditana* growth and lipid accumulation with increased inorganic carbon delivery," *Journal of Applied Phycology*, vol. 30, no. 4, pp. 2155–2166, 2018.
- [26] D. Simionato, M. A. Block, N. La Rocca et al., "The response of *Nannochloropsis gaditana* to nitrogen starvation includes de novo biosynthesis of triacylglycerols, a decrease of chloroplast galactolipids, and reorganization of the photosynthetic apparatus," *Eukaryotic Cell*, vol. 12, no. 5, pp. 665–676, 2013.
- [27] F. Smedes and T. K. Thomassen, "Evaluation of the Bligh & Dyer lipid determination method," *Marine Pollution Bulletin*, vol. 32, no. 8-9, pp. 681–688, 1996.
- [28] H. S. Basily, M. M. Nassar, G. I. El Diwani, and S. A. A. El-Enin, "Extraction of algal lipid as a natural cosmetic component," *Egyptian Pharmaceutical Journal*, vol. 17, no. 1, pp. 13–20, 2018.
- [29] M. Mahali and G. Sibi, "Extraction methods and functional properties of protein from *Arthrospira platensis* for bioavailability of algal proteins," *International Journal of Pharmacy and Chemistry*, vol. 5, no. 2, pp. 20–25, 2019.
- [30] S. Jayakumar, L. Y. Peng, M. H. A. Rahim, G. P. Maniam, and N. Govindan, "Studies about the effect of different photosynthetic light intensity on two isolated species of marine microalgae (diatom) for high lipid production," *Advances in Environmental Biology*, vol. 9, no. 27, pp. 332–340, 2015.
- [31] B. P. Lim, G. P. Maniam, and S. A. Hamid, "Biodiesel from adsorbed waste oil on spent bleaching clay using CaO as a heterogeneous catalyst," *European Journal of Scientific Research*, vol. 33, no. 2, pp. 347–357, 2009.
- [32] J. Y. Lee, C. Yoo, S. Y. Jun, C.-Y. Ahn, and H. M. Oh, "Comparison of several methods for effective lipid extraction from microalgae," *Bioresource Technology*, vol. 101, no. 1, pp. S75–S77, 2010.
- [33] K. M. Palanisamy, P. Paramasivam, G. P. Maniam, M. H. A. Rahim, N. Govindan, and Y. Chisti, "Production of lipids by *Chaetoceros affinis* in media based on palm oil mill effluent," *Journal of Biotechnology*, vol. 327, pp. 86–96, 2021.
- [34] C. Latsos, J. V. Houcke, and K. R. Timmermans, "The effect of nitrogen starvation on biomass yield and biochemical constituents of *Rhodomonas* sp.," *Frontiers in Marine Science*, vol. 7, 2020.
- [35] M. K. Enamala, S. Enamala, M. Chavali et al., "Production of biofuels from microalgae - a review on cultivation, harvesting, lipid extraction, and numerous applications of microalgae," *Renewable and Sustainable Energy Reviews*, vol. 94, pp. 49–68, 2018.
- [36] D. de Jesús-Campos, J. A. López-Elías, L. Á. Medina-Juarez, G. Carvallo-Ruiz, D. Fimbres-Olivarria, and C. Hayano-Kanashiro, "Chemical composition, fatty acid profile and molecular changes derived from nitrogen stress in the diatom *Chaetoceros muelleri*," *Aquaculture Reports*, vol. 16, article 100281, 2020.
- [37] A. Abomohra, M. Li, S. Faisal, L. Li, and M. Elsayed, "Maximizing nitrogen removal and lipid production by microalgae under mixotrophic growth using response surface methodology: towards enhanced biodiesel production," *Fermentation*, vol. 8, no. 12, p. 682, 2022.
- [38] V. Pozzobon, N. Cui, A. Moreaud, E. Michiels, and W. Levasseur, "Nitrate and nitrite as mixed source of nitrogen for *Chlorella vulgaris*: growth, nitrogen uptake and pigment contents," *Bioresource Technology*, vol. 330, article 124995, 2021.
- [39] X. Wang, H. K. Fosse, K. Li, M. S. Chauton, O. Vadstein, and K. I. Reitan, "Influence of nitrogen limitation on lipid accumulation and EPA and DHA content in Four Marine microalgae for possible use in aquafeed," *Frontiers in Marine Science*, vol. 6, no. 95, 2019.
- [40] Y. Li, W. Zhou, B. Hu, M. Min, P. Chen, and R. R. Ruan, "Integration of algae cultivation as biodiesel production feedstock with municipal wastewater treatment: strains screening and significance evaluation of environmental factors," *Bioresource Technology*, vol. 102, no. 23, pp. 10861–10867, 2011.
- [41] M. Cecchin, S. Berteotti, S. Paltrinieri et al., "Improved lipid productivity in *Nannochloropsis gaditana* in nitrogen-replete conditions by selection of pale green mutants," *Biotechnology for Biofuels*, vol. 13, no. 1, pp. 1–14, 2020.
- [42] S. Y. Chiu, C. Y. Kao, M. T. Tsai, S. C. Ong, C. H. Chen, and C. S. Lin, "Lipid accumulation and CO<sub>2</sub> utilization of *Nannochloropsis oculata* in response to CO<sub>2</sub> aeration," *Bioresource Technology*, vol. 100, no. 2, pp. 833–838, 2009.
- [43] E. M. Fakhry and D. M. El Maghraby, "Lipid accumulation in response to nitrogen limitation and variation of temperature in *Nannochloropsis salina*," *Botanical Studies*, vol. 56, no. 1, pp. 1–8, 2015.
- [44] X. Peng, F. Meng, Y. Wang, X. Yi, and H. Cui, "Effect of pH, temperature, and CO<sub>2</sub> concentration on growth and lipid accumulation of *Nannochloropsis* sp. MASCC 11," *Journal of Ocean University of China*, vol. 19, no. 5, pp. 1183–1192, 2020.
- [45] Y. Ma, Z. Wang, C. Yu, Y. Yin, and G. Zhou, "Evaluation of the potential of 9 *Nannochloropsis* strains for biodiesel production," *Bioresource Technology*, vol. 167, pp. 503–509, 2014.
- [46] M. Ashour, M. E. Elshobary, R. El-Shenody, A. W. Kamil, and A. E. F. Abomohra, "Evaluation of a native oleaginous marine microalga *Nannochloropsis oceanica* for dual use in biodiesel production and aquaculture feed," *Biomass and Bioenergy*, vol. 120, pp. 439–447, 2019.
- [47] N. Arora, E. Lo, and G. P. Philippidis, "A two-prong mutagenesis and adaptive evolution strategy to enhance the temperature tolerance and productivity of *Nannochloropsis oculata*," *Bioresource Technology*, vol. 364, article 128101, 2022.
- [48] Q. Hu, W. Xiang, S. Dai et al., "The influence of cultivation period on growth and biodiesel properties of microalga *Nannochloropsis gaditana* 1049," *Bioresource Technology*, vol. 192, pp. 157–164, 2015.
- [49] H. Hindarso, A. Aylilianawati, and M. E. Sianto, "Biodiesel production from the microalgae *nannochloropsis* by microwave using CaO and MgO catalysts," *International Journal of Renewable Energy Development*, vol. 4, no. 1, pp. 72–76, 2015.