

# Compatible Pair of Nontrivial Actions for Some Cyclic Groups of 2-Power Order

Sahimel Azwal Sulaiman<sup>a</sup>, Mohd Sham Mohamad<sup>b</sup> and Yuhani Yusof<sup>c</sup>

<sup>a,b,c</sup>Futures and Trends Research Groups, Faculty of Industrial Sciences & Technology,  
Universiti Malaysia Pahang, Lebuhraya Tun Razak, 26300 Gambang, Kuantan, Pahang Darul Makmur  
<sup>a</sup>titus1704@yahoo.com, <sup>b</sup>mohdsham@ump.edu.my, <sup>c</sup>yuhani@ump.edu.my

**Abstract.** Compatible action played a very important verification before the nonabelian tensor product can be computed. This paper gives the some exact number of compatible pairs of actions for some cyclic groups of 2-power order. Some necessary and sufficient number theoretical conditions for a pair of cyclic groups of 2-power order with nontrivial actions act compatibly on each other are used to investigate some properties to find the exact number of compatible pairs of actions. Algorithms in Groups, Algorithms and Programming (GAP) software are constructed to create more examples on selected case. New results on compatible pair of nontrivial actions of order two and four are given.

**Keywords:** Compatible Actions, Cyclic Groups, Nonabelian Tensor Product

## INTRODUCTION

Let  $G$  and  $H$  be groups which act on each other and each of which acts on itself by conjugation, then the actions are compatible if  ${}^{(s_h)}g' = {}^s({}^h(g^{-1}g'))$  and  ${}^{(h_g)}h' = {}^h({}^s(h^{-1}h'))$ . This compatible action played a very important verification before the nonabelian tensor product can be computed. Brown and Loday [1] were the first to introduce the concept of the nonabelian tensor product of groups with the compatible actions. Compatible conditions are the prerequisite before nonabelian tensor product can be calculated.

This construction of the nonabelian tensor product has its origins in the algebraic K-theory and in homotopy theory. Several authors, starting with Brown, Johnson and Robertson [2], have studied group theoretical aspects of nonabelian tensor products extensively. A paper by Kappe [3] gives an overview of known results and literature.

Many authors are working with nonabelian tensor square (where  $H = G$ ), and the compatibly conditions can be checked easily. It is interesting to check the compatible conditions when calculating tensor product since the groups are not the same or when the actions are nontrivial. Ellis and McDermott [4] had done this but specifically for quaternion group of order 32. Visscher [5] in 1998 had provided necessary and sufficient conditions for a pair of cyclic groups to act compatibly. He gave complete compatible conditions when one of the actions is trivial or both actions are trivial in cyclic groups of  $p$ -power order. In this research, all compatible actions involving at least one of the actions having order greater than two will be determined as a continuation from Visscher's.

Groups, Algorithms and Programming (GAP) software [6] is used to determine the conditions in which the actions satisfy the compatibility conditions in some finite cyclic groups.

## SOME PREPARATORY RESULTS ON COMPATIBILITY CONDITIONS

The nonabelian tensor product  $G \otimes H$  is defined only if  $g$  and  $h$  act on each other in such way that the actions satisfy certain compatibility condition. Start with the definition of an action a group  $g$  on a group  $h$ .

### Definition 2.1 [5]

Let  $G$  and  $H$  be groups. An action of  $G$  on  $H$  is a mapping,  $\Phi : G \rightarrow \text{Aut}(H)$  such that

$$\Phi(gg')(h) = \Phi(g)(\Phi(g')(h))$$

for all  $g, g' \in G$  and  $h \in H$ .

For a group  $g$  to act on a group  $h$ , by using the fact that the identity in  $g$  acts as the identity mapping on  $h$ , thus this implies that all elements in  $g$  act as automorphism of  $h$  on  $h$ . The definition of a compatible action is given as follows.

**Definition 2.1 [5]**

Let  $G$  and  $H$  be groups which act on each other. These mutual actions are said to be compatible with each other and with the actions of  $G$  and  $H$  on themselves by conjugation if

$$({}^s h)g' = {}^s ({}^h (g^{-1} g')) \text{ and } ({}^h s)h' = {}^h ({}^s (h^{-1} h'))$$

for all  $g, g' \in G$  and  $h \in H$ .

Since we are focusing on finite cyclic groups, the compatibility conditions which hold for abelian groups will be reduce to  $({}^s h)g' = {}^h g'$  and  $({}^h s)h' = {}^s h'$ , for all  $g, g' \in G$  and  $h \in H$ . For compatibility of actions of  $G$  and  $H$  on each other, it suffices for the compability conditions to hold on the generators of  $G$  and  $H$ . Visscher [7] has done this and he gave some necessary and sufficient number theoretical conditions for a pair of finite cyclic groups to act compatibly on each other as stated in Proposition 2.3.

**Proposition 2.3 [5]**

Let  $G = \langle x \rangle = C_m$  and  $H = \langle y \rangle = C_n$  be finite cyclic groups. Then there exist mutual actions of  $G$  and  $H$  on each other such that  ${}^y x = x^k$  and  ${}^x y = y^l$  for  $k, l \in \mathbb{Z}$  if and only if conditions (i) and (ii) below are satisfied. These action are compatible if and only condition (iii) is satisfied as well.

- (i)  $(k, m) = (l, n) = 1$
- (ii)  $k^n \equiv 1 \pmod m$  and  $l^m \equiv 1 \pmod n$
- (iii)  $k^{l-1} \equiv 1 \pmod m$  and  $l^{k-1} \equiv 1 \pmod n$ .

In the next section, all compatible actions involving at least one of the actions having order greater than two will be determined.

## COMPATIBLE CONDITIONS FOR NONTRIVIAL ACTION

In the previous section, we can see that Visscher had done the classification of compatibility condition focusing on cyclic groups and provide necessary and sufficient conditions for the general case without any specific condition.

Now, our aim is to determine the compatible conditions if the order of the action is more than two. All computations are doing by GAP. New GAP coding has been built to calculate the compatible actions.

Let  $G = \langle x \rangle = C_m$  and  $H = \langle y \rangle = C_m$  be finite cyclic groups. Let the actions of  $G$  and  $H$  on each other be such that  ${}^y x = x^k$  and  ${}^x y = y^l$  for  $k, l \in \mathbb{Z}$ . Now consider that both actions have order four. The compatible conditions that are satisfied by  $k$  and  $l$  are given in **Table 1** when  $G = H$  and in **Table 2** when  $G \neq H$ .

**TABLE 1.** Values of  $k$  and  $l$  in which the action of order four satisfying the compatible conditions when  $G = H$ .

Groups	$k$	$l$		Groups	$k$	$l$
$G = H = C_{2^4}$	5	5		$G = H = C_{2^7}$	33	33
	13	13			97	97
$G = H = C_{2^5}$	9	9		$G = H = C_{2^8}$	65	65
	25	25			193	193
$G = H = C_{2^6}$	17	17		$G = H = C_{2^9}$	129	129
	49	49			385	385

**TABLE 2.** Values of  $k$  and  $l$  in which the action of order four satisfying the compatible conditions when  $G \neq H$ .

Groups	$k$	$l$	Groups	$k$	$l$	Groups	$k$	$l$
$G = C_{2^4}$	5	9	$G = C_{2^5}$	9	5	$G = C_{2^6}$	17	5
and	5	25	and	9	13	and	17	13
$H = C_{2^5}$	13	9	$H = C_{2^4}$	25	5	$H = C_{2^4}$	49	5
	13	25		25	13		49	13
$G = C_{2^4}$	5	17	$G = C_{2^5}$	9	17	$G = C_{2^6}$	17	9
and	5	49	and	9	49	and	17	25
$H = C_{2^6}$	13	17	$H = C_{2^6}$	25	17	$H = C_{2^5}$	49	9
	13	49		25	49		49	25
$G = C_{2^4}$	5	33	$G = C_{2^5}$	9	33	$G = C_{2^6}$	17	33
and	5	97	and	9	97	and	17	97
$H = C_{2^7}$	13	33	$H = C_{2^7}$	25	33	$H = C_{2^7}$	49	33
	13	97		25	97		49	97
$G = C_{2^4}$	5	65	$G = C_{2^5}$	9	65	$G = C_{2^6}$	17	65
and	5	193	and	9	193	and	17	193
$H = C_{2^8}$	13	65	$H = C_{2^8}$	25	65	$H = C_{2^8}$	49	65
	13	193		25	193		49	193
$G = C_{2^4}$	5	129	$G = C_{2^5}$	9	129	$G = C_{2^6}$	17	129
and	5	385	and	9	385	and	17	385
$H = C_{2^9}$	13	129	$H = C_{2^9}$	25	129	$H = C_{2^9}$	49	129
	13	385		25	385		49	385
Groups	$k$	$l$	Groups	$k$	$l$	Groups	$k$	$l$
$G = C_{2^7}$	33	5	$G = C_{2^8}$	65	5	$G = C_{2^9}$	129	5
and	33	13	and	65	13	and	129	13
$H = C_{2^4}$	97	5	$H = C_{2^4}$	193	5	$H = C_{2^4}$	385	5
	97	13		193	13		385	13
$G = C_{2^7}$	33	9	$G = C_{2^8}$	65	9	$G = C_{2^9}$	129	9
and	33	25	and	65	25	and	129	25
$H = C_{2^5}$	97	9	$H = C_{2^5}$	193	9	$H = C_{2^5}$	385	9
	97	25		193	25		385	25
$G = C_{2^7}$	33	17	$G = C_{2^8}$	65	17	$G = C_{2^9}$	129	17
and	33	49	and	65	49	and	129	49
$H = C_{2^6}$	97	17	$H = C_{2^6}$	193	17	$H = C_{2^6}$	385	17
	97	49		193	49		385	49
$G = C_{2^7}$	33	65	$G = C_{2^8}$	65	33	$G = C_{2^9}$	129	33
and	33	193	and	65	97	and	129	97
$H = C_{2^8}$	97	65	$H = C_{2^7}$	193	33	$H = C_{2^7}$	385	33
	97	193		193	97		385	97
$G = C_{2^7}$	33	129	$G = C_{2^8}$	65	129	$G = C_{2^9}$	129	65
and	33	385	and	65	385	and	129	193
$H = C_{2^9}$	97	129	$H = C_{2^9}$	193	129	$H = C_{2^8}$	385	65
	97	385		193	385		385	193

This lead the following results:

**Theorem 3.1**

Let  $C_{2^\alpha} = \langle x \rangle$  and  $C_{2^\beta} = \langle y \rangle$  be cyclic groups of 2-power order. Furthermore, let  $C_{2^\alpha}$  and  $C_{2^\beta}$  act on each other so that  ${}^y x = x^k$  and  ${}^x y = y^l$  for  $k, l \in \mathbb{N}$  with  $(k, 2) = (l, 2) = 1$ . Then  $C_{2^\alpha}$  and  $C_{2^\beta}$  act compatibly on each other when  $\alpha = \beta$  with  ${}^{y^4} x = x$  and  ${}^{x^4} y = y$ , if and only if  $k$  and  $l$  are congruent to one of the following:

- (i)  $k = l \equiv 1 + 2^{\alpha-2} \pmod{2^\alpha}$  ;
- (ii)  $k = l \equiv 1 - 2^{\alpha-2} \pmod{2^\alpha}$  .

**Theorem 3.2**

Let  $C_{2^\alpha} = \langle x \rangle$  and  $C_{2^\beta} = \langle y \rangle$  be cyclic groups of 2-power order. Furthermore, let  $C_{2^\alpha}$  and  $C_{2^\beta}$  act on each other so that  ${}^y x = x^k$  and  ${}^x y = y^l$  for  $k, l \in \mathbb{N}$  with  $(k, 2) = (l, 2) = 1$ . Then  $C_{2^\alpha}$  and  $C_{2^\beta}$  act compatibly on each other when  $\alpha \neq \beta$  with  ${}^{y^4} x = x$  and  ${}^{x^4} y = y$ , if and only if  $k$  and  $l$  are congruent to one of the following:

- (i)  $k \equiv 1 + 2^{\alpha-2} \pmod{2^\alpha}$  and  $l \equiv 1 + 2^{\beta-2} \pmod{2^\beta}$  ;
- (ii)  $k \equiv 1 + 2^{\alpha-2} \pmod{2^\alpha}$  and  $l \equiv 1 - 2^{\beta-2} \pmod{2^\beta}$  ;
- (iii)  $k \equiv 1 - 2^{\alpha-2} \pmod{2^\alpha}$  and  $l \equiv 1 + 2^{\beta-2} \pmod{2^\beta}$  ;
- (iv)  $k \equiv 1 - 2^{\alpha-2} \pmod{2^\alpha}$  and  $l \equiv 1 - 2^{\beta-2} \pmod{2^\beta}$  .

Next, let the actions of  $G$  and  $H$  be on each other such that  ${}^y x = x^k$  and  ${}^x y = y^l$  for  $k, l \in \mathbb{Z}$ . Now consider that one of the actions has order four and the other one has order two. The compatible conditions that are satisfied by  $k$  and  $l$  are given in **Table 3** when  $G = H$  and in **Table 4** when  $G \neq H$ .

**TABLE 3.** Values of  $k$  for which the action has order four and  $l$  for which the action had order two satisfying the compatible conditions when the groups  $G$  and  $H$  are the same.

Groups	$k$	$l$	Groups	$k$	$l$	Groups	$k$	$l$
$G = H = C_{2^4}$	3	9	$G = H = C_{2^4}$	3	9	$G = H = C_{2^4}$	3	9
	5	9		5	9		5	9
	11	9		11	9		11	9
	13	9		13	9		13	9
$G = H = C_{2^5}$	7	17	$G = H = C_{2^5}$	7	17	$G = H = C_{2^5}$	7	17
	9	17		9	17		9	17
	23	17		23	17		23	17
	25	17		25	17		25	17

**TABLE 4.** Values of  $k$  for which the action has order four and  $l$  for which the action has order two satisfying the compatible conditions when the groups  $G$  and  $H$  are different.

Groups	$k$	$l$	Groups	$k$	$l$	Groups	$k$	$l$
$G = C_{2^4}$	3	17	$G = C_{2^5}$	7	9	$G = C_{2^6}$	15	9
and	5	17	and	9	9	and	17	9
$H = C_{2^5}$	11	17	$H = C_{2^4}$	23	9	$H = C_{2^4}$	47	9
$G = C_{2^4}$	13	17	$G = C_{2^5}$	25	9	$G = C_{2^6}$	49	9
$G = C_{2^4}$	3	33	$G = C_{2^5}$	7	33	$G = C_{2^6}$	15	17
and	5	33	and	9	33	and	17	17
$H = C_{2^6}$	11	33	$H = C_{2^6}$	23	33	$H = C_{2^5}$	47	17
$G = C_{2^4}$	13	33	$G = C_{2^5}$	25	33	$G = C_{2^6}$	49	17
$G = C_{2^4}$	3	65	$G = C_{2^5}$	7	65	$G = C_{2^6}$	15	65
and	5	65	and	9	65	and	17	65
$H = C_{2^7}$	11	65	$H = C_{2^7}$	23	65	$H = C_{2^7}$	47	65
$G = C_{2^4}$	13	65	$G = C_{2^5}$	25	65	$G = C_{2^6}$	49	65
$G = C_{2^4}$	3	129	$G = C_{2^5}$	7	129	$G = C_{2^6}$	15	129
and	5	129	and	9	129	and	17	129
$H = C_{2^8}$	11	129	$H = C_{2^8}$	23	129	$H = C_{2^8}$	47	129
$G = C_{2^4}$	13	129	$G = C_{2^5}$	25	129	$G = C_{2^6}$	49	129
$G = C_{2^4}$	3	257	$G = C_{2^5}$	7	257	$G = C_{2^6}$	15	257
and	5	257	and	9	257	and	17	257
$H = C_{2^9}$	11	257	$H = C_{2^9}$	23	257	$H = C_{2^9}$	47	257
$G = C_{2^4}$	13	257	$G = C_{2^5}$	25	257	$G = C_{2^6}$	49	257
$G = C_{2^4}$	31	9	$G = C_{2^8}$	63	9	$G = C_{2^9}$	127	9
and	33	9	and	65	9	and	129	9
$H = C_{2^4}$	95	9	$H = C_{2^4}$	191	9	$H = C_{2^4}$	383	9
$G = C_{2^7}$	97	9	$G = C_{2^8}$	193	9	$G = C_{2^9}$	385	9
$G = C_{2^7}$	31	17	$G = C_{2^8}$	63	17	$G = C_{2^9}$	127	17
and	33	17	and	65	17	and	129	17
$H = C_{2^5}$	95	17	$H = C_{2^5}$	191	17	$H = C_{2^5}$	383	17
$G = C_{2^7}$	97	17	$G = C_{2^8}$	193	17	$G = C_{2^9}$	385	17
$G = C_{2^7}$	31	33	$G = C_{2^8}$	63	33	$G = C_{2^9}$	127	33
and	33	33	and	65	33	and	129	33
$H = C_{2^6}$	95	33	$H = C_{2^6}$	191	33	$H = C_{2^6}$	383	33
$G = C_{2^7}$	97	33	$G = C_{2^8}$	193	33	$G = C_{2^9}$	385	33
$G = C_{2^7}$	31	129	$G = C_{2^8}$	63	65	$G = C_{2^9}$	127	65
and	33	129	and	65	65	and	129	65
$H = C_{2^8}$	95	129	$H = C_{2^7}$	191	65	$H = C_{2^7}$	383	65
$G = C_{2^7}$	97	129	$G = C_{2^8}$	193	65	$G = C_{2^9}$	385	65
$G = C_{2^7}$	31	257	$G = C_{2^8}$	63	257	$G = C_{2^9}$	127	129
and	33	257	and	65	257	and	129	129
$H = C_{2^9}$	95	257	$H = C_{2^9}$	191	257	$H = C_{2^8}$	383	129
$G = C_{2^7}$	97	257	$G = C_{2^8}$	193	257	$H = C_{2^8}$	385	129

The result is given in the following theorem.

**Theorem 3.3**

Let  $G = \langle x \rangle \cong C_{2^\alpha}$  and  $H = \langle y \rangle \cong C_{2^\beta}$  be cyclic groups of 2-power order. Furthermore, let  $G$  and  $H$  act on each other so that  ${}^y x = x^k$  and  ${}^x y = y^l$  for  $k, l \in \mathbb{N}$  with  $(k, 2) = (l, 2) = 1$ . Then  $G$  and  $H$  act compatibly on each other with the action of  $H$  on  $G$  is order 4 and the action of  $H$  on  $G$  of order 2, if and only if  $k$  and  $l$  are congruent to one of the following:

- (i)  $k \equiv 1 + 2^{\alpha-2} \pmod{2^\alpha}$  and  $l \equiv 1 - 2^{\beta-1} \pmod{2^\beta}$  ;
- (ii)  $k \equiv 1 - 2^{\alpha-2} \pmod{2^\alpha}$  and  $l \equiv 1 - 2^{\beta-1} \pmod{2^\beta}$  ;
- (iii)  $k \equiv -1 + 2^{\alpha-2} \pmod{2^\alpha}$  and  $l \equiv 1 - 2^{\beta-1} \pmod{2^\beta}$  ;
- (iv)  $k \equiv -1 - 2^{\alpha-2} \pmod{2^\alpha}$  and  $l \equiv 1 - 2^{\beta-1} \pmod{2^\beta}$  .

## CONCLUSION

In this paper, three theorems have been stated accordingly to the order of the actions that satisfy compatible actions, which covered when the both actions have order 4, and when one of the actions has order 4 while the other has order 2

## ACKNOWLEDGMENTS

The authors would like to thank the Research & Innovation Department, Universiti Malaysia Pahang for the financial funding through RACE Grant, RDU141307.

## REFERENCES

1. R. Brown, J.-L. Loday, "Excision homotopique en basse dimension," *C.R. Acad. Sci. Ser. I Math. Paris*, vol. **298**, pp. 353-356, 1984.
2. R. Brown, D.L. Johnson and E.F. Robertson, "Some Computations of Non-Abelian Tensor Products of Groups," *Journal of Algebra*, vol. **111**, pp. 177-202, 1987.
3. L-C. Kappe, "Nonabelian Tensor Products of Groups: The Commutator Connection," *Proceeding Groups St. Andrews 1997 at Bath*, LMS Lecture Notes.
4. G. Ellis and A. McDermott, "Tensor products of prime-power groups," *Journal of Pure and Applied Algebra*, vol. **132**, pp. 119-128, 1998.
5. M. Visscher, "On the Nonabelian Tensor Products of Groups," Dissertation, State University of New York at Binghamton, 1998.
6. <http://www.gap-system.org>. 2005. The GAP Group, GAP – Groups, Algorithms, and Programming, Version 4.4.