Paddy Disease Detection System Using Image Processing

Radhiah Binti Zainon

THESIS SUBMITTED IN FULFILMENT OF THE DEGREE OF COMPUTER SCIENCE

FACULTY OF COMPUTER SYSTEM AND SOFTWARE ENGINEERING

2012

SUPERVISOR DECLARATION

"I hereby declare that I have checked this project and in my opinion this project is satisfactory in term of scope and quality for the award of degree of Bachelor Computer Science (Software Engineering)".

Supervisor Signature:Supervisor Name: Dr. Lee Ho CheongDate: 13/12/2012

DECLARATION

I hereby declare that the work in this thesis is my own except for quotations and summaries which have been duly acknowledged.

Signature :

Name of Candidate: Radhiah Binti ZainonDate: 13/12/2012

ACKNOWLEDGMENTS

First and foremost praise is to Almighty Allah for all his blessings for giving me patience and good health throughout the duration of this research.

I am deeply indebted to all the people who, directly and indirectly, are responsible for this project to become success. I am most grateful to Dr. Lee Ho Cheong as my supervisor for his encouragement, positive criticism of my research in this project, and most all his guidance and support for the whole project.

I am also thankful to Dr. Mohammad Masroor Ahmed for his guidance and helping developing the system.

And finally, a lot of thanks to my parents, family and friends who endured this long process together with me within guidance and full supportive.

ABSTRACT

The main objectives of this research is to develop a prototype system for detect the paddy disease which are Paddy Blast Disease, Brown Spot Disease, Narrow Brown Spot Disease. This paper concentrate on the image processing techniques used to enhance the quality of the image and neural network technique to classify the paddy disease. The methodology involves image acquisition, pre-processing and segmentation, analysis and classification of the paddy disease. All the paddy sample will be passing through the RGB calculation before it proceed to the binary conversion. If the sample is in the range of normal paddy RGB, then it is automatically classify as type 4 which is Normal. Then, all the segmented paddy disease sample will be convert into the binary data in excel file before proceed through the neural network for training and testing. Consequently, by employing the neural network technique, the paddy diseases are recognized about 92.5 percent accuracy rates. This prototype has a very great potential to be further improved in the future.

ABSTRAK

Objektif utama kajian ini adalah untuk membangunkan satu sistem prototaip bagi mengesan penyakit padi iaitu Penyakit Karah Padi, Penyakit Spot Coklat, Penyakit Spot Sempit Coklat . Karya ini menumpukan perhatian kepada teknik-teknik pemprosesan imej yang digunakan untuk meningkatkan kualiti imej dan teknik rangkaian neural untuk mengklasifikasikan penyakit padi. Metodologi melibatkan pemerolehan imej, pra-pemprosesan dan segmentasi, analisis dan klasifikasi penyakit padi. Semua sampel padi akan melalui pengiraan RGB sebelum ia meneruskan untuk penukaran binari. Jika sampel adalah dalam julat RGB padi normal, maka ia secara automatik mengklasifikasikan sebagai jenis 4 iaitu Normal. Kemudian, semua sampel penyakit padi yang telah disegmentasi akan menukar ke data binari dalam bentuk fail excel sebelum meneruskan melalui rangkaian neural untuk latihan dan ujian. Oleh itu, dengan menggunakan teknik rangkaian neural, kadar ketepatan penyakit padi diiktiraf kira-kira 92,5 peratus. Prototaip ini mempunyai potensi yang sangat besar untuk terus meningkat pada masa hadapan.

LIST OF TABLE

Table No	Title	Page
2.1	Paddy Disease, Symptom and Management	10
3.1	Software requirement	26
3.2	Hardware requirement	27
5.1	Testing result	63
5.2	RGB calculation of Sample 1	104
5.3	RGB calculation of Sample 2	104
5.4	RGB calculation of Sample 3	104
5.5	RGB calculation of Sample 4	105
5.6	RGB calculation of Sample 5	105
5.7	RGB calculation of Sample 6	105
5.8	RGB calculation of Sample 7	106
5.9	RGB calculation of Sample 8	106
5.10	RGB calculation of Sample 9	106
5.11	RGB calculation of Sample 10	107

LIST OF FIGURE

Figure No Title		Page
2.1	Blast Disease on leaf	7
2.2	Paddy Brown Spot Disease	9
2.3	Narrow Brown spot Disease	9
2.4	The methodology	12
2.5	(a) Blast Disease	13
	(b) Brown Spot Disease	
	(c) Narrow Brown Spot Disease	
2.6	(a) RGB image	13
	(b) Binary image with noise	
	(c) Binary image with noise free	
2.7	Example of grape leaf color extraction	16
2.8Example of grape leaf disease segmentation		16
	3.1 Iterative and incremental development method	
3.2	Flow chart of Paddy Disease Detection System	23 25
3.3	(a) Blast Disease	
	(b) Brown Spot Disease	28-29
	(c) Narrow Brown Spot Disease	
3.4	(a) RGB Image	
	(b) Binary Image with Noise	30-31
	(c) Binary Image with Noise free	
3.5	(a) Sample of Paddy Blast	
5.5	(b) Sample of brown Spot Disease	32-33
	(c) Sample of Narrow Brown Spot Disease	52-33
	(c) Sample of Martow Brown Spot Disease	
3.6	Prototype of paddy disease detection system	34

4.1	Image Acquisition source code	38
4.2	Grayscale source code	39
4.3	Filtering source code	40
4.4	Threshold source code	41
4.5	(a) Calculation of range RGB for Normal leaf	41-42
	(b) Binary image is converted into excel files	
4.6	Neural Network source code	43-45
5.1	Original image	47
5.2	5.2 Grayscale image	
5.3	5.3 Filtering image	
5.4 Binary image		49
5.5 Training the neural network		50
5.6	Neural network performance plot	50
5.7 Neural network training state plot		51
5.8	5.8Neural network regression plot51	
5.9	5.9GUI of Paddy Disease Detection System52	
5.10	Gantt Chart	60
5.11 Plagiarism checker		61

LIST OF APPENDIX

Appendix	Title	Page
А	Gantt chart	59
В	Plagiarism	61
C Testing result		62
D Calculation of RGB for Normal paddy 103		103

TABLE OF CONTENT

CHAPTER	TITLE			PAGE	
	Supe	rvisor Declaration		i	
	Stud	ent Declaration		ii	
	Ackr	owledgement		iii	
	Abst	ract		iv	
	Abst	rak		V	
	List	of Table		vi	
	List	of Figure		vii	
	List	of Abbreviation		viii	
1	INTI	RODUCTION			
		Introduction of Stud	dy	1	
	1.1	Problem Backgrour	nd	2	
	1.2	Problem Statement		3	
	1.3	Objectives		3	
	1.4	Scope		3	
	1.5	Thesis Organization	1	4	
2	LITI	CRATURE REVIEW	7		
	2.1	Paddy Overviews		6	
		2.1.1 Definition o	f Paddy	6	
		2.1.2 Paddy Disea	ase, Symptom and Management	6	
	2.2	Case Study on Exis	ting System	11	
		2.2.1 Investigatin	g on Image Processing Technique		
		for Diagnos	ing Paddy Disease	11	
		2.2.1.1 Meth	nod	11	
		А	Image Acquisition	12	
		В	Image Segmentation and Pre		
			Processing	13	

		С	Feature Extraction using Texture	
			Analysis	14
		D	Image Classification	14
	2.2.2	Grape leaf I	Disease Detection from Color	
		Imagery usi	ng Hybrid Intelligent System	15
		2.2.2.1 Met	hod	15
		А	Grape leaf color segmentation	15
		В	Grape leaf disease segmentation	16
		С	Analysis and Classification of	
			Disease	17
	2.2.3	A Framewo	rk for Detection and Classification	
		of Plant Lea	ıf	19
		2.2.3.1 Met	hod	19
		А	Clustering Method	19
		В	Feature Extraction	20
2.3	Propo	se System		21
2.4	Conclu	usion		21

3 METHODOLOGY

3.1	Introduction 2			
3.2	The Ju	The Justification of Iterative and Incremental		
	Devel	opment	23	
3.3	Iterati	ve and incremental Development Method	23	
	3.3.1	Planning Stage	24	
	3.3.2	Software and Hardware Requirement/		
		Specification	26	
		3.3.2.1 Software Requirement	26	
		3.3.2.2 Hardware Requirement	27	
	3.3.3	Analysis and Design Stage	28	
	5.5.5	Anarysis and Design Stage	20	
		3.3.3.1 Formulation	28	
		3.3.3.2 Interface Design	34	

3.3.4	Implementation Stage	35
3.3.5	Testing Stage	35
3.3.6	Deployment and Maintenance	36

4 IMPLEMENTATION

4.1	Introduction	38
4.2	Image Acquisition	
4.3	Image Segmentation and Pre Processing	39
	4.2.1 Grayscale	40
	4.2.2 Filtering	41
	4.2.3 Threshold	41
4.4	Analysis and Classification of Paddy Disease	42

5 **RESULT, DISCUSSION AND CONCLUSION**

5.1	Introduction	47	
5.2	Image Acquisition	48	
5.3	Image Segmentation and Pre Processing	48	
	4.2.1 Grayscale	48	
	4.2.2 Filtering	49	
	4.2.3 Threshold	49	
5.4	Analysis and Classification of Paddy Disease	50	
5.5	Interface of the system		
5.6	Advantages and Disadvantages	53	
	5.6.1 Advantages	54	
	5.6.2 Disadvantages	54	
5.7	Achieve objectives	54	
5.8	Constraints	55	
5.9	Assumptions and Further Research 53		

5.10 Conclusion	56
REFERENCES	57
APPENDIX	60

CHAPTER I

INTRODUCTION

This chapter discuss about an overview of the study conducted. The title of my project is Paddy Disease Detection System Using Image Processing .It consists of background, problem statements, objectives and the scope of study. The background briefly describes the identification of projects and related issues. Problem statements describes the problems that arise and make the selected projects to be undertaken. The objectives are the goals list for the projects to be achieves. Scope of study discuss about the limitations of projects and users. Lastly, thesis organization gives a summary of the sequence for each chapter in the thesis.

1.1 Background

A product quality control is fundamentally required in order to gain more valueadded products[2]. Many studies show that quality of agricultural products can be reduced from many causes. One of the most important factors of such quality is plant diseases. Consequently, minimizing plant diseases allows substantially improving quality of the products.

Rice known as Oryza Sativa (specific name), is one of the most utilized food plants and widely grown originated in ASIA. [4] Rice is an important crop worldwide and over half of the world population relies on it for food. Many people in the world including Malaysia eat rice as staple food. However, there are many factors that make paddy rice production become slow and less productive. One of the main factors is paddy disease.

An abnormal condition that injures the plant or leads it to function improperly is called as a disease. Diseases are readily recognized by their symptoms. There are a lot of paddy disease types which are Bakanae, red disease virus, brown spot disease and many more.[1] Image processing and computer vision technology are very beneficial to the agricultural industry. They are more potential and more important to many areas in agricultural technology [1].

Paddy Disease Detection System is one of the very beneficial systems. It can help the paddy farmer detect the disease faster. This study aims to develop a prototype system to automatically detect and classify the paddy diseases by using image processing technique as an alternative or supplemental to the traditional manual method.

1.2 Problem Statement

Paddy will be harvest twice in a year. Most of paddy farmer faces many problems to harvest their paddy because they had been attack by snail, worm and fungi. Furthermore, when the paddy had been infected or attacked, the others areas had been exposed to be infected. Thus, it will decrease paddy farmer's income and lead to significance losses to farmer. Currently, the paddy farmer determines the type of disease manually. The errors might occur in order to determine the type of diseases. Paddy farmer also have to spend a lot of time to detect the type of disease. It also takes a time as the paddy farmers manually check the disease since the paddy field is in wide area.

1.3 Objective

There are three objectives to achieve in this project:

i. to develop the prototype of paddy disease detection system

ii. to detect the paddy disease by using image processing

iii. to apply image processing technique to analyze the pattern of paddy disease

1.4 Scope

- The users of this system are paddy farmers.
- The prototype will be develop by using MATLAB 7.10
- 10 samples each of the normal, brown spot disease, narrow brown spot disease and blast disease will be used in this project.

1.5 Thesis Organization

This thesis consists of 6 chapters ranging from Chapter 1 until Chapter 6. Chapter 1 gives an overview of the study conducted. It also consists of problem statements, objectives and the scope of study. Meanwhile, Chapter 2 reviews the previous research works that was conducted by others researches. All the relevant technical paper, journals and books taken from those researches will be discussed in detail. Chapter 3 focuses about the methodology for system development and process flow in detail of this research. It reveals the technique and the algorithms that will be used in performing this study. Chapter 4 concentrates on details of the implementation of the system which consists of coding phase and their explanation of the coding function. Results of the testing are to be expounding in Chapter 5 that consist of expected result or output, constraint of project and further research. Lastly, Chapter 6 concludes the overall study and research.

CHAPTER II

LITERATURE REVIEWS

This chapter briefly reviews, explains and discusses on existing literature review related with the current project which is "Paddy Disease Detection System Using Image Processing" that will be developed later. This chapter comprises three sections. The first section describes the overviews of paddy. The subsections are the definition, type of paddy disease, paddy symptom and paddy management. The second section is the review of some existing system that used same techniques and methods. The third section discusses the review on technique and method used by the system. The subsections are image acquisition, image segmentation and artificial neural network.

2.1 Paddy Overviews

In this section, firstly presents a definition of paddy. After that, this subsection briefly discusses on type of paddy disease, symptoms and management of paddy disease.

2.1.1 Definition of Paddy

Paddy also known as rice is the starchy seeds of an annual southeast Asian cereal grass (*Oryza sativa*) that are cooked and used for food. This cereal grass that is widely cultivated in warm climates for its seeds and by-products [8]. Rice is one of the most utilized food plants and widely grown originated in ASIA. Rice is an important crop worldwide and over half of the world population relies on it for food. Many people in the world including Malaysia eat rice as staple food.

2.1.2 Paddy Diseases, Symptoms and Management

There are many factors that make paddy rice production become slow and less productive. One of the main factors is paddy disease. The table below will show you type of paddy disease, the symptom of paddy disease and the management of paddy disease. This researches focus on three types of diseases, which are paddy blast, brown spot disease and narrow brown spot disease. The following paddy diseases are shown below:

No	Disease	Disease Symptoms
1	Paddy Blast	Paddy Blast Symptoms
	Fig. 2.1 : Blast disease on leaf	 Disease can infect paddy at all growth stages and all aerial parts of plant (Leaf, neck and node). Among the three leaves and neck infections are more severe. Small specks originate on leaves - subsequently enlarge into spindle shaped spots(0.5 to 1.5cm length, 0.3 to 0.5cm width) with ashy center. Several spots coalesce -> big irregular patches
		Management
		 Avoid excess N - fertilizer application Apply nitrogen in three split doses. Removes weed hosts from bunds. Use of tolerant varieties (Penna, Pinakini, Tikkana, Sreeranga, Simphapuri, Palghuna, Swarnamukhi, Swathi, Prabhat, Co 47, IR - 64, , IR - 36, Jaya) Burning of straw and stubbles after harvest Dry seed treatment with <i>Pseudomonas</i> <i>fluorescens</i>talc formulation @10g/kg of seed. Stagnate water to a depth of 2.5cm over an area of 25m2 in the nursery. Sprinkle 2.5 kg of <i>P</i>. <i>fluorescens</i> (talc) and mix with stagnated water. Soak the root system of seedlings for 30 min and transplant. Spray <i>P. fluorescens</i> talc formulation @ 0.5% from 45 days after transplanting at 10 day intervals, three times.

2	Paddy Brown Disease	Spot	 Edifenphos @ 0.1 %. Pre-Tillering to Mid-Tillering Light at 2 to 5 % disease severities - Apply Edifenphos or Carbendazim @ 0.1 %. Delay top dressing of N fertilizers when infection is seen. Panicle initiation to booting At 2 to 5% leaf area damage spray Edifenphos or Carbendazim or Tricyclazole @ 0.1 %. Flowering and after At 5 % leaf area damage or 1 to 2 % neck infection spray Edifenphos or Carbendazim or Tricyclazole @ 1 g /lit of water. Paddy Brown Spot Disease
	Disease		 Initial lesions are water-soaked to greenish gray and later become grayish white with brown margin Lesions on leaf sheaths near waterline Presence of sclerotia

		Partially filled or empty grains
	0 80	Management
	Fig 2.2: Paddy Brown Spot Disease	 Apply FYM 12.5 t/ha or green manure 6.25 t/ha to promote antagonistic microflora. Soil application of <i>P. fluorescens</i> @ 2.5 kg/ha mixed with 50 kg FYM after 30 days of transplanting. Foliar spraying of <i>P. fluorescens</i> @0.2% at boot leaf stage and 10 days later. Avoid flow of irrigation water from infected to healthy field. Carbendazim (1 g/lit), Propiconazole (1ml/lit) may be applied. Spraying of infected plants with fungicides, such as Benomyl and Iprodione, and antibiotics, such as Validamycin and Polyoxin, is effective against the disease Reduce Nitrogen dosage and skip top dressing
3	Narrow Brown Spot	Symptoms
	Disease Fig 2.3 : Narrow Brown Spot Disease	 Short, narrow, elliptical to linear brown lesions usually on leaf blades but may also occur on leaf sheaths, pedicels, and glumes or rice hulls Lesions about 2-10 mm long and 1 mm wide Lesions narrower, shorter, and darker brown on resistant varieties Lesions wider and lighter brown with gray necrotic centers on susceptible varieties Leaf necrosis may also occur on susceptible varieties Lesions occur in large numbers during the later growth stages

Why Occurs?
 The disease is observed on rice crops grown on soil deficient in potassium. Temperature ranging from 25-28° C is favorable for the optimum growth of the disease. Susceptibility of the variety to the fungus and the growth stage of the rice crop are other factors that affect the development of the disease. Although rice plants are susceptible to the fungus at all stages of growth, they are more susceptible from panicle emergence to maturity, thus, becoming more severe as rice approaches maturity.
Management
 Cultural practices, such as the use of potassium and phosphorus fertilizers, and planting of early maturing cultivars early in the growing season, are recommended to manage the narrow brown leaf spot. The use of resistant varieties is the most effective approach to manage the disease. However, the resistant varieties and lines are only grown in United States and India. Spraying of fungicides such benomyl, propicanazole, carbendazim, propiconazole, and iprodione, when the disease is observed in the field is effective.

2.2.1 Investigating on Image Processing Techniques for Diagnosing Paddy Diseases

A study conducted by Nunik Noviana Kurniawati, Siti Norul Huda Sheikh Abdullah, Salwani Abdullah, Saad Abdullah from Universiti Kebangsaan Malaysia aims to develop a prototype system to automatically and correctly detect and classify the paddy diseases with Blast Disease (BD), Brown Spot Disease (BSD), and Narrow Brown Spot Disease (NBSD) using image processing technique as an alternative or supplemental to the traditional manual method.[1]

2.2.1.1 Method

The methodology for diagnosing paddy diseases can be simplified as Fig. 2.4. This process involves several tasks, such as image acquisition and collection, image segmentation and pre-processing, shape feature extraction and color feature extraction, and paddy diseases classification based on lesion type, boundary color, spot color, and broken paddy leaf color. Cunha [18] used recognition technique to analyze the pathological stress conditions and characterization of the fruits or plant leafs. Runtz and Dave [19,20] applied image processing technique for classification and identifying of the plant species.




Figure 2.4: The methodology

A Image Acquisition

The RGB colour images of paddy leaf are captured using a Canon PowerShot G2 digital camera, with pixel resolution 768x1024. The digitized images are about 225 KB size each. Those images are cropped into a smaller image with dimension of 109 x 310 pixels. There have collected about 94 data samples. It consists of three types of paddy diseases as shown in Fig. 2.5. Images are stored in BMP format. The prototype uses Matlab image processing library.




Fig 2.5 : (a) Blast Disease ; (b) Brown Spot Disease; (c) Narrow Brown Spot Disease

B Image Segmentation and Pre-Processing

The main objective of this process is to obtain the binary image with less noise or noise free. In order to achieve high accuracy, an appropriate silhouette should be obtained. The RGB image (Fig. 2.6(a)) is converted into a binary image using threshold method, as shown in Fig. 2.6(b). They used local entropy threshold methods of Eliza and Chang [11] and Otsu method. A morphological algorithm is used to remove noises by using region filling technique.




Fig 2.6 : (a) RGB image ; (b) Binary image with noise; (c) Binary image with noise free

C Feature Extraction using Texture Analysis

The image analysis focused on the shape feature extraction and color based segmentation.

D Image Classification

The production rules have been developed through serial interviews with agricultural expert based on above characteristics, such as lesion type, boundary color, spot color, and broken paddy leaf color, paddy diseases is recognized using production rule method with forward-chaining method.

2.2.2 Grape Leaf Disease Detection from Color Imagery using Hybrid Intelligent System

A study conducted by A. Meunkaewjinda, P. Kumasawat, K. Attakitmongcol and A. Srikew on Grape Leaf Disease Detection from Color Imagery using Hybrid Intelligent System [2]. Vegetable and fruits are the most important export agricultural products of Thailand. In order to obtain more value added products, a product quality control is essentially required. Many studies show that quality of agricultural products may be reduced from many causes. One of the most important factors of such quality is plant disease. Consequently, minimizing plant diseases allows substantially improving quality of the products. This system shows an automatic plant disease diagnosis using multiple artificial intelligent techniques. The system can analyze or diagnosis plant leaf disease without maintain any expertise once the system is trained.

2.2.2.1 Method

There are several steps must be achieve of this system which are grape leaf color segmentations, grape leaf disease segmentation and analysis and classification of the disease.

A Grape leaf color segmentation

The grape leaf color segmentation is pre-processing module which segments out of any irrelevant background information. A self-organizing features map together with a back-propagation neural network is deployed to recognize colors of grape leaf. This information is used to segment grape leaf pixels within the image. The input is enhanced by using anisotropic diffusion technique [12] to preserve information of the affected pixels before extracting grape color from the background.





(a) Original image(b) Grape leaf with color extractionFig. 2.7 : Example of grape leaf color extraction

B Grape leaf disease segmentation


The grape leaf disease segmentation is performed using modified selforganizing features map with genetic algorithms for optimization and support vector machines for classification. Using modified self-organizing feature map (MSOFM) [13], the clustering process does not require any training nor predefined number of color groups.



(a) Original image




(b)Grape leaf disease color extraction




C Analysis and Classification of Disease

The resulting segmented image is filtered by Gabor wavelet [14] which allows the system to analyze leaf disease color features more efficient. The support vector machines are then again applied to classify types of grape leaf diseases. The image can be able to categorize the image of grape leaf into three classes:

- i) Scab disease
- ii) Rust disease
- iii) No disease



(a) Example of rust disease detection



(b) Example of scab disease detection

2.2.3 A Framework for Detection and Classification of Plant Leaf and Stem Disease

A study conducted by Dheeb Al Bashish, Malik Braik, and Sulieman Bani-Ahmad on a framework for Detection and Classification of Plant Leaf and Stem Diseases. Studies show that relying on pure naked-eye [15] observation of experts to detect such diseases can be prohibitively expensive, especially in developing countries. Providing fast, automatic, cheap and accurate image processing-based solutions for that task can be of great realistic significance.

2.2.3.1 Method

The proposed framework is image processing-based and is composed of the following main steps;

a) The images at hand are segmented using the K-Means technique

b) The segmented images are passed through a pre-trained neural network.

A Clustering method

K-means clustering is used to partition the leaf image into four clusters in which one or more clusters contain the disease in case when the leaf is infected by more than one disease.The k-means clustering algorithms tries to classify *objects* (pixels in our case) based on a set of features into K number of classes. The classification is done by minimizing the *sum of squares* of distances between the objects and the corresponding cluster or class *centroid*. On this experiment, the K-means clustering is set to use squared Euclidean distances.

B Feature Extraction

The method followed for extracting the feature set is called the *Color Co-occurrence Method* or CCM method in short. It is a method, in which both the color and texture of an image are taken into account, to arrive at unique features, which represent that image.

- Co-occurrence Methodology for Texture Analysis
- Normalizing the CCM matrices
- Texture features identification

As a testbed, D. Al Bashish (2001) use a set of leaf images taken from Al-Ghor area in Jordan. There are five diseases which effect on the plants;

- a) Cottony mold
- b) Early scorch
- c) Ashen mold
- d) Late scorch
- e) Tiny whiteness.

The experimental results indicate that the proposed approach can significantly support accurate and automatic detection of leaf diseases. The developed Neural Network classifier that is based on statistical classification perform well and could successfully detect and classify the tested diseases with a precision of around 93%.

2.3 Propose System

There is a growing demand of image processing in diverse application areas, such as multimedia computing, secured image data communication, biomedical imaging, biometrics, remote sensing, texture understanding, pattern recognition, content-based image retrieval, compression and so on. To improve the quality of paddy, there must be a system that can accurately detect the disease so that the paddy farmer can cure it as soon as possible. The methodology consists of the pre-processing and segmentation of paddy disease and classification of the disease. The techniques that will be applied for the classification of paddy disease in this system are feed-forward neural network technique.

2.4 Conclusion

Based on the previous system, there are a lot of methods and techniques can be applied to the system such as rule method with forward-chaining method, modified selforganizing feature (MSOFM), k-means and pre-trained neural network. The details of methodology and techniques will be discussed on the next chapter. **CHAPTER III**

METHODOLOGY

3.1 Introduction

The purpose of this chapter is to discuss the approach and framework for the project. Method, technique or approach that will be or has been used while designing and implementing the project included in the content. This chapter also explains about the justification of method or approach used and hardware and software necessity. This project will be conducted based on the Iterative and Incremental Method.

3.2 The justification of Iterative and Incremental Development Methodology

Typically agile software development combines iterative and incremental techniques to deliver functionality early and often [21]. Iterative means perform repeatedly. Iterative also can be refers as repeated application of a procedure, applying it each time to the result of the previous application. Besides, iterative development refactors code repeatedly, making progress through successive refinement. Increment means an increase or addition. Incremental development builds and delivers software to a production environment as a series of small and regular releases with expanding functionality. Agile software development employs both iterative and incremental development techniques but also applies feedback across releases or increments or iteration. Moreover, incremental development becomes evolutionary, where the previous iteration guides the next iteration. Iterative and Incremental development developed in response to the weakness of the waterfall model. It starts with an initial planning and ends with deployment with the cyclic interaction in between [22].

3.3 Iterative and Incremental Development Method

Paddy Disease Detection System contains some stage or phase of development method such as planning stage, requirement, analysis and design, implementation, testing, evaluation and maintenance.




Fig 3.1 : Iterative and incremental development method

3.3.1 Planning Stage

The first stage to start the development of Paddy Disease Detection System using Image Processing is planning. Planning is the process of gathering all the information including problem statements, objectives and the scope of the system from the journal and article reading from the internet.




Fig 3.2: Flow chart of Paddy Disease Detection System

3.3.2 Software and Hardware Requirement/Specification

There are several things that should be considered to make sure the development stage of the system can run successfully. There are software and hardware specification. Software is a conceptual entity which is a set of computer programs, procedures, and associated documentation concerned with the operation of a data processing system. Hardware is component devices which are typically installed into or peripheral to a computer case to create a personal computer upon which system software is installed including a firmware interface.

3.3.2.1 Software Requirement

The software requirement for Paddy Disease Detection System development has been identified as shown in table below:-

	Software	Purpose	
Operating System	Microsoft Windows Vista	As the operating system	
	™ Ultimate service pack 1		
	Microsoft Office Word	For documentation	
	2010		
	Microsoft Office Project	Gantt chart	
	2010		
	Microsoft Office Visio	Flowchart	
Software	2010		
	Microsoft Power Point	For slide presentation	
	2010		
	Adobe Acrobat Reader X	For reading from the internet	
		resources	
	MATLAB 7.10	Developments tools	

Table 3.1 : Software requirement

3.3.2.2 Hardware Requirement

The hardware requirement for Paddy Disease Detection System development has been identified as shown in table below:-

	Component	Minimum Requirement	
Laptop	Processor	Intel® Core(TM)2 duo CPU T6500	
		@ 2.10 GHz	
	RAM	3 GB	
	CPU	32-bit operating system	
Camera	Sensor	12 megapixel	
	Zoom	4x optical zoom.	
	Display	2.7-inch	
	Card Slot	SD/SDHC	

Table 3.2 : Hardware requirement

3.3.3 Analysis and Design Stage

Analysis is the process determining the needs or conditions to meet for a new or altered the system. Design is the process of problem solving and planning for a software solution. It includes low-level component and algorithm implementation issues as well as architectural view. There is a growing demand of image processing in diverse application areas, such as multimedia computing, secured image data communication, biomedical imaging, biometrics, remote sensing, texture understanding, pattern recognition, content-based image retrieval, compression and so on.

3.3.3.1 Formulation

A. Image Acquisition

In this process, it is a preparation process to obtain paddy leaf images. The RGB color images of paddy leaf are captured using Olympus *fe*-4050 digital camera with pixel resolution 2048x1024. Those image are cropped into smaller image. Images are stored in BMP format. The prototype uses Matlab image processing library.



Figure 3.3(a): Blast Disease



Figure 3.3(b): Brown Spot Disease



Figure 3.3(c): Narrow Brown Spot Disease

B. Image Segmentation and Image Pre-Processing

i. Converting RGB to gray scale image

The segmentation and pre-processing task are the initial stage before the image is used for the next process. The main objective of this process is to obtain the binary image with Otsu method. The Otsu method is based on selecting the lowest point between two classes of the histogram by considering the between-class variance.



Figure 3.4 (a): RGB Image



Figure 3.4(b): Binary Image with Noise



Figure 3.4(c): Binary Image with Noise free

ii. Filtering

Averaging filter is implemented in this process. The average filter computes the mean (average) of the gray-scale values within a rectangular filter window surrounding each pixel. This has the effect of smoothing the image (eliminating noise). The filtered pixel will be calculated by:

$$r = (a1 + a2 + \dots + a9)/9 \tag{1}$$

C. Feed Forward Neural Network

The neural network is applied to pattern recognition, system identification and system control. The feed forward neural network was the first and arguably simplest type of artificial neural network devised. In this network, the information moves in only one direction, forward, from the input nodes, through the hidden nodes (if any) and to the output nodes. There are no cycles or loops in the network.

i. Training and testing of neural network

In this stage, there is training database, initialize neural network layer, train network. 10 samples of Blast Disease image, 10 samples of Brown Spot Disease image and 10 samples of Narrow-Brown Spot Disease image is used as training image.




Figure 3.5(a): Sample of Blast Disease




Figure 3.5(b): Sample of Brown Spot Disease




Figure 3.5(c): Sample of Narrow Brown Spot Disease

D. Image Classification

Based on above training and testing of neural network, the image will be classify weather it is paddy blast, brown spot disease and narrow brown spot disease with the feed forward neural network.

3.3.3.2 Interface Design

Paddy Disease Detection System				
Browse Process	Grayscale Image	Filtering	Threshold	Edge Detection
	Type of Disease and Management :		Type1: Blast Disease	
	e e e e e e e e e e e e e e e e e e e	Reset Exit		

Fig 3.6: Prototype of paddy disease detection system

3.3.4 Implementation Stage

A software implementation method is a systematically structured approach to effectively integrate a software [23]. In this stage, the paddy system must be tested to make sure it is free from error. Besides, the implementation of the system includes the achievement of the goals/objective of the system and also to make sure the system can run smoothly. The designer should know and understand all the navigation key, coding and design.

3.3.5 Testing Stage

Software testing is an investigation conducted to provide information about the quality of the product or service under test[24]. Testing stage is the process of executing a program or application with the intent of finding software bugs (errors or other defects). A primary purpose of testing is to detect software failures so that defects may be discovered and corrected. The scope of software testing often includes examination of code as well as execution of that code in various environments and conditions as well as examining the aspects of code There are some criteria of validating and verifying that a software program/application/product of paddy system:

- 1. meets the requirements that guided its design and development;
- 2. works as expected; and
- 3. can be implemented with the same characteristics.

3.3.6 Deployment and Maintenance

Deployments are all of the activities that make a software system available for use[25]. Maintenance is the modification of a system after delivery to correct faults, to improve performance or other attributes[26]. The paddy system should be maintained from time to time by the administrator, if there is any correction or addition to the system in order to keep the user satisfaction.[27]

CHAPTER IV

IMPLEMENTATION

4.1. Introduction

This chapter briefly discusses the implementation of the project. As the software progresses, the implementation of Paddy Disease Detection System using Image Processing will need to test the software for improvement and proper implementation.

4.2 Image Acquisition

Image acquisition is the first step in system development to get the sample or image. For Paddy Disease Detection System using Image Processing, some of the sample is already taken from the internet and some of the sample manually captured in Kedah. The sample is already has kept into the folder in computer. So, there is no process of image acquisition.

The software platform that is used to execute the system is MATLAB 2010. In this system, the image is loaded from the folder in computer named "Paddy" by the user. User can choose which image to be load.

When the user browse the image by clicking the *Browse Button*, the system start to search the image stored.

```
% browse the image into the system
[filename1,pathname1] =
uigetfile({'*.jpg';'*.gif';'*.bmp';'*.*'}); ('Please
pick the paddy sample');
global imageSelected;
global sameImage;
imageSelected=(strcat(pathname1,filename1));
sameImage=(strcat(pathname1,filename1));
axes(handles.originalaxes);
imshow(imageSelected);
guidata(hObject, handles);
```

4.3 **Image segmentation and pre-processing**

4.3.1 Grayscale

First, the image is converted to the grayscale image. The function of G = rgb2gray(Z) is used to convert into the grayscale image. Then, the intensity of image is adjusted by calling function that specifies lower (bottom 1% of all pixel values) and upper limits (top 1% of all pixel values) that are used for contrast stretching the grayscale image.

```
% convert to grayscale image
global imageSelected;
global sameImage;
% convert to grayscale image
Z= imread(sameImage);
G = rgb2gray(Z);
axes(handles.grayscaleaxes);
imshow(G);
```

Figure 4.2 Grayscale source code

Filtering is one of the processes in image enhancement. The function of filtering is to remove noise and to smooth out the image acquired. The average filtering is applied in this system. Filter h = fspecial('average'); is applied to the image by function F = imfilter(G, h, 'symmetric'). Symmetric means input array values outside the bounds of the array are computed by mirror-reflecting the array across the array border.


```
% filtering
h = fspecial('average');
F = imfilter(G,h,'symmetric');
axes(handles.filtering);
imshow (F);
```

Figure 4.3 Filtering source code

4.3.3 Threshold

Threshold image is consist only with binary pixels where an object pixel is given a value '1' (remarked by white color) while background pixel is given value of '0' (remarked by black color). The image of grayscale image with fltering is convert to binary image by function BW = im2bw(F, level) as shown in figure 4.4.

```
% threshold
BW = im2bw (F,graythresh(F));
BW = ~ BW;
new = imfill(BW,'holes');
axes(handles.threshold);
imshow(new);
```



4.4 Analysis and Classification of Paddy Disease

Before the image is convert into the binary data, the image will be pass through the RGB calculation. This is for detecting the normal paddy leaf by using the range of RGB calculation. Figure 4.5 (a) shows the calculation of RGB. After that, the resulting segmented image allows the system to analyze leaf disease features more efficient. The binary images are convert into the binary data excel files where the column is 1 and the row is 240. The format of the excel files is .csv.

```
img=imread(inputImage);
Red = img(:,:,1);
Green = img(:,:,2);
Blue = img(:,:,3);
if (Red(:) >=93 & Red(:) <=204 & Green(:) >= 142 &
Green(:) <=210 & Blue(:) >= 64 & Blue(:) <=155)
result= 'Type 4 :Normal';
```

Figure 4.5 (a) Calculation of range RGB for Normal leaf

```
Row1 = 1;
Column1 = 240;
new = imresize(new, [Row1 Column1]);
new = new'; %Take transpose.
% Get the current folder fullpath name
p = mfilename('fullpath');
% break the fill path of ur .m file into parts
[pathStr, excelFile, ext] = fileparts(p);
% Set the file path for the excel file that will be
create later
newExcelFile=(strcat(pathStr,'\ConvertedData\','csvDa
ta.csv'));
%output data from MatLab to a Text file with data 0
and 1.
dlmwrite( newExcelFile, new,'delimiter', ',');
```

Figure 4.5(b) Binary image is converted into excel files.


The outputs of the data are 0 and 1. The excel files are then again applied to neural network to classify types of paddy diseases. The neural network can be able to categorize the image of paddy disease into four classes:

- i) Paddy blasé disease
- ii) Narrow Brown Spot Disease
- iii) Brown Spot Disease
- iv) Normal paddy leaf

```
Sample = 20; %number of samples made
Sample Attribute = 240; % (That means Size: 60x4);
% Get the current folder fullpath name
 p = mfilename('fullpath');
% break the fill path of ur .m file into parts
 [pathStr, excelFile, ext] = fileparts(p);
imageClass=(strcat(pathStr,'\csvFile\','mask Monochro
me Combined.csv'));
imageTarget=(strcat(pathStr,'\csvFile\','mask Monochr
ome Target.csv'));
% importdata from folder TrainFile
alphabet = csvread(imageClass);
targets = csvread(imageTarget);
%% create neural network using feedforward
net = newff(alphabet,targets,25);
net1 = net;
net1.divideFcn = '';
%% Train the Network
[net1,tr] = train(net1,alphabet,targets);
numNoisy = 2; %10;
% SET TESTING PARAMETERS
noise range = 0:.05:.5;
max test = 100;
network1 = [];
%network2 = [];
```

```
% PERFORM THE TEST
for noiselevel = noise range
  errors1 = 0;
  errors2 = 0;
  for i=1:max test
     x = alphabet +
randn(Sample Attribute,Sample)*noiselevel;
    %% TEST NETWORK 1
    %simulate the network
    y = sim(net1, x);
    yy = compet(y);
    errors1 = errors1 + sum(sum(abs(yy-targets)))/2;
   end
  % AVERAGE ERRORS FOR 100 SETS OF 26 TARGET VECTORS.
  network1 = [network1 errors1/Sample/max test];
end
 %Types of Paddy
            ['Type1: Blast Disease.'...
Tvpe1 =
             'Avoid excess N - fertilizer application.
Apply nitrogen in three split doses.'...
             'Removes weed hosts from bunds. Use of
tolerant varieties. '...
             'Burning of straw and stubbles after
harvest.'];
Type2 =
            ['Type2: Narrow Brown Spot Disease '...
              'Cultural practices, such as the use of
potassium and phosphorus fertilizers, '...
              'and planting of early maturing cultivars
early in the growing season, '...
              'are recommended to manage the narrow brown
leaf spot.'...
              'The use of resistant varieties is the most
effective approach to manage the disease. '...
              'Spraying of fungicides such benomyl,
propicanazole, carbendazim, propiconazole, and iprodione';]
```

```
Туре3 =
            ['Type3: Brown Spot Disease'...
              'Apply FYM 12.5 t/ha or green manure 6.25
t/ha. '...
              'Soil application of P. fluorescens @ 2.5
kg/ha mixed with 50 kg FYM after 30 days of
transplanting.'...
              'Foliar spraying of P. fluorescens @0.2percent
at boot leaf stage and 10 days later.'
              'Avoid flow of irrigation water from infected
to healthy field.'];
Type4 = 'Type4: Normal';
%Testing
noisyJ = double(column1);
%plotchar60x40(noisyJ);
y = sim(net1, noisyJ);
y = compet(y);
answer = find(compet(y) == 1);
%plotchar60x40(alphabet(:,answer));
if answer == 1
    result=Type1;
elseif answer == 2
   result=Type2;
elseif answer == 3
    result=Type3;
else
    result=Type4;
end;
```



CHAPTER V

RESULT, DISCUSSION AND CONCLUSION

5.1 Introduction

There are 30 samples of paddy image used as sample data in the testing phase of this development. The paddy images samples had gone through the phases as discuss in the chapter 3 and 4. This chapter will briefly describe about the output result of each phases.

5.2 Image Acquisition

Some of the paddy images for testing are collected from the paddy field in Kedah, and some of them were collected from the internet. The samples of paddy were taken based on the 3 types of disease including the normal leaf. Figure 5.1 show the sample captured image for testing phase.



Fig. 5.1 : Original image

5.3 Image Segmentation and Pre-Processing

After the image is captured and cropped, it then will undergo to the next process. The sample will converted into gray scale image, filtered with average filter, and threshold.

5.3.1 RGB to Grayscale

The captured image which is in RGB format was converted to grayscale format as shown in figure 5.2. The output of grayscale image also display in the GUI.




Fig 5.2 : Grayscale image

5.3.2 Filtering

After the image was converted to grayscale format, it will then filter using average filter to remove noise as shown in figure 5.3. The noise of the image cannot been seen by naked eye.




Fig 5.3 : Filtering image

5.3.3 Threshold

After the filtering process completed, the image then will be converted into the binary image as shown in figure 5.4.



Fig 5.4 : Binary image

5.4 Analysis and classification of paddy disease

The next step is create a neural network that will learn to identify the paddy disease. The feed forward network is used in this part for training and testing. Below is a picture of MLP structure with two hidden layer [25 10] neurons and sigmoid, linear activation function. Now the network is ready to be trained. The samples are automatically divided into training, validation ad test sets. The training set is used to teach network. Training continues as long as the network continues improving on the validation set. The test set provides a completely independent measure of network accuracy.

Neural Network		
Layer	Layer	_
	w · /	Output
Algorithms		
Training: Levenberg-Marqu	ardt (trainIm)	
Performance: Mean Squared Erro		
Progress		
Epoch: 0	8 iterations	1000
Time:	0:02:03	
Performance: 1.18	4.19e-31	0.00
Gradient: 1.00	2.95e-15	1.00e-10
Mu: 0.00100	1.00e-08	1.00e+10
Validation Checks: 0	0	6
Plots		
Performance (plotperform)		
Training State (plottrainstate	e)	
Regression (plotregressio	un)	
(prostegressie		
Plot Interval:	1 ерс	ochs
)pening Performance Plot		
	<i>c</i>	

Figure 5.5 : Training the Neural Network




Figure 5.6 : Neural Network performance plot




Figure 5.7: Neural network training state plot




Figure 5.8: Neural network regression plot

5.5 Interface of the system




Figure 5.9 shows GUI that help the user to check the output of the prototype.

Figure 5.9: GUI of Paddy Disease Detection System

5.6 Advantages and disadvantages

The system has some advantages and disadvantages which should be enhance in the future.

i) This system used the neural network to classify the paddy disease. Neural network is one of the accurate techniques that can be applied in the system for detecting
 disease

- ii) This system can detect and classify the paddy disease automatically
- iii) This is an alternative or supplement to replace the traditional manual method
- iv) Human sight error can be eliminate when determine the paddy disease

5.6.2 Disadvantages

 The image produce in threshold process gain a lot of noises that make the system confuse to classify the paddy.

5.7 Achieved objectives

The objectives of this project are:

- i. to develop the prototype of paddy disease detection system
- ii. to detect the paddy disease by using image processing
- iii. to apply image processing technique to analyze the pattern of paddy disease

All the objectives stated above are achieved. The GUI has been developed for detect the paddy disease. This system also use some image processing in order to enhance the image and this system also applied the neural network technique to analyze the pattern of paddy disease which then classify it into 3 diseases.

5.8 Constraints

There are some constraints happen that effect the system when it is running. First, the image produces from the threshold process gain a lot of noise which affect the neural network training and testing. The quality of the image samples was blurring and sometimes it is not able to be process.

5.9 Assumption and further research

The system should be published and used in the agricultural sector especially in paddy industry to help the farmer detect the disease early. Farmer can protect their crop from being spread to the other crop area. Therefore, some assumptions have to be considered along with the development of this system. The camera must have a good pixel so that the picture was clear and easy to the system extract the feature.

In the view of the disadvantages stated, further research should be carried out to enhance the current research. There are some suggestion and recommendation for further research to publish the system in the agriculture industry. This system must be improved on the threshold method so that there is less noise or free noise. There still a lot of technique beside the Otsu method that can be implement in this system. Furthermore, this system should be implementing in the mobile application development. So that the paddy farmers just carry their phone to know the types of paddy disease.

5.10 Conclusion

A system for diagnosis the paddy disease has been developed using the Matlab application. The image processing techniques is applied to improve and enhance the image to a better quality. Besides, the neural network is used to classify the paddy diseases which are paddy blast, brown spot disease, narrow brown spot disease and normal paddy leaf. The methodology involves image acquisition, pre-processing and segmentation, analysis and classification of the paddy disease. All the paddy sample will be passing through the RGB calculation before it proceed to the binary conversion. If the sample is in the range of normal paddy disease sample will be convert into the binary data in excel file before proceed through the neural network for training and testing. Consequently, by employing the neural network technique, the paddy diseases are recognized about 92.5 percent accuracy rates. This prototype has a very great potential to be further improved in the future.

REFERENCES

1.Kurniawati, N.N. Abdullah, S.N.H.S. Abdullah.S. Investigation on Image Processing Techniques for Diagnosing Paddy Diseases". International Conference of Soft Computing and Pattern Recognition pg 273-277, 2009.

2.Meunkaewjinda, A. Kumsawat, P. Attakitmongcol, K. and Srikaew, A. "Grape Leaf Disease Detection from Color Imagery using Hybrid Intelligent System" Poceeding of ECTI-CON pg 513-516, 2008.

3.Ju-Won L., Han-Wook L., Jong-Hoe L., Ick-Tae K., Gun-Ki L., "A Study on Lung Nodule Detection usirlg Neural Networks" IEEE TENCON (1150-1153), 1999
Zhihao, Q. Minghua, Z. (2005). "Detection of rice sheath blight for in-season disease management using multispectral remote sensing"
Retrieved October 6, 2011 from

4.http://www.deepdyve.com/lp/elsevier/detection-of-rice-sheath-blight-for-in-season-diseasemanagement-using-tbRKqaZuHk?key=elsevier#Download

5.http://en.wikipedia.org/wiki/Neural_network Retrieved on 8 Disember 2011

6.Zulhadi, Zakarian. Nor Ashidi Mat Isa and Shahrel A. Suandi, Member IEEE "A Study on Neural Network Training Algorithm for Multiface Detection in Static Images" World Academy of Science, Engineering and Technology 62 2010, pg 170-173, 2010

7.http://en.wikipedia.org/wiki/Image_processing Retrived on 8 Disember 2011

8.http://www.merriam-webster.com/dictionary/rice?show=0&t=1320089188

9.http://wiki.tnau.ac.in/groups/paddycroppinginpapanasam/wiki/23961/Paddy_Diseases_Sympt oms_and_Management.html

10. http://en.wikipedia.org/wiki/Segmentation_(image_processing)

11. Eliza Yingzi Du and Chein-I Chang. 2003. An Unsupervised Approach to Colour Video Thresholding. IEEE Proc. International Conference on Multimedia and Expo ICME'03 3:III-337-40.

2. Meunkaewjinda, A. Kumsawat, P. Attakitmongcol, K. and Srikaew, A. "Grape Leaf Disease Detection from Color Imagery using Hybrid Intelligent System" Poceeding of ECTI-CON pg 513-516, 2008.

 Perona, P. and Malik, J. "Scale-space and edge detection using anisotropic diffusion," IEEE Transaction on pattern Analysis and Machine Intelligence, vol. 12, pp. 629-639, 1990
 Doungchatom, B. Kumsawat, P. Attakitmongcol, K. and A. Srikeaw, "Modified Self-Organizing Map for Optical Flow Clustering System," Proceeding of the 7th WSEAS International Conference on Signal, Speech and Image Processing, pp.61-69, 2007.

14. Liu C. and Wechsler H, "Gabor feature based classification using the enhanced fisher discriminant model for face recognition," IEEE Transaction on Image Processing, vol. 11, pp467-476, 2002.

15. Shen Weizheng, Wu Yachun, Chen zhanliang, Wei Hongda. "Grading Method of Leaf Spot Disease Based on Image Processing". In the proceedings of CSSE 2008.

16. S.C. Scardaci et al., Rice Blast: A New Disease in California, Agronomy Fact Sheat Series 1997-2, Department of Agronomy and Range Science University of California, Davis.

17. Robert W. Herdt, Research Priorities for Rice Biotechnology, in Rice Biotechnology, G.S. Khush and G.H. Toenniessen (eds.), Alden Press Ltd., London, 1991, pp. 35-37.

18. J.B. Cunha. 2003. Application of Image Processing Techniques in the Characterization of Plant Leafs. Proc. IEEE Intl' Symposium on Industrial Electronics.

19. K.J. Runtz. 1991. Electronic Recognition of Plant Spesies for Machine Vision Sprayer Control Systems. Proc. IEEE Western Canada Conference on Computer, Power, and Communication System in a Rural Environment (WESCANEX'91), pp. 84-88.

20. S.Dave and K. Runtz. 1995. Image Processing Methods for Identifying Species of Plants. Proc. IEEE Western Canada Conference on Computer, Power, and Communication System in a Rural Environment (WESCANEX'95), pp.403-408.

21.http://www.energizedwork.com/weblog/2005/07/iterative-and-incremental-development
Retrieved on 1 Disember 2011
22.http://en.wikipedia.org/wiki/Iterative_and_incremental_development
Retrieved on 1 Disember 2011

23.http://en.wikipedia.org/wiki/Product_software_implementation_method\ Retrieved on 1 Disember 2011

24,http://en.wikipedia.org/wiki/Software_testing\ Retrieved on 8 Disember 2011

25.http://en.wikipedia.org/wiki/Software_deployment Retrieved on 8 Disember 2011

26.http://en.wikipedia.org/wiki/Software_maintenance Retrieved on 8 Disember 2011

27.EBADI. GHASEM, "Evaluation of the Current Maintenance System and Development of an Improved Maintenance Management System in KIMIA Co" MSc thesis, NPC, GGSB,IUT, 2009

APPENDIX

A Gantt Chart

Project Task														We															
	1 2 Septe 20	mber	C	2 3 Octobe 2011		No	2 3 oveml 2011	ber		nber		Jan	3 nuar 012		2 epte 20	3 mbe 12		C	2 Octol 201	per		2 Nove 20	mbe		De	2 3 cemb 2012		1	lary
Identify project (problem to investigate) and scope of research																	Q- 12							6 10					
Discuss title with SV								-										č											
Plan work schedule						222			2-32	-8-	-20	-3	-2)	3(S	8—8)X23	-0				3—3		252			2)		
Discussion on related literature															<u>1</u> - 25										1				
Resubmission of literature review chapter 2		20						- 23 - 1	8-3	565			21-1	3-03	81-18		97 - 93 	- (2)				302		5 - ME			21	2	x13
Determine methodology															8		41	- 52			-				4.1 		5.6 · · :		
Submission methodology chapter 3									<u>11-1</u> 5	2 2			3									2-12							
Resubmission chapter 3																													+
Results and		3	1		1	838	1			1.8			3					-2		8	1	1 2					8		


Discussion			1 T	T II	1 1	12	1	Ĩ	- 10	ř-	Ē	1	-	12	Ĩ		Ť	1	19	- 19	Ť	1	Ť	3	n 1	T	-1
Write final draft and project overview																											
Submission project overview and final draft																											
Presentation PSM I																											
Submit written research proposal and abstract																											
Presentation of project progress to SV															1												
Initial stage of implementati on																											
Implementat ion																											
Result and Discussion							1 10															1		2			
Submission chapter 5 result and discussion	8 8	8 8 4	8	3 S			3:S			-5×73	;	23 é				2				\$—S		2		8			SZ

Submission chapter 6 conclusion			2	8 – 8 – 8 – 8		2)32-	8		x—3	- (3		i		-0			2		2	22	10
Submission of report draft and log book to faculty																					
Submission of project overview to faculty			2	2502		2)32	-2	0-0	× - 23	-9	, e			-9			43 - 54		2		
Distribution report draft and log book to supervisor							0		x - 21												
Distribution project overviewto evaluator																					
Presentation of PSM II															1						
Submission ofreport and CD			20				24 82			- 18				- 18	20						

Figure 5.10 Gantt Chart

B Plagiarism

		Assignment Inbox: PSM21213	
	Info	Dates	Similarity
psm2 draft submission	٦	Start 09-Dec-2012 8:58AM Due 26-Dec-2012 11:59PM Post 26-Dec-2012 12:00AM	42% Resubmit View




Result: 42 %

C Testing Result

No	Type of Paddy	Total	Sample	Sample	Accuracy for each
	Disease	Sample	Pass	Fail	sample
1.	Paddy Blast	10	9	1	90 %
2.	Narrow Brown Spot	10	10	-	100 %
	Disease				
3.	Brown Spot Disease	10	8	2	80 %
4.	Normal	10	10	-	100 %
	Total	40	37	3	92.5 %

Table 5.1 : Testing result



Type of Disease : Paddy Blast Disease

Paddy	Paddy Disease De	etection System	
Browse Process Original Image	Grayscale Image	Filtering	Threshold
	Paddy Disease and Management :	nitrogen in three split doses.Remo	ess N - fertilizer application. Apply oves weed hosts from bunds. Use of straw and stubbles after harvest.

Type of Disease : Paddy Blast Disease

dy	Paddy Disease De	etection System	
Browse Process Original Image	Grayscale Image	Fitering	Threshold
	Paddy Disease and Management : Reset	nitrogen in three split doses.Rem	cess N - fertilizer application. Apply noves weed hosts from bunds. Use of f straw and stubbles after harvest.

Type of Disease : Paddy Blast Disease

Paddy	Paddy Disease D	etection Syster	n
Browse Process Original Image	Grayscale Image	Filtering	Threshold
	Paddy Disease and Management :	of potassium and phosphorus f cultivars early in the growing se narrow brown leaf spot. The use approach to manage the diseas	ease. Cultural practices, such as the use fertilizers, and planting of early maturing ason, are recommended to manage the of resistant varieties is the most effective se. Spraying of fungicides such benomyl, zim, propiconazole, and iprodione
	Reset	Exit	

Type of Disease : Narrow Brown Spot Disease

	Paddy Disease D	etection System	
Browse Process	Grayscale Image	Filtering	Threshold
	Paddy Disease and Management :	of potassium and phosphorus fer cultivars early in the growing sea narrow brown leaf spot. The use of approach to manage the disease.	se. Cultural practices, such as the use tilizers, and planting of early maturing son, are recommended to manage the resistant varieties is the most effective Spraying of fungicides such benomyl, m, propiconazole, and iprodione
0	Reset	Exit	

Type of Disease : Narrow Brown Spot Disease

ddy	Paddy Disease D	etection Syste	:m
Browse Process Original Image	Grayscale Image	Filtering	Threshold
18.	181	of potassium and phosphoru:	sease. Cultural practices, such as the use sfertilizers, and planting of early maturing
	Paddy Disease and Management :	narrow brown leaf spot. The us approach to manage the dise	season, are recommended to manage the e of resistant varieties is the most effective ase. Spraying of fungicides such benomyl, lazim, propiconazole, and iprodione

Type of Disease : Narrow Brown Spot Disease

and the second sec	Paddy Disease De	etection Systen	l
Browse Process Original Image	Grayscale Image	Filtering	Threshold
	Paddy Disease and Management :	t/ha. Soil application of P. fluor FYM after 30 days of transplar @0.2percent at boot leaf st	ly FYM 12.5 t/ha or green manure 6.25 escens @ 2.5 kg/ha mixed with 50 kg ting, Foliar spraying of P. Huorescens age and 10 days later.Avoid flow of n infected to healthy field.
	Reset	Exit	

Type of Disease: Brown Spot Disease

Paddy	Paddy Disease D	etection System	
Browse Process	Grayscale Image	Filtering	Threshold
	Paddy Disease and Management :	of potassium and phosphorus fert cultivars early in the growing seas narrow brown leaf spot. The use of approach to manage the disease.	e. Cultural practices, such as the use ilizers, and planting of early maturing on, are recommended to manage the resistant varieties is the most effective Spraying of fungicides such benomyl, n, propiconazole, and iprodione
	Reset	Exit	

Type of Disease: Brown Spot Disease

Result : Fail

Paddy Disease Detection System				
Browse Process Original Image	Grayscale Image	Filtering	Threshold	
0 0	Paddy Disease and Management :	t/ha. Soil application of P. fluor FYM after 30 days of transpla @0.2percent at boot leaf st	oly FYM 12.5 t/ha or green manure 6.25 escens @ 2.5 kg/ha mixed with 50 kg niting.Foliar spraying of P. fluorescens age and 10 days later.Avoid flow of m infected to healthy field.	

Type of Disease: Brown Spot Disease

Paddy	Paddy Disease D	etection Syste	em	
Browse Process Original Image	Grayscale Image	Filtering	Threshold	
	Paddy Disease and Management		Type 4 :Normal	
	Reset	Exit		

Type of Disease: Normal Paddy

Paddy Disease Detection System					
Browse Process Original Image	Grayscale Image	Filtering	Threshold		
	Paddy Disease and Management :	Т	Type 4 : Normal		
	Reset	Exit			

Type of Disease: Normal Paddy

Paddy	Paddy Disease De	tection System		
Browse Process Original Image	Grayscale Image	Filtering	Threshold	
	Paddy Disease and Management :	Type 4 :	Normal	
	Reset	Exit		

Type of Disease: Normal Paddy

Paddy	Paddy Disease De	etection Syste	em
Browse Process Original Image	Grayscale Image	Filtering	Threshold
	Paddy Disease and Management :	nitrogen in three split doses	id excess N - fertilizer application. Apply Removes weed hosts from bunds. Use of ng of straw and stubbles after harvest.
	Reset	Exit	

Type of Disease: Paddy Blast

Browse Process Grayscale Ima Grayscale Ima Grayscale Ima	sease Det	tection System	n
Paddy Disease and	age	Fitering	Threshold
		t/ha. Soil application of P. fluor FYM after 30 days of transpla @0.2percent at boot leaf st	ply FYM 12.5 t/ha or green manure 6.25 rescens @ 2.5 kg/ha mixed with 50 kg nting. Foliar spraying of P. fluorescens tage and 10 days later.Avoid flow of m infected to healthy field.

Type of Disease: Paddy Blast

Result : Fail

Browse Process	Grayscale Image	Filtering	Threshold
Original Image	12		
	Paddy Disease and Management :	nitrogen in three split doses.Remo	ess N - fertilizer application. Apply wes weed hosts from bunds. Use of straw and stubbles after harvest.

Type of Disease: Paddy Blast

	Paddy Disease Detection System				
Browse Process Original Image	Grayscale Image	Filtering	Threshold		
	Paddy Disease and Management :	of potassium and phosphoru cultivars early in the growing narrow brown leaf spot.The u approach to manage the dise	Disease. Cultural practices, such as the use us fertilizers, and planting of early maturing s season, are recommended to manage the se of resistant varieties is the most effective ease. Spraying of fungicides such benomyl, idazim, propiconazole, and iprodione		
	Reset	Exit			

Type of Disease: Narrow Brown Spot Disease

Paddy Disease Detection System					
Browse Process Original Image	Grayscale Image	Filtering	Threshold		
	Paddy Disease and Management :	of potassium and phosphorus for cultivars early in the growing se narrow brown leaf spot. The use approach to manage the diseas	ease. Cultural practices, such as the use ertilizers, and planting of early maturing ason, are recommended to manage the of resistant varieties is the most effective e. Spraying of fungicides such benomyl, zim, propiconazole, and iprodione		
	Reset	Exit			

Type of Disease: Narrow Brown Spot Disease

and here any he	Paddy Disease D	etection Syster	n
Browse Process Original Image	Grayscale Image	Filtering	Threshold
	Paddy Disease and Management :	of potassium and phosphorus f cultivars early in the growing se narrow brown leaf spot. The use approach to manage the diseas	ase. Cultural practices, such as the use ertilizers, and planting of early maturing tason, are recommended to manage the of resistant varieties is the most effective e. Spraying of fungicides such benomyl, zim, propiconazole, and iprodione
	Reset	Exit	


Type of Disease: Narrow Brown Spot Disease

Paddy	Paddy Disease De	etection System	
Browse Process Original Image	Grayscale Image	Filtering	Threshold
	Paddy Disease and Management :	t/ha. Soil application of P. fluores FYM after 30 days of transplanti @0.2percent at boot leaf stag	FYM 12.5 t/ha or green manure 6.25 .cens @ 2.5 kg/ha mixed with 50 kg ng.Foliar spraying of P. fluorescens e and 10 days later.Avoid flow of infected to healthy field.
	Reset	Exit	

Type of Disease: Brown Spot Paddy Disease

Paddy	Paddy Disease D	etection System	
Browse Process Original Image	Grayscale Image	Fitering	Threshold
	Paddy Disease and Management	t/ha. Soil application of P. fluores FYM after 30 days of transplanti @0.2percent at boot leaf stag	r FYM 12.5 t/ha or green manure 6.25 scens @ 2.5 kg/ha mixed with 50 kg ing.Foliar spraying of P. fluorescens je and 10 days later.Avoid flow of infected to healthy field.

Type of Disease: Brown Spot Paddy Disease



Type of Disease: Brown Spot Paddy Disease

Paddy	Paddy Disease D	etection System	m
Browse Process Original Image	Grayscale Image	Filtering	Threshold
	Paddy Disease and Management		pe 4 :Normal
	Reset	Exit	

Type of Disease: Normal Paddy

Paddy Paddy Disease Detection System					
Browse Process Original Image	Grayscale Image	Filtering	Threshold		
	Paddy Disease and Managemen		ype 4 :Normal		
Reset					

Type of Disease: Normal Paddy

Browse Process original mage Filtering Treshold Signal mage Signal	Paddy	Paddy Disease De	etection Syste	em
		Grayscale Image	Filtering	Threshold
· · · · · · · · · · · · · · · · · · ·		Paddy Disease and Management :		Type 4 :Normal

Type of Disease: Normal Paddy

Browse Process original Image Image Image Image I	And property of the second	Paddy Disease De	etection Syster	n
Paddy Disease and Management: Paddy Disease and Management: of potassium and phosphorus fertilizers, and planting of early maturing cultivars early in the growing season, are recommended to manage the narrow brown leaf spot. The use of resistant varieties is the most effective approach to manage the disease. Spraying of fungicides such benomyl,		Grayscale Image	Filtering	Threshold
	1.1.1	Paddy Disease and Management :	of potassium and phosphorus for cultivars early in the growing se narrow brown leaf spot. The use approach to manage the diseas	ertilizers, and planting of early maturing ason, are recommended to manage the of resistant varieties is the most effective e. Spraying of fungicides such benomyl,

Type of Disease: Narrow Brown Spot Disease

ły	Paddy Disease D	etection Syster	n
Browse Process Original Image	Grayscale Image	Filtering	Threshold
() (後)	Paddy Disease and Management :	of potassium and phosphorus f cultivars early in the growing se narrow brown leaf spot. The use approach to manage the diseas	ease. Cultural practices, such as the use fertilizers, and planting of early maturing eason, are recommended to manage the of resistant varieties is the most effective se. Spraying of fungicides such benomyl, zim, propiconazole, and iprodione
	Reset	Exit	

Type of Disease: Narrow Brown Spot Disease

Paddy Disease Detection System				
Browse Process Original Image	Grayscale Image	Filtering	Threshold	
	Paddy Disease and Management :	of potassium and phosphorus fert cultivars early in the growing seas narrow brown leaf spot. The use of approach to manage the disease.	se. Cultural practices, such as the use tilizers, and planting of early maturing son, are recommended to manage the resistant varieties is the most effective Spraying of fungicides such benomyl, m, propiconazole, and iprodione	
	Reset	Exit		


Type of Disease: Narrow Brown Spot Disease

Paddy	Paddy Disease D	etection System	
Browse Process Original Image	Grayscale Image	Filtering	Threshold
and the	Paddy Disease and Management :	of potassium and phosphorus ferti cultivars early in the growing seas narrow brown leaf spot. The use of r approach to manage the disease.	e. Cultural practices, such as the use lizers, and planting of early maturing on, are recommended to manage the esistant varieties is the most effective Spraying of fungicides such benomyl, I, propiconazole, and iprodione
	Reset	Exit	

Type of Disease: Narrow Brown Spot Disease

ły	Paddy Disease De	etection System	n
Browse Process	Grayscale Image	Filtering	Threshold
3	Paddy Disease and Management :	t/ha. Soil application of P. fluor FYM after 30 days of transpla @0.2percent at boot leaf st	bly FYM 12.5 t/ha or green manure 6.25 escens @ 2.5 kg/ha mixed with 50 kg tring. Foliar spraying of P. fluorescens age and 10 days later.Avoid flow of m infected to healthy field.
	Reset	Exit	

Type of Disease: Brown Spot Paddy Disease



Type of Disease: Brown Spot Paddy Disease

Paddy Disease Detection System					
Browse Process Original Image	Grayscale Image	Filtering	Threshold		
	Paddy Disease and Management :	of potassium and phosphorus fer cultivars early in the growing sea narrow brown leaf spot. The use of approach to manage the disease	se. Cultural practices, such as the use tilizers, and planting of early maturing son, are recommended to manage the f resistant varieties is the most effective . Spraying of fungicides such benomyl, m, propiconazole, and iprodione		
	Reset	Exit			

Type of Disease: Brown Spot Paddy Disease

Result : Fail

Paddy Disease Detection System					
Browse Process Original Image	Grayscale Image	Filtering	Threshold		
	Paddy Disease and Management :	t/ha. Soil application of P. fluoresc FYM after 30 days of transplantin @0.2percent at boot leaf stage	FYM 12.5 t/ha or green manure 6.25 cens @ 2.5 kg/ha mixed with 50 kg g.Foliar spraying of P. fluorescens e and 10 days later.Avoid flow of nfected to healthy field.		
	Reset	Exit			

Type of Disease: Brown Spot Paddy Disease

Pade	dy	Paddy Disease I	Detection System		
	Browse Process Original Image	Grayscale Image	Fittering	Threshold	
		Paddy Disease and Managemen		e 4 :Normal	
		Reset	Exit		

Type of Disease: Normal Paddy

Paddy	Paddy Disease De	tection System				
Browse Process Original Image	Grayscale Image	Filtering	Threshold			
	Paddy Disease and Management :	Туре	4 :Normal			
	Reset					

Type of Disease: Normal Paddy

Paddy Paddy Disease Detection System							
Browse Process Original Image	Grayscale Image	Filtering	Threshold				
	Paddy Disease and Management :	1	Type 4 :Normal				
	Reset Exit						

Type of Disease: Normal Paddy

Paddy	Paddy Disease [Detection System	n
Browse Process Original Image	Grayscale Image	Filtering	Threshold
	Paddy Disease and Managemen		e 4 :Normal
	Reset	Exit	

Type of Disease: Normal Paddy

↓ Paddy	Paddy Disease De	etection System	
Browse Process Original Image	Grayscale Image	Filtering	Drreshold
	Paddy Disease and Management :	nitrogen in three split doses.Remo	ess N - fertilizer application. Apply oves weed hosts from bunds. Use of straw and stubbles after harvest.
	Reset	Exit	

Type of Disease: Paddy Blast

dy	Paddy Disease De	etection System	n
Browse Process Original Image	Grayscale Image	Filtering	Threshold
9.	Paddy Disease and Management :	nitrogen in three split doses.Rer	xcess N - fertilizer application. Apply moves weed hosts from bunds. Use of of straw and stubbles after harvest.
	Reset	Exit	

Type of Disease: Paddy Blast

Paddy	Paddy Disease De	etection System		
Browse Process Original Image	Grayscale Image	Filtering	Treshold	
	Paddy Disease and Management :	nitrogen in three split doses.Remo	ess N - fertilizer application. Apply oves weed hosts from bunds. Use of straw and stubbles after harvest.	

Type of Disease: Paddy Blast

Paddy	Paddy Disease De	etection System	
Browse Process Original Image	Grayscale Image	Filtering	Treshold
	Paddy Disease and Management :	nitrogen in three split doses.Remo	ess N - fertilizer application. Apply wes weed hosts from bunds. Use of traw and stubbles after harvest.
	Reset	Exit	

Type of Disease: Paddy Blast

D Calculation of RGB for Normal paddy

Sample 1

value	Max val	Min value	RGB	
23	123	93	Red	
71	171	142	Green	
24	124	89	Blue	

Table 5.2: RGB calculation of Sample 1

Sample 2

 RGB	Min value	Max value
Red	102	141
Green	143	184
Blue	111	155

Table 5.3: RGB calculation of Sample 2

Sample 3

RGB	Min value	Max value
Red	147	172
Green	169	191
Blue	69	89

Table 5.4: RGB calculation of Sample 3

Sample 4

RGB	Min value	Max value
Red	157	187
Green	179	209
Blue	95	126

Table 5.5: RGB calculation of Sample 4

Sample 5

RGB	Min value	Max value
Red	102	141
Green	143	184
Blue	111	155

Table 5.6: RGB calculation of Sample 5

Sample 6

RGB	Min value	Max value
Red	144	188
Green	168	210
Blue	80	136

Table 5.7: RGB calculation of Sample 6

Sample 7

RGB	Min value	Max value
Red	148	204
Green	167	205
Blue	74	122

Table 5.8: RGB calculation of Sample 7

Sample 8

RGB	Min value	Max value
Red	150	188
Green	173	210
Blue	83	132

Table 5.9: RGB calculation of Sample 8

Sample 9

RGB	Min value	Max value
Red	145	185
Green	168	209
Blue	80	125

Table 5.10: RGB calculation of Sample 9

Sample 10

RGB	Min value	Max value
Red	136	181
Green	151	199
Blue	64	117

Table 5.11: RGB calculation of Sample 10

Total range RGB of 10 samples are:

Red : 93 - 204

Green: 142 - 210

Blue : 64 - 155