

**PRODUCTION OF PRAVASTATIN BY FILAMENTOUS FUNGI
ISOLATED FROM SOIL**

EMINE SEYDAMETOVA

Thesis submitted in fulfilment of the requirements
for the award of the degree of
Doctor of Philosophy (Bio-process Engineering)

Faculty of Chemical and Natural Resources Engineering
UNIVERSITI MALAYSIA PAHANG

AUGUST 2013

ABSTRACT

Pravastatin is a clinically useful cholesterol-lowering agent, selectively inhibiting 3-hydroxy-3-methylglutaryl-coenzyme A reductase, the regulatory enzyme in cholesterol biosynthesis. Currently, industrial production of this statin is based on a two-step fermentation process: the initial production of compactin and its subsequent biotransformation to pravastatin. The development of a one-step fermentation process using pravastatin-producing microfungi may be a commercially attractive approach. To facilitate this, isolation of novel fungal strains from different natural sources and their screening for pravastatin production is required. Soil being a reservoir for a wide variety of filamentous fungi has been recognized for a long time. In this study, 54 fungal cultures were obtained from soil samples collected in Pahang State (Malaysia). Isolates were cultivated in submerged fermentation and tested for their ability to produce pravastatin using high-performance liquid chromatography. Five selected pravastatin producers were identified to species level using cultural and morphological characteristics, physiological and biochemical tests, and molecular techniques. Screening of the important variables affecting pravastatin production by the best of these producers was initially carried out using 2^{7-3} fractional factorial design and these selected variables were then optimized using rotatable central composite design. Kinetic studies of substrate uptake, fungal growth and pravastatin production in shake flask culture under optimized conditions were also conducted. Among 25 *Penicillium* isolates that were capable of producing pravastatin directly by fermentation, only five (ESF2M, ESF19M, ESF20P, ESF21P, and ESF26P) did so in relatively high concentrations, with *Penicillium* sp. ESF21P being the most active pravastatin producer, achieving a concentration of 196.83 mg/L. Fungal identification methods used in this study confirmed that the isolates *Penicillium* sp. ESF2M and ESF19M are referable to *Penicillium citrinum*, *Penicillium* sp. ESF20P and ESF26P were most closely related to *Penicillium janthinellum*, and *Penicillium* sp. ESF21P showed the highest homology with *Eupenicillium brefeldianum*. All sequence data from this study have been deposited in the GenBank database. A maximum concentration of 234.36 mg/L of pravastatin was produced by *E. brefeldianum* ESF21P under the optimized conditions suggested by the Design-Expert 6.0.8 software. Pravastatin fermentation using this fungus showed the typical kinetics of a secondary metabolite, with maximum yield obtained after about 288 h of fermentation.

ABSTRAK

Pravastatin merupakan satu agen perendah-kolesterol yang merencat 3-hidroksi-3-methylglutaril-koenzim A reductase secara klinikal. 3-hidroksi-3-methylglutaril-koenzim A reductase merupakan enzim pengatur dalam biosintesis kolesterol. Sehingga kini penghasilan statin adalah berasaskan satu proses fermentasi dua-langkah: penghasilan kompaktin pada peringkat permulaan dan diikuti dengan biotransformasinya kepada pravastatin. Pembangunan satu proses fermentasi satu-langkah menggunakan mikrofungi penghasil-pravastatin adalah satu pendekatan komersil yang menarik. Pemencilan strain fungi novel daripada sumber semulajadi dan penyaringannya adalah perlu untuk penghasilan pravastatin. Sejak dahulu lagi tanah merupakan sumber terbaik untuk pelbagai fungi berfilamen. Dalam kajian ini, 54 kultur fungi didapati daripada sampel-sampel tanah yang dikutip dalam negeri Pahang (Malaysia). Pencilan-pencilan ini dibiakkan dalam fermentasi tenggelam dan diuji kemampuannya untuk menghasilkan pravastatin menggunakan kromatografi cecair berprestasi tinggi. Lima spesis fungi penghasil pravastatin telah dikenal pasti daripada pencirian kultur dan morfologi, ujian-ujian fisiologi dan biokimia, dan teknik-teknik molekul. Penyaringan pembolehubah-pembolehubah penting yang mempengaruhi penghasilan pravastatin oleh penghasil terbaik telah dilakukan pada peringkat awal dengan menggunakan rekabentuk faktorial 2^{7-3} . Kemudian aras pembolehubah-pembolehubah yang dipilih melalui saringan ini dioptimumkan dengan menggunakan rekabentuk boleh-putar komposit berpusat. Kajian kinetik penggunaan substrat, tumbesaran fungi dan penghasilan pravastatin dalam kultur kelalang goncang pada keadaan optimum telah juga dilakukan. Daripada 25 pencilan *Penicillium* yang boleh menghasilkan pravastatin, hanya lima (ESF2M, ESF19M, ESF20P, ESF21P, dan ESF26P) boleh menghasilkannya dalam kepekatan yang relatif tinggi. Didapati *Penicillium* sp. ESF21P merupakan penghasil pravastatin yang paling aktif, mencapai kepekatan 196.83 mg/L. Kaedah-kaedah pengenalpastian fungi yang digunakan dalam kajian ini telah mengesahkan bahawa pencilan-pencilan *Penicillium* sp. ESF2M dan ESF19M boleh dirujuk kepada *Penicillium citrinum*. *Penicillium* sp. ESF20P dan ESF26P adalah paling berkait rapat dengan *Penicillium janthinellum*, dan *Penicillium* sp. ESF21P menunjukkan homologi tertinggi dengan *Eupenicillium brefeldianum*. Semua data jujukan daripada kajian ini telah telah disimpan di pengkalan data GenBank. Pada keadaan optimum yang dicadangkan oleh perisian Design-Expert 6.0.8, *E. brefeldianum* ESF21P telah mencatatkan kepekatan maksimum pravastatin (234.36 mg/L). Fermentasi pravastatin menggunakan fungi ini menghasilkan nilai kinetik tipikal bagi metabolit sekunder di mana hasil maksimum dicapai selepas kira-kira 288 jam fermentasi.

TABLE OF CONTENTS

	Page
SUPERVISOR'S DECLARATION	ii
STUDENT'S DECLARATION	iii
DEDICATION	iv
ACKNOWLEDGEMENTS	v
ABSTRACT	vi
ABSTRAK	vii
TABLE OF CONTENTS	viii
LIST OF TABLES	xiv
LIST OF FIGURES	xviii
LIST OF ABBREVIATIONS	xxiv
GLOSSARY	xxx

CHAPTER 1 INTRODUCTION

1.1	Background	1
1.2	Problem Statement	4
1.3	Aim of the Study	6
1.4	Research Objectives	7
1.5	Scope of the Study	7
1.6	Rationale and Significance	8
1.7	Novelty of the Study	8

CHAPTER 2 LITERATURE REVIEW

2.1	Introduction	10
2.2	Cholesterol Biosynthesis Pathway	11

2.3	Historical Development of Statins	14
2.4	Chemical Structure of Statins	18
2.5	Mechanism of Action	20
2.6	Current and Potential Biomedical Applications of Statins	22
2.7	Soil Microfungi Identification Approaches	31
	2.7.1 Oil Palm Plantation and Mangrove Soils in Malaysia as Valuable Sources of Microfungi	31
	2.7.2 Various Identification Techniques Applied in Mycology	34
2.8	Biosynthesis of Pravastatin, a Natural HMG-CoA Reductase Inhibitor	40
	2.8.1 Biosynthesis of Compactin (Precursor of Pravastatin)	40
	2.8.2 Microbial Hydroxylation of Compactin to Pravastatin	44
2.9	Strain Improvement Strategies for Increased Compactin Synthesis and Hydroxylation	46
2.10	Optimization of Pravastatin Production	50
	2.10.1 Some Strategies for Optimization	50
	2.10.2 Nutritional Variables Improvement	53
	2.10.3 Optimization of Levels of the Significant Fermentation Variables	59
2.11	Fermentative Production of Pravastatin	65
	2.11.1 Batch Fermentation Processes for Compactin Production	65
	2.11.2 Synthesis of Compactin Using Immobilized Fungal Spores	67
	2.11.3 Solid State Fermentation for the Manufacture of Compactin	69
	2.11.4 Bioconversion of Compactin into Pravastatin	73
	2.11.5 Direct Microbial Production of Pravastatin and Its Kinetics	77
2.12	Concluding Remarks	79

CHAPTER 3 GENERAL MATERIALS AND METHODS

3.1	Chemical Reagents and Nutrient Media	81
3.2	Spore Inoculum Preparation	82
3.3	Fermentation Procedure	82
3.4	Analytical Determination of Statins	83

3.5	Dry Cell Weight Estimation	84
3.6	Statistical Analysis	84

**CHAPTER 4 ISOLATION AND MORPHOLOGICAL
CHARACTERIZATION OF SOIL FILAMENTOUS
FUNGI**

4.1	Introduction	85
4.2	Materials and Methods	86
	4.2.1 Chemical Reagents and Nutrient Media	86
	4.2.2 Soil Sampling	86
	4.2.3 Isolation Technique for Soil Filamentous Fungi	89
	4.2.4 Morphological Identification of Fungal Isolates	90
	4.2.5 Statistical Analysis	90
4.3	Results	91
	4.3.1 Isolation of Filamentous Fungi from Oil Palm Plantation Soils	91
	4.3.2 Isolation of Filamentous Fungi from Mangrove Soils	95
	4.3.3 Macro- and Micro-morphology of Isolated Fungi	98
4.4	Discussion	129
	4.4.1 Fungal Isolates from Oil Palm Plantation Soils	129
	4.4.2 Fungal Isolates from Mangrove Soils	130
	4.4.3 Morphological Identification of Isolated Fungi	133
4.5	Concluding Remarks	134

**CHAPTER 5 SCREENING OF *PENICILLIUM* ISOLATES FOR
STATIN-PRODUCING ABILITY**

5.1	Introduction	135
5.2	Materials and Methods	136
	5.2.1 Chemical Reagents and Nutrient Media	136
	5.2.2 Fermentation Procedure	136
	5.2.3 Analytical Determination of Statins	136
	5.2.4 Dry Cell Weight Estimation	137
	5.2.5 Statistical Analysis	137

5.3	Results	137
	5.3.1 Statin-producing Ability of Fungal Isolates	137
5.4	Discussion	141
	5.4.1 Screening for Natural Statin Production	141
5.5	Concluding Remarks	144

CHAPTER 6 SPECIES LEVEL IDENTIFICATION OF ACTIVE PRAVASTATIN PRODUCERS

6.1	Introduction	145
6.2	Materials and Methods	146
	6.2.1 Chemical Reagents and Nutrient Media	146
	6.2.2 Fungal Isolates	146
	6.2.3 Cultural and Morphological Studies	146
	6.2.4 Physiological Characterization	147
	6.2.5 Ehrlich Test	148
	6.2.6 Enzyme Assays	148
	6.2.7 Micro-scale Extraction and Analysis for Secondary Metabolite Profiling	149
	6.2.8 Fungal Identification Using the Biolog System	151
	6.2.9 Molecular Identification	153
6.3	Results	156
	6.3.1 Cultural and Morphological Features of the Fungal Isolates	156
	6.3.2 Physiological Separation of <i>Penicillium</i> Isolates	158
	6.3.3 Ehrlich Reaction	161
	6.3.4 Enzyme Activities of the Fungal Isolates	162
	6.3.5 Secondary Metabolite Profiling	164
	6.3.6 Identification of the Fungal Isolates Using Biolog System	170
	6.3.7 Molecular Identification of the Fungal Isolates Based on ITS1-5.8S-ITS2 Region of rDNA	174
	6.3.8 Classification of the Identified Species	182
6.4	Discussion	184
	6.4.1 Classical Morphological Identification of the Fungal Isolates	184
	6.4.2 Physiological Criteria as an Aid for Fungal Isolates Separation	189

6.4.3 Biochemical Tests as an Aid to the Rapid Identification of <i>Penicillium</i> Isolates	192
6.4.4 Molecular Identification of the <i>Penicillium</i> isolates	200
6.5 Concluding Remarks	201

CHAPTER 7 OPTIMIZATION OF PRAVASTATIN PRODUCTION BY *EUPENICILLIUM BREFELDIANUM* ESF 21P

7.1	Introduction	203
7.2	Materials and Methods	204
	7.2.1 Chemical Reagents and Nutrient Media	204
	7.2.2 Fungal culture	205
	7.2.3 Fermentation Procedure for Media Screening	205
	7.2.4 The 2^{7-3} Fractional Factorial Experiments	206
	7.2.5 Optimization of Pravastatin Production Using Rotatable Central Composite Design	209
	7.2.6 Analytical Determination of Pravastatin	213
7.3	Results	213
	7.3.1 Fermentation Media Screening	213
	7.3.2 Screening of Variables with Significant Effects on Pravastatin Production by <i>Eupenicillium brefeldianum</i> ESF 21P	214
	7.3.3 Optimization of Pravastatin Production Using Response Surface Methodology	224
7.4	Discussion	243
	7.4.1 Selection of Optimal Fermentation Medium for Pravastatin Production	243
	7.4.2 Assessment of the Effects of Different Variables	245
	7.4.3 Determination and Validation of Optimal Conditions	252
7.5	Concluding Remarks	257

CHAPTER 8 KINETIC STUDY OF PRAVASTATIN FERMENTATION BY *EUPENICILLIUM BREFELDIANUM* ESF 21P

8.1	Introduction	259
8.2	Materials and Methods	260
	8.2.1 Chemical Reagents and Nutrient Media	260
	8.2.2 Fungal Culture and Fermentation Procedure	261
	8.2.3 Analytical Determination of Pravastatin	261
	8.2.4 Analytical Determination of Substrates	261
	8.2.5 Dry Cell Weight Estimation	262
	8.2.6 Statistical Analysis	262
	8.2.7 Determination of Kinetic Parameters	262
8.3	Results	265
	8.3.1 Time Course Profiles of Fungal Growth, Substrate Uptake and Pravastatin Production	265
	8.3.2 Analysis of Kinetic Parameters of <i>Eupenicillium brefeldianum</i> ESF 21P Fermentation	267
8.4	Discussion	269
	8.4.1 Kinetics of Growth and Pravastatin Fermentation by <i>Eupenicillium brefeldianum</i> ESF 21P	269
8.5	Concluding Remarks	273

CHAPTER 9 CONCLUSIONS AND RECOMMENDATIONS

9.1	Conclusions	275
9.2	Contribution	277
9.3	Recommendations for the Future Research	279
REFERENCES		281
APPENDICES		314
A	Recipe of Agar Media	314
B	High-performance Liquid Chromatograms	319
C	API ZYM Test Results	323
D	Biolog Data	326

E	Commercial Kits and Reagents for Molecular Identification of Fungi	334
F	List of Publications	339
G	Production of Lovastatin by <i>Penicillium</i> spp. Soil Microfungi (manuscript)	342
H	Pravastatin: Microbial Production and Biomedical Applications (monograph's preface)	346

LIST OF TABLES

Table No.	Title	Page
2.1	Commercially available HMG-CoA reductase inhibitors (statins) (adapted from U.S. Food and Drug Administration: approved drug products, 2012)	17
2.2	Comparison of various reports on compactin production (adapted from Chakravarti and Sahai, 2004)	58
4.1	Soil samples obtained from oil palm plantations in Gambang (3.72°-N 103.12°-E)	87
4.2	Rhizosphere soil sampling from different plants in mangrove forests (Pahang State, Malaysia)	88
4.3	Characteristics of soil samples obtained from oil palm plantations, and the number of recovered isolates	91
4.4	Characteristics of composite soil samples obtained from mangrove forests, and the number of recovered isolates	96
5.1	Natural statins production by <i>Penicillium</i> spp. microfungi isolated from oil palm plantation soils, detected by HPLC analysis (ND=not detected, DCW=dry cell weight)	137
5.2	Natural statins production by <i>Penicillium</i> spp. microfungi isolated from mangrove soils, detected by HPLC analysis (ND=not detected, DCW=dry cell weight)	139
6.1	List of examined <i>Penicillium</i> isolates	156
6.2	Cultural properties of <i>Penicillium</i> isolates (CD = colony diameter; COC = colony obverse color; CRC = colony reverse color)	157
6.3	Morphological properties of <i>Penicillium</i> isolates	157
6.4	Eco-physiological data of the examined <i>Penicillium</i> isolates	159

6.5	Results of physiological tests carried out on the five selected isolates	161
6.6	Enzymes detected by the API ZYM test and their respective substrates ((n) = negative reaction (0); (m) = moderate reaction (scale 1 or 2); (p) = positive reaction (scale 3-5))	163
6.7	Species identification for <i>Penicillium</i> sp. ESF19M isolate after 10 d of incubation (PROB=probability; SIM=similarity index; DIST=distance)	171
6.8	Species identification for <i>Penicillium</i> sp. ESF20P isolate after 3 d of incubation (PROB=probability; SIM=similarity index; DIST=distance)	171
6.9	Species identification for <i>Penicillium</i> sp. ESF21P isolate after 4 d of incubation (PROB=probability; SIM=similarity index; DIST=distance)	172
6.10	Identified fungal isolates with relationship to the species listed in the National Center for Biotechnology Information (NCBI) website (http://www.ncbi.nlm.nih.gov/ , 30 July 2012)	181
6.11	Cultural and morphological characteristics of <i>Penicillium</i> sp. ESF2M and ESF19M in comparison with those reported in the literature for <i>P. citrinum</i> and <i>P. brevicompactum</i> (CD = colony diameter; CRC = colony reverse color)	184
6.12	Cultural and morphological characteristics of <i>Penicillium</i> sp. ESF20P and ESF26P in comparison with those reported in the literature for <i>P. janthinellum</i> (CD = colony diameter; CRC = colony reverse color)	185
6.13	Cultural and morphological characteristics of <i>Penicillium</i> sp. ESF21P in comparison with those reported in the literature for <i>E. brefeldianum</i> (CD = colony diameter; CRC = colony reverse color)	186
6.14	Physiological characteristics for <i>Penicillium</i> sp. ESF2M and ESF19M in comparison with those reported in the literature for <i>P. citrinum</i> and <i>P. brevicompactum</i>	191

6.15	API ZYM results for <i>Penicillium</i> sp. ESF2M and <i>Penicillium</i> sp. ESF19M in comparison with those reported in the literature for <i>P. citrinum</i> and <i>P. brevicompactum</i> ((n) = negative reaction (0); (m) = moderate reaction (scale 1 or 2); (p) = positive reaction (scale 3-5))	193
6.16	API ZYM results for <i>Penicillium</i> sp. ESF20P and ESF26P in comparison with those reported in the literature for <i>P. janthinellum</i> ((n) = negative reaction (0); (m) = moderate reaction (scale 1 or 2); (p) = positive reaction (scale 3-5))	194
7.1	Variables and their coded and actual levels used in the method of 2^{7-3} fractional factorial experiments	207
7.2	The design of the 2^{7-3} fractional factorial experiments and the overall randomization of runs (SA=slant age; SC=spore concentration; IV=inoculum volume; FT=fermentation time; T=temperature; pH=initial pH of the medium; AR=agitation rate)	207
7.3	The levels of the variables in the rotatable central composite design based on the full 2^4 factorial experiments	210
7.4	The rotatable central composite design based on the full 2^4 factorial experiments (SA=slant age; FT=fermentation time; pH=initial pH of the medium; AR=agitation rate)	211
7.5	Pravastatin production in different fermentation media	214
7.6	Matrix corresponding to the 2^{7-3} fractional factorial experiments together with the observed experimental and predicted data (SA=slant age; SC=spore concentration; IV=inoculum volume; FT=fermentation time; T=temperature; pH=initial pH of the medium; AR=agitation rate)	215
7.7	The percentage contributions of the main effect of each variable and of the interactive effects between variables on the response (pravastatin concentration)	216
7.8	Evaluation of the factorial model using analysis of variance (ANOVA)	219

7.9	The rotatable central composite design based on full 2^4 factorial experiments with experimental results and corresponding predictions (SA=slant age; FT=fermentation time; pH=initial pH of the medium; AR=agitation rate)	225
7.10	Results of the rotatable central composite design after square root transformation (SA=slant age; FT=fermentation time; pH=initial pH of the medium; AR=agitation rate)	226
7.11	Lack of fit tests from Design-Expert 6.0.8 software	229
7.12	Analysis of variance (ANOVA) for evaluation of the second-order model	231
7.13	Suggested solution from the Design-Expert 6.0.8 software for maximum pravastatin production	240
7.14	The validation experiments in 250-mL shake flasks (actual versus predicted values)	242
8.1	Shake flask fermentation data of <i>E. brefeldianum</i> ESF21P	263
8.2	Calculated specific growth rate values	267
8.3	Comparison between pravastatin yield coefficients on different substrates (glucose and glycerol)	269

LIST OF FIGURES

Figure No.	Title	Page
2.1	The cholesterol biosynthesis pathway (adapted from Tobert, 2003)	12
2.2	Statin side chains linked at C-8 (R_1) and C-6 (R_2) of the base structure (adapted from Manzoni and Rollini, 2002)	18
2.3	Chemical structures of the synthetic statins: atorvastatin and fluvastatin (adapted from Manzoni and Rollini, 2002)	19
2.4	Structural analogy between HMG-CoA and the HMG-like moiety of statins (adapted from Manzoni and Rollini, 2002)	21
2.5	Model of the biological effects of statins (adapted from Barrios-González and Miranda, 2010)	27
2.6	Oil palm cultivation area in south-east Asia (adapted from Fitzherbert <i>et al.</i> , 2008)	32
2.7	Diagram of the ITS region including the intergenic transcribed spacers (ITS1 and ITS2) and the 5.8S rRNA gene (adapted from http://www.biology.duke.edu/fungi/mycolab/primers.htm , 24 May 2010)	39
2.8	Structure of compactin ($R_1=H$, $R_2=H$) and pravastatin ($R_1=OH$, $R_2=H$) (adapted from Chakravarti and Sahai, 2004)	41
2.9	Schematic diagram of biosynthetic pathways for lovastatin and compactin (modified from Manzoni and Rollini, 2002; Endo, 2004)	42
2.10	Compactin biosynthetic gene cluster (adapted from Abe <i>et al.</i> , 2002)	43
2.11	Pravastatin production using a two-step fermentation process (adapted from Ykema <i>et al.</i> , 1999)	66
4.1	Sampling locations of mangrove soils in Pahang State (Malaysia) (adapted from www.malaxi.com)	88

4.2	Fungal isolates from oil palm plantation soils grown on PDA (COC = colony obverse color, CRC = colony reverse color)	92
4.3	Fungal isolates from mangrove soils grown on PDA (COC = colony obverse color, CRC = colony reverse color)	96
4.4	Colony morphology of fungal isolates from oil palm plantation soils on CZ and CYA25 agar media (COC = colony obverse color, CRC = colony reverse color)	99
4.5	Colony morphology of fungal isolates from oil palm plantation soils on MEA and YES agar media (COC = colony obverse color, CRC = colony reverse color)	105
4.6	Colony morphology of fungal isolates from mangrove soils on CZ and CYA25 agar media (COC = colony obverse color, CRC = colony reverse color)	111
4.7	Colony morphology of fungal isolates from mangrove soils on MEA and YES agar media (COC = colony obverse color, CRC = colony reverse color)	115
4.8	Light micrographs of fungal isolates from oil palm plantation soils (image is magnified 400×)	120
4.9	Light micrographs of fungal isolates from mangrove soils (image is magnified 400×)	126
6.1	Growth of <i>Penicillium</i> isolates on CYA25 and CYAS agar media (COC = colony obverse color; CRC = colony reverse color)	158
6.2	Growth of <i>Penicillium</i> isolates on NO ₂ agar	159
6.3	Growth, acid and base production of <i>Penicillium</i> isolates on CREA agar	160
6.4	Ehrlich results on CYA25 after 7d at 25°C	162
6.5	Dendrogram based on the Euclidean distance and centroid clustering method showing the relationship between clusters of <i>Penicillium</i> isolates	164

6.6	HPLC analysis of plug extract from <i>Penicillium</i> sp. ESF2M isolate grown on YES	165
6.7	HPLC analysis of plug extract from <i>Penicillium</i> sp. ESF19M isolate grown on YES	166
6.8	HPLC analysis of plug extract from <i>Penicillium</i> sp. ESF20P isolate grown on YES	167
6.9	HPLC analysis of plug extract from <i>Penicillium</i> sp. ESF21P isolate grown on YES	168
6.10	HPLC analysis of plug extract from <i>Penicillium</i> sp. ESF26P isolate grown on YES	169
6.11	Dendrogram of the <i>Penicillium</i> isolates studied using Euclidean distance and centroid clustering method	170
6.12	Macroscopic and microscopic photographs of the proposed species taken directly from the Biolog FF Database	173
6.13	Genomic DNA of the fungal isolates separated on a 1% agarose gel. Lane L is the VC 100-bp plus DNA ladder (Vivantis Technologies, U.S.A.) used as size marker. Lanes 1-5: 1) <i>Penicillium</i> sp. ESF2M; 2) <i>Penicillium</i> sp. ESF19M; 3) <i>Penicillium</i> sp. ESF20P; 4) <i>Penicillium</i> sp. ESF21P; 5) <i>Penicillium</i> sp. ESF26P.	174
6.14	PCR products separated on a 1% agarose gel. Lane L is the VC 100-bp plus DNA ladder (Vivantis Technologies, U.S.A.) used as size marker. Lanes 1-5: 1) <i>Penicillium</i> sp. ESF2M; 2) <i>Penicillium</i> sp. ESF19M; 3) <i>Penicillium</i> sp. ESF20P; 4) <i>Penicillium</i> sp. ESF21P; 5) <i>Penicillium</i> sp. ESF26P.	175
6.15	Aligned rDNA sequences of the ITS1-5.8S-ITS2 regions of <i>Penicillium</i> sp. ESF2M and <i>Penicillium citrinum</i> KACC43900 (EU821333.1). Gaps indicate differences between the sequences of <i>Penicillium</i> sp. ESF2M (query sequence) and <i>Penicillium citrinum</i> KACC43900 (subject sequence). The numbers on the right margin show the actual number of bases in each line.	176

6.16	Aligned rDNA sequences of the ITS1-5.8S-ITS2 regions of <i>Penicillium</i> sp. ESF19M and <i>Penicillium citrinum</i> KACC43900 (EU821333.1). Gaps indicate differences between the sequences of <i>Penicillium</i> sp. ESF19M (query sequence) and <i>Penicillium citrinum</i> KACC43900 (subject sequence). The numbers on the right margin show the actual number of bases in each line.	177
6.17	Aligned rDNA sequences of the ITS1-5.8S-ITS2 regions of <i>Penicillium</i> sp. ESF20P and <i>P. janthinellum</i> CBS:340.48 (GU981585.1). Gaps indicate differences between the query (<i>Penicillium</i> sp. ESF20P) and subject (<i>P. janthinellum</i> CBS:340.48) sequences. The numbers on the right margin show the actual number of bases in each line.	178
6.18	Aligned rDNA sequences of the ITS1-5.8S-ITS2 regions of <i>Penicillium</i> sp. ESF26P and <i>P. janthinellum</i> P1 (JQ727998.1). Gaps indicate differences between the query (<i>Penicillium</i> sp. ESF26P) and subject (<i>P. janthinellum</i> P1) sequences. The numbers on the right margin show the actual number of bases in each line.	179
6.19	Aligned rDNA sequences of the ITS1-5.8S-ITS2 regions of <i>Penicillium</i> sp. ESF21P and <i>E. brefeldianum</i> 3352 (FJ527878.1). Gaps indicate differences between the query (<i>Penicillium</i> sp. ESF21P) and subject (<i>E. brefeldianum</i> 3352) sequences. The numbers on the right margin show the actual number of bases in each line.	180
6.20	Taxonomic classification of the identified <i>Penicillium</i> species (Pitt, 1979)	183
6.21	Taxonomic classification of <i>E. brefeldianum</i> ESF21P (Pitt, 1979)	183
7.1	Normal probability plot of the studentized residuals	220
7.2	Outlier T versus run order	221
7.3	Box-Cox plot for power transformations	222
7.4	Ramps of pre-optimization from Design-Expert 6.0.8 software	223

7.5	Normal probability plot of the studentized residuals after square root transformation	227
7.6	Outlier T versus run order after square root transformation	228
7.7	Box-Cox plot for power transformations	229
7.8	Effect of interaction between slant age (SA) and fermentation time (FT) on pravastatin production	233
7.9	Effect of interaction between slant age (SA) and initial pH of the medium on pravastatin production	234
7.10	Effect of interaction between slant age (SA) and agitation rate (AR) on pravastatin production	235
7.11	Effect of interaction between fermentation time (FT) and initial pH of the medium on pravastatin production	236
7.12	Effect of interaction between fermentation time (FT) and agitation rate (AR) on pravastatin production	237
7.13	Effect of interaction between initial pH of the medium and agitation rate (AR) on pravastatin production	238
7.14	Effects of the different agitation speeds on morphological characteristics of <i>E. brefeldianum</i> ESF21P in submerged fermentation: (a) filamentous form at 180 rpm; (b) pellet form at 210 rpm	238
7.15	Perturbation plot showing influence of significant variables on pravastatin production by <i>E. brefeldianum</i> ESF21P (SA=slant age; FT=fermentation time; pH=initial pH of the medium; AR=agitation rate)	239
7.16	Ramps of the optimization experiment using CCD (SA=slant age; FT=fermentation time; pH=initial pH of the medium; AR=agitation rate)	241
8.1	Growth profile of <i>E. brefeldianum</i> ESF21P in shake flask fermentation	265

8.2	Substrate uptake during shake flask fermentation of <i>E. brefeldianum</i> ESF21P	266
8.3	Pravastatin production by <i>E. brefeldianum</i> ESF21P in shake flask fermentation	267
8.4	The maximum dry cell weight (X_{\max}) obtained during shake flask fermentation of <i>E. brefeldianum</i> ESF21P	268

LIST OF ABBREVIATIONS

ANOVA	Analysis of variance
AR	Agitation rate
bp	Base pair
CCD	Central composite design
CD	Colony diameter
cm	Centimeter
CoA	Coenzyme A
COC	Colony obverse color
CRC	Colony reverse color
CREA agar	Creatine-Sucrose Agar
$\text{CuSO}_4 \cdot 5\text{H}_2\text{O}$	Copper (II) sulphate pentahydrate
CYA	Czapek Yeast Autolysate Agar
CYAS	Czapek Yeast Autolysate with 5% NaCl
CZ	Czapek-Dox Agar
d	Day
2D	Two-dimensional
3D	Three-dimensional
DAD	Diode array detection
DCW	Dry cell weight
DIST	Distance

DMSO	Dimethylsulfoxide
DNA	Deoxyribonucleic acid
$\text{FeSO}_4 \cdot 7\text{H}_2\text{O}$	Ferrous sulfate heptahydrate
FF	Filamentous fungi
FFD	Fractional factorial design
2FI	2- Factor interaction model
3FI	3-Factor interaction model
FT	Fermentation time
g	Gram
GC	Gas chromatography
Glu	Glucose
Gly	Glycerol
h	Hour
ha	Hectare
HCl	Hydrochloric acid
HDL	High-density lipoprotein
HMG	3-Hydroxy-3-methylglutaryl
HMG-CoA	3-Hydroxy-3-methylglutaryl-coenzyme A
HPLC	High- performance liquid chromatography
H_3PO_4	Phosphoric acid
ID	Internal diameter
IFO	Institute for Fermentation

ITS 1	Internal transcribed spacer 1
ITS 2	Internal transcribed spacer 2
IUPAC	International Union of Pure and Applied Chemistry
IV	Inoculum volume
kb	Kilobase
KCl	Potassium chloride
kg	Kilogram
KH_2PO_4	Potassium dihydrogen phosphate
$\text{K}_2\text{HPO}_4 \cdot 3\text{H}_2\text{O}$	Dipotassium hydrogen phosphate
L	Litre
LC	Liquid chromatography
LDL	Low-density lipoprotein
m	Metre
mAU	Milliabsorbance unit
MEA	Malt Extract Agar
mg	Milligram
MgCl_2	Magnesium chloride
$\text{MgSO}_4 \cdot 7\text{H}_2\text{O}$	Magnesium sulfate heptahydrate
min	Minute
mL	Millilitre
ML-236B	Mevastatin
mm	Millimetre

MS	Mass spectrometry
N	Normality
NaCl	Sodium chloride
NADPH	Nicotinamide adenine dinucleotide phosphate
NaNO ₂	Sodium nitrite
NaNO ₃	Sodium nitrate
NaOH	Sodium hydroxide
Na ₂ SO ₄	Sodium sulphate
ND	Not detected
ng	Nanogram
NHMS	National Health and Morbidity Survey
nm	Nanometre
NO ₂ agar	Nitrite-Sucrose Agar
NTG	<i>N</i> -methyl- <i>N'</i> -nitro- <i>N</i> -nitrosoguanidine
OD	Optical density
PCR	Polymerase Chain Reaction
PDA	Potato-Dextrose Agar
pH	Hydrogen ion concentration
PROB	Probability
PTFE	Polytetrafluoroethylene
rDNA	Ribosomal deoxyribonucleic acid
RID	Refractive index detector