## Utilization of Ion-Exclusion Chromatography for Water Quality Monitoring in a Suburban River in Jakarta, Indonesia

Daisuke Kozaki <sup>ab</sup>, Takashi Ozaki<sup>c</sup>, Nobutake Nakatani<sup>d</sup>, Masanobu Mori<sup>e</sup> and Kazuhiko Tanaka<sup>c</sup> <sup>a</sup>Faculty of Industrial Sciences and Technology, University Malaysia Pahang, Lebuhraya Tun Razak, 26300 Gambang Kuantan, Pahang Darul Makmur, Malaysia

<sup>b</sup>Centre for Earth Resources Research & Management, University Malaysia Pahang, University Malaysia Pahang, Lebuhraya Tun Razak, 26300 Gambang Kuantan, Pahang Darul Makmur, Malaysia <sup>c</sup>Graduate School for International Development and Cooperation, Hiroshima University, 1-5-1 Kagamiyama, Higashi-hiroshima 739-8529, Japan

<sup>d</sup>Department of Environmental and Symbiotic Sciences, College of Agriculture, Food and Environmental Science, Rakuno Gakuen University, 582 Bunkyodai-Midorimachi, Ebetsu, Hokkaido 069–8501, Japan

<sup>e</sup>Faculty of Science and Technology, Gunma University, 1-5-1 Tenjin, Kiryu, Gunma 376–8515, Japan

## ABSTRACT

We evaluated the use of ion-exclusion chromatographic systems for analyzing the behavior of inorganic ions (e.g., bicarbonate, sulfate, chloride, nitrate, phosphate, dissolved silicate, sodium, ammonium, potassium, magnesium, and calcium ions) in a suburban river located in Jakarta, Indonesia. Carbonate, phosphate, and silicate ion concentrations were determined using ionexclusion chromatography (IEC) on a weakly acidic cation-exchange resin column (WCX) in the H<sup>+</sup>form with water eluent. Other ions were identified by ion-exclusion/cation-exchange chromatography (IEC/CEC) on a WCX column with tartaric acid eluent. The use of IEC systems for water quality monitoring was advantageous for the following reasons: (1) the concentrations of analyte ions, except  $NO_3^-$  and silicate ions, increased from upstream to downstream; and (2) the speciation of inorganic nitrogen ions could be analyzed by single injection into the IEC/CEC. The IEC approach provided beneficial information for the construction of sewage treatment facilities in our study area. Results showed that (1) the analyte concentrations for samples obtained in the downstream area were higher than those in the upstream area owing to contamination by domestic sewage; (2) the concentrations of  $NO_3^-$  and  $NH_4^+$  correlated with the concentration of dissolved oxygen; and (3) bicarbonate concentrations increased downstream, likely due to respiration of bacteria and dissolution of concrete under low-oxygen conditions.

**KEYWORDS**: ion-exclusion chromatography; inorganic ions; microbial reaction; suburban river; water in developing countries

## DOI: 10.3390/w6071945